Bioavailability of Oral Pyridostigmine and Inhibition of Red Blood Cell Acetylcholinesterase by Oral and Intravenous Pyridostigmine

TASK ORDER #2

FINAL REPORT

David M. Kornhauser, M.D.
Brent G. Petty, M.D.
Paul S. Lietman, M.D., Ph.D.

22 March 1989

Supported by:

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Fort Detrick, Maryland 21701-5012

Contract No. DAMD17-85-C-5133

Division of Clinical Pharmacology
The Johns Hopkins University School of Medicine
600 N. Wolfe Street
Baltimore, Maryland 21205

Approved for public release; distribution is unlimited.

The views, opinions, and/or findings contained in this repent are those of the authors and should not be construed as an official Department of Army position, policy, or decision, unless so designated by other documentation.

20100915142

19. ABSTRACT (continued)

The mean bioavailability of oral pyridostigmine syrup was 29.2%. Considerable interindividual variability was noted, with the range of bioavailability being 14.7% to 51.1%. The mean rate constant of absorption was 0.373/hr with a range of 0.149 to 1.441/hr. These absorption rate constants are equivalent to a mean absorption half-time of 1.06 hours with a range of 0.48 to 4.65 hours.

The pharmacokinetics of both oral and intravenous pyridostigmine have been defined using a two-compartment model. Considerable interindividual variability in elimination of pyridostigmine was found here as well. The mean total clearance was 779 ml/min with a range of 382 ml/min to 1511 ml/min, and the mean beta half-life was 0.8 hours with a range of 0.36 to 3.2 hours.

The subjects were monitored for toxicity with clinical laboratory tests of hematology and chemistry variables, electrocardiograms, vital signs, and coordination testing. All of the subjects tolerated pyridostigmine well, with no adverse symptoms. One subject developed a significant increase in serum creatine kinase and 2 subjects had trivial elevations in liver enzymes with no symptoms. No significant changes occurred in electrocardiograms, vital signs, coordination, or other clinical laboratory tests.

The relationship between pyridostigmine plasma levels and erythrocyte acetylcholinesterase inhibition was also defined. The erythrocyte acetylcholinesterase inhibition was delayed in both onset and dissipation compared to plasma pyridostigmine levels after both oral and intravenous pyridostigmine administration. The delay in effect was modelled by assuming the presence of an effect compartment and an E_{max} concentration-effect relationship. According to this model, the mean rate constant of elimination from the effect compartment was 3.42/hr with a range of 1.22 to 5.76/hr. Thus, the mean half-time of drug effect at a constant plasma concentration was 0.20 hours (12 minutes) with a range of 0.12 to 0.57 hours. The mean plasma concentration of pyridostigmine required to produce 50% enzyme inhibition at steady state was estimated to be 29.5 ng/ml with a range of 17.3 to 41.1 ng/ml.

The time delay in enzyme inhibition could also be accounted for by a model which assumed that carbamylation of the acetylcholinesterase enzyme inactivated it and that hydrolysis of the inactivated enzyme restored activity. The mean estimate of the rate constant of inactivation was 0.087 ml/ng/hr with a range of 0.058 to 0.164 ml/ng/hr, while the mean estimate for the rate constant of reactivation was 2.736/hr with a range of 1.652 to 4.107/hr. These estimates are similar to estimates of the rates of the inactivation and reactivation of purified acetylcholinesterase previously measured in vitro by others.

Considerable intersubject variability in the drug's effect occurred after oral dosing. The peak erythrocyte acetylcholinesterase inhibition varied from 20 to 39% of the baseline enzyme activity, and the duration of at least 20% inhibition varied from 0.33 to 5.0 hours. The variation appeared to be due to interindividual differences in the amount of absorption of the drug, in the rate of elimination of the drug, and in the sensitivity of the acetylcholinesterase to inhibition by pyridostigmine.

SECURITY CLASSIFICATION OF THIS PAGE	DOCUMENTATIO	N PAGE			rm Approved
			OA	AB No. 0704-0188	
ia. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE	MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION	/AVAILABILITY OF	REPORT	
2b. DECLASSIFICATION / DOWNGRADING SCHE	DITE		ed for Publi		
20. DECEASSIFICATION/DOWNGRADING SCHE	DOLE	Distribution Unlimited			
4. PERFORMING ORGANIZATION REPORT NUM	ABER(S)	5. MONITORING	ORGANIZATION RE	PORT NUMBER	(S)
6a. NAME OF PERFORMING ORGANIZATION Div. of Clinical Pharmacology	6b. OFFICE SYMBOL (If applicable)		ONITORING ORGAN Army Medical		
The Johns Hopkins U. Sch. of M			pment Center		
6c. ADDRESS (City, State, and ZIP Code)			y, State, and ZIP C		
600 N. Wolfe Street - Osler Baltimore, Maryland 21205	527				
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	Bb. OFFICE SYMBOL (If applicable)	9. PROCUREMENT	INSTRUMENT IDE	NTIFICATION N	IUMBER
USAMRDC		DAMD17-	85-C-5133		
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF F	UNDING NUMBERS	\$	
Fort Detrick Frederick, Maryland 21701-	5012	PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
riederick, haryland 21701-	3012	NA	NA	NA	NA
11. TITLE (Include Security Classification)					
Phase I Clinical Pharmacolo	gy Studies				
12. PERSONAL AUTHOR(S) David M. Kornhauser, M.D.;					
	COVERED 0ct21,8500Mar22,89	14. DATE OF REPO		-	E COUNT 505
16. SUPPLEMENTARY NOTATION	00000 Mar 22.09	. 1969 Marci	11 22		505
Subtitle: Bioavailability	of Oral Pyridosti	gmine and In	nibition of	Red Blood	Cell
Acetylcholineste	rase by Oral and	Intravenous	Pyridostiemi	ine	
17. COSATI CODES FIELD GROUP SUB-GROUP	18. SUBJECT TERMS (C	continue on reverse mine, Acetyle			
FIELD GROOF 30B-GROOF		netics, Human			
19. ABSTRACT (Continue on reverse if necessary Pyridostigmine bromic			to reduce t	the mortal	ity from
organophosphate exposure ir	animal studies.	This human	study was pe	erformed (1) to
determine the bioavailabili					
kinetics and safety of pyri the relationship between py					
inhibition after the oral a					normescerase
The first six subject					а
dose-ranging phase) in an e	ffort to approach	, but not ex	ceed, 40% in	nhibition	of
erythrocyte acetylcholinest					
compromise between effective and toxicity from the drug					
40% erythrocyte acetylcholi					
received oral and intravend	ous pyridostigmine	to address	the objectiv	ves mentio	ned above.
20. DISTRIBUTION / AVAILABILITY OF ABSTRAC		21. ABSTRACT SEC	CURITY CLASSIFICA	ATION	Cont
22a. NAME OF RESPONSIBLE INDIVIDUAL	The Colonia	22b. TELEPHONE (Include Area Code) 22c. OFFICE !	SYMBOL
Schuster, Brian, Col., MC		(301) 427-	-5346	SCRD-I	JWWU

Bioavailability of Oral Pyridostigmine and Inhibition of Red Blood Cell Acetylcholinesterase by Oral and Intravenous Pyridostigmine

TASK ORDER #2

FINAL REPORT

David M. Kornhauser, M.D.
Brent G. Petty, M.D.
Paul S. Lietman, M.D., Ph.D.

22 March 1989

Supported by:

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND Fort Detrick, Maryland 21701-5012

Contract No. DAMD17-85-C-5133

Division of Clinical Pharmacology
The Johns Hopkins University School of Medicine
600 N. Wolfe Street
Baltimore, Maryland 21205

Approved for public release; distribution is unlimited.

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as an official Department of Army position, policy, or decision, unless so designated by other documentation.

SUMMARY

Pyridostigmine bromide pretreatment has been shown to reduce the mortality from organophosphate exposure in animal studies. This human study was performed (1) to determine the bioavailability of oral pyridostigmine syrup, (2) to examine the pharmacokinetics and safety of pyridostigmine given orally and intravenously, and (3) to determine the relationship between pyridostigmine plasma levels and erythrocyte acetylcholinesterase inhibition after the oral and intravenous administration of pyridostigmine.

The first six subjects received oral and intravenous pyridostigmine (in a dose-ranging phase) in an effort to approach, but not exceed, 40% inhibition of erythrocyte acetylcholinesterase. This target level of inhibition was chosen as a compromise between effective protection against organophosphate poisoning on the one hand and toxicity from the drug on the other. After the proper doses to achieve approximately 40% erythrocyte acetylcholinesterase inhibition were determined, 18 additional subjects received oral and intravenous pyridostigmine to address the objectives mentioned above.

The mean bioavailability of oral pyridostigmine syrup was 29.2%. Considerable interindividual variability was noted, with the range of bioavailability being 14.7% to 51.1%. The mean rate constant of absorption was 0.373/hr with a range of 0.149 to 1.441/hr. These absorption rate constants are equivalent to a mean absorption half-time of 1.06 hours with a range of 0.48 to

4.65 hours.

The pharmacokinetics of both oral and intravenous pyridostigmine have been defined using a two-compartment model.

Considerable interindividual variability in elimination of pyridostigmine was found here as well. The mean total clearance was 779 ml/min with a range of 382 ml/min to 1511 ml/min, and the mean beta half-life was 0.8 hours with a range of 0.36 to 3.2 hours.

The subjects were monitored for toxicity with clinical laboratory tests of hematology and chemistry variables, electrocardiograms, vital signs, and coordination testing. All of the subjects tolerated pyridostigmine well, with no adverse symptoms. One subject developed a significant increase in serum creatine kinase and 2 subjects had trivial elevations in liver enzymes with no symptoms. No significant changes occurred in electrocardiograms, vital signs, coordination, or other clinical laboratory tests.

The relationship between pyridostigmine plasma levels and erythrocyte acetylcholinesterase inhibition was also defined. The erythrocyte acetylcholinesterase inhibition was delayed in both onset and dissipation compared to plasma pyridostigmine levels after both oral and intravenous pyridostigmine administration. The delay in effect was modelled by assuming the presence of an effect compartment and an E_{max} concentration-effect relationship. According to this model, the mean rate constant of elimination from the effect compartment was 3.42/hr with a range of 1.22 to 5.76/hr. Thus, the mean half-time of drug effect at a

constant plasma concentration was 0.20 hours (12 minutes) with a range of 0.12 to 0.57 hours. The mean plasma concentration of pyridostigmine required to produce 50% enzyme inhibition at steady state was estimated to be 29.5 ng/ml with a range of 17.3 to 41.1 ng/ml.

The time delay in enzyme inhibition could also be accounted for by a model which assumed that carbamylation of the acetyl-cholinesterase enzyme inactivated it and that hydrolysis of the inactivated enzyme restored activity. The mean estimate of the rate constant of inactivation was 0.087 ml/ng/hr with a range of 0.058 to 0.164 ml/ng/hr, while the mean estimate for the rate constant of reactivation was 2.736/hr with a range of 1.652 to 4.107/hr. These estimates are similar to estimates of the rates of the inactivation and reactivation of purified acetylcholinesterase previously measured in vitro by others.

Considerable intersubject variability in the drug's effect occurred after oral dosing. The peak erythrocyte acetylcholinesterase inhibition varied from 20 to 39% of the baseline enzyme activity, and the duration of at least 20% inhibition varied from 0.33 to 5.0 hours. The variation appeared to be due to interindividual differences in the amount of absorption of the drug, in the rate of elimination of the drug, and in the sensitivity of the acetylcholinesterase to inhibition by pyridostigmine.

FOREWORD

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

For the protection of human subjects, the investigators have adhered to the policies of applicable Federal Law 45 CFR 46.

TABLE OF CONTENTS

I.	REP	ORT		Page
	1.	INTRODUCTION		
	2.	MATE	RIALS AND METHODS	12
		2.1	Pyridostigmine	12
		2.2	Subjects 2.2.1 Inclusion Criteria 2.2.2 Exclusion Criteria 2.2.3 Recruitment 2.2.4 Informed Consent 2.2.5 Compensation 2.2.6 Liability	13 13 14 14 14 15
		2.3	Experimental Protocol 2.3.1 Objectives 2.3.2 Design	15 15 16
		2.4	Clinical Laboratory Examinations 2.4.1 Hematology 2.4.2 Chemistry 2.4.3 Electrocardiography 2.4.4 Urine Analysis 2.4.5 Chest x-ray	17 18 18 18 18
		2.5	Pyridostigmine and Erythrocyte Acetylcholinesterase Determinations 2.5.1 Pyridostigmine Analysis 2.5.2 Erythrocyte Acetylcholinesterase Determinations	19 19 20
		2.6	Pharmacokinetic and Pharmacodynamic Analyses 2.6.1 Pharmacokinetic Analyses 2.6.2 Pharmacodynamic Modeling	20 20 22
		2.7	Statistical Methods	26
	3.	RESU	LTS AND DISCUSSION	26
		3.1	Amendments and Compliance 3.1.1 Amendments 3.1.2 Compliance	26 26 26
		3.2	Description of Population of Subjects	29
		3.3	Clinical Results 3.3.1 Symptomatic 3.3.2 Pulse 3.3.3 Coordination Testing	30 30 30 31

		3.3.5	Hand Grip Laboratory 3.3.5.1 Liver Function Tests 3.3.5.2 Creatine Kinase 3.3.5.3 Other Clinical Chemistry Tests 3.3.5.4 Hematological Testing 3.3.5.5 Electrocardiograms Clinical Conclusions	31 31 31 33 34 34 35
	3.4	3.4.1	acokinetics and Pharmacodynamics Pharmacokinetics Pharmacodynamics	3 5 3 5 4 5
	4. CON	CLUSION	NS .	54
II.	REFEREN	CES		57
T T T	TABLES	AND ETC	TUDEC	()
III.	Table		Sample Times Omitted From Pharmacokinetic Curve Fitting	63 63
	Table	2	Equations to which Plasma Pyridostigmine Concentrations were Fit	64
	Table	3	Vital Statistics of Subjects	65
	Table		Sampling Times with Intravenous Pyrido- stigmine	66
	Table	5	Pyridostigmine Concentrations with	67
			Intravenous Pyridostigmine	
	Table		Sampling Times with Oral Pyridostigmine	68
	Table	7	Pyridostigmine Concentrations with Oral Pyridostigmine	69
	Table	8	"Goodness of Fit" Indicators for Pharmaco- kinetic Curve Fitting	70
	Table	9	Estimates of Pharmacokinetic Microconstants	71
	Table		Estimates of Pharmacokinetic Macroconstants	72
	Table		Estimates of Area Under the Concentration- Time Curve and Clearance of Pyridostigmine	73
	Table	12	Urinary Excretion of Pyridostigmine over Time	74
	Table		Total Urinary Pyridostigmine Recovery	75
	Table		Pyridostigmine Renal Clearance and Bioavail-	76
			ability using Urinary Excretion Data	
	Table	15	Acetylcholinesterase Inhibition with Intravenous Pyridostigmine	77
	Table	16	Acetylcholinesterase Inhibition with Oral Pyridostigmine	78
	Table	17	Duration of Acetylcholinesterase Inhibition with Intravenous Pyridostigmine	79
	Table	18	Duration of Acetylcholinesterase Inhibition with Oral Pyridostigmine	80
	Table	19	Maximal Acetylcholinesterase Inhibition with	81
	Table	20	Intravenous Pyridostigmine Maximal Acetylcholinesterase Inhibition with	82
	Table	21	Oral Pyridostigmine Area within the Hysteresis Loop	83

	lable 22	"Goodness of Fit" indicators for the Effect	84
	T 1 1 00	Model	
	Table 23	Estimates of K _{E0} and IC ₅₀	85
	Table 24	"Goodness of Fit" Indicators for the	86
		Biochemical Model	
	Table 25	Estimates of Rate Constants for Acetyl-	87
		cholinesterase Inactivation and Reactivation	
	Figure 1a	Schematics for Effect Compartment and	88
	and 1b	Biochemical Models	
	Figure 2	Average Pyridostigmine Plasma Concentrations	89
	-	For Each Sampling Time	
	Figure 2a	Pyridostigmine Concentrations with Intravenous	90
	3	Dosing of Subjects 7-12	
	Figure 2b	Pyridostigmine Concentrations with Oral	91
		Dosing of Subjects 7-12	
	Figure 2c	Pyridostigmine Concentrations with Intravenous	92
	119410 20	Dosing of Subjects 13-18	22
	Figure 2d	Pyridostigmine Concentrations with Oral	93
	rigare za	Dosing of Subjects 13-18))
	Figure 2e	Pyridostigmine Concentrations with Intravenous	94
	rigure ze	Dosing of Subjects 19-24	94
	Figure 26		0 =
	Figure 2f	Pyridostigmine Concentrations with Oral	95
	7.	Dosing of Subjects 19-24	
	Figure 3	Pyridostigmine Absorbed as a Function of Oral	96
		Dose	
	Figure 4	Average Acetylcholinesterase Inhibition for	97
		Each Sampling Time	
	Figure 4a	Acetylcholinesterase Inhibition with	98
		Intravenous Dosing of Subjects 7-12	
	Figure 4b	Acetylcholinesterase Inhibition with Oral	99
		Dosing of Subjects 7-12	
	Figure 4c	Acetylcholinesterase Inhibition with	100
	-	Intravenous Dosing of Subjects 13-18	
	Figure 4d	Acetylcholinesterase Inhibition with Oral	101
	3	Dosing of Subjects 13-18	
	Figure 4e	Acetylcholinesterase Inhibition with	102
	9	Intravenous Dosing of Subjects 19-24	
	Figure 4f	Acetylcholinesterase Inhibition with Oral	103
	rigure 41	Dosing of Subjects 19-24	103
	Figure 5a	Acetylcholinesterase Inhibition Plotted	104
	rigure sa		104
		Against Pyridostigmine Concentration with Intravenous Dose	
	71		205
	Figure 5b	Acetylcholinesterase Inhibition Plotted	105
		Against Pyridostigmine Concentration with	
		Oral Dose	
IV.	APPENDICES		106
		on of Pyridostigmine Dose Administered	106
		Timetable and Study Flow Charts	109
	C. Clinical La	boratory Normal Values	119
		ythrocyte Acetylcholinesterase at The Johns	121
		versity Division of Clinical Pharmacology	
	E. Original Co		128
	F. Amended Con		129

G.	Publication Supported by this Contract	131
Н.	Personnel Receiving Contract Support	132
I.	Distribution List	133
J.	Case Report Forms	134

1. INTRODUCTION

Studies in animals have indicated that carbamate acetylcholinesterase inhibitors have protective effects against organophosphate poisoning. Pretreatment of animals with carbamates prior to nerve agent exposure has improved survival and effectively increased the LD50 of nerve gas agents (1). Experimental carbamate pretreatment has been most effective when used in conjunction with atropine and oximes, both given following exposure (1,2). Of the carbamates studied, pyridostigmine has been found to be a useful agent with a duration of protective action of about two hours. The putative mechanism of action of carbamates is thought to be carbamylation of a fraction of the tissue acetylcholinesterase, thereby protecting the enzyme from irreversible inhibition by the organophosphate (3). relatively rapid hydrolysis of the inactivated carbamylated enzyme has two sequelae. First, following a single dose of pyridostigmine, there is only a relatively short period of protection since pyridostigmine is rapidly removed from the body and the inactivated enzyme is rapidly reactivated by hydrolysis. Thus, if long term protection from possible organophosphate exposure is required, then repeated administration of pyridostigmine is required. Second (and on the positive side), if exposure to an organophosphate were to occur, the relatively short duration of action of pyridostigmine allows for reactivation of carbamylated acetylcholinesterase in a short

period of time, possibly minimizing the time period required for intensive therapy of the effects of cholinergic excess.

The dose of pyridostigmine and degree of acetylcholinesterase inhibition which provide adequate protection against organophosphate poisoning while at the same time remaining free of acceptable toxicity in man are not known. Studies in rats have indicated that inhibition of twenty-five percent of blood acetylcholinesterase does not affect muscle twitch tension, even after several days of therapy. On the other hand, abnormalities in twitch tension occurred in animals treated with enough pyridostigmine to produce sixty-eight percent inhibition of the enzyme (4). In addition, there are ultrastructural changes at the neuromuscular junction associated with this greater degree of inhibition of acetylcholinesterase (5). The changes appear to be dose related and, at least at low levels of enzyme inhibition, largely reversible.

The optimal use of pyridostigmine in man as prophylaxis for organophosphate poisoning thus presupposes that pyridostigmine is present in the body at the time of exposure and that a level of drug sufficient to provide protection while having few toxic effects can be achieved. With the fixed positive charge on the pyridostigmine molecule likely to make transdermal delivery difficult, multiple oral doses of drug appear to be, at least for the short term, the most feasible route of administration for prolonged protection. This study was designed to determine the amount of pyridostigmine required to produce modest levels of

erythrocyte acetylcholinesterase inhibition (approximately 40%) and to determine the bioavailability of oral pyridostigmine administered as the syrup that is currently commercially available. Once this fundamental information is obtained, the efficacy and systemic availability of other preparations can be evaluated and compared to the syrup as a standard.

2. MATERIALS AND METHODS

2.1 Pyridostigmine

Pyridostigmine was administered as the bromide salt. Mestinon^R, Roche Laboratory's brand of pyridostigmine bromide syrup, lot #0204, was provided by the U.S. Army for oral dosing. The oral doses were administered by expelling the syrup from a syringe directly onto the subject's tongue. The subject swallowed the dose immediately. Blood samples were obtained by means of an indwelling catheter in the arm which was kept patent by a dilute heparin solution. Organon brand pyridostigmine bromide, lot #485460A, was given intravenously to the first six subjects. A single lot, Roche lot #102, U.S. Army lot #BL10992, of injectable Mestinon was used for the intravenous infusions in subjects 07 through 24. Drug was diluted to 20 ml with normal saline and infused over 30 minutes with an infusion pump. Blood samples were drawn from an indwelling catheter in the contralateral arm. The volume of pyridostigmine administered was determined by weighing the syringe full of medication immediately prior to dosing and subtracting the weight of the empty syringe after dosing, then dividing the difference by the specific gravity of the solution. Oral and intravenous dosing solutions were assayed for pyridostigmine concentration. The administered dose was then calculated from the product of the volume administered times the assayed concentrations (Appendix A).

2.2 Subjects

Healthy men who were able to give written informed consent were eligible to volunteer for the study. The study was approved by the Joint Committee on Clinical Investigation of The Johns Hopkins Medical Institutions and the Human Subjects Research Review Board of the U.S. Army.

2.2.1 Inclusion Criteria

To participate in the study the volunteer had to be male, between 18 and 35 years of age, and within 10% of his ideal body weight. A detailed health history and physical examination were performed by a physician. Serum chemistries, hematology and urine analysis had to be within normal ranges (except creatine kinase), as defined by The Johns Hopkins Hospital Department of Laboratory Medicine. The creatine kinase (CK) could be above the "normal range" and not exclude the subject because of the frequent finding of elevated CK in healthy subjects who are especially active physically (6-13). A chest x-ray within 6 months of entry and an electrocardiogram had to be

normal.

2.2.2 Exclusion Criteria

Women were excluded from this study. Men were excluded if they did not meet the criteria listed above (2.2.1) or if they had a known or suspected allergy to pyridostigmine or related drugs. Once accepted as candidates for the study, subjects were not permitted to take any medication for one week prior to admission to the study.

2.2.3 Recruitment

Advertisements were placed in the help wanted classified sections of metropolitan Baltimore newspapers. A special telephone line was dedicated to volunteer recruitment. Interested candidates were screened on the telephone by a research nurse who described the details of the study, took a brief history and scheduled the appropriate screening examinations.

2.2.4 Informed Consent

Written informed consent was obtained from each participant upon admission to the Clinical Research Unit. The consent document described in detail the purpose of the study, the research protocol, and the potential risks.

2.2.5 Compensation

A payment schedule was designed to compensate volunteers for their participation. Each volunteer was compensated as follows:

Screening phase No compensation In-hospital phase \$375.00

Total per subject for entire study \$375.00

For the entire study, \$9000.00 was distributed to 24 volunteers.

2.2.6 Liability

Liability coverage for unexpected toxicity was provided by the U.S. Army, and for malpractice by the Johns Hopkins Medical Institutions.

2.3 Experimental Protocol

2.3.1 Objectives

There were three objectives of the study. The primary objective was to determine the absolute bioavailability of pyridostigmine bromide when given as the oral syrup. However, since for safety reasons erythrocyte acetylcholinesterase inhibition could not exceed 40%, and because the assay of pyridostigmine base was possible only at concentrations greater than 1.5 ng/ml, it was first necessary to determine the oral and intravenous doses of pyridostigmine which would produce plasma

levels of drug high enough to measure but which inhibited the enzyme less than 40%. Once the appropriate doses were found, a formal bioavailability study with eighteen subjects was conducted. The other two objectives of the study were to examine the pharmacokinetics and safety of pyridostigmine given orally and intravenously, and to determine the relationship between pyridostigmine plasma levels and erythrocyte acetylcholinesterase inhibition after the oral and intravenous administration of pyridostigmine.

2.3.2 Design

The study was conducted as an open design study with each subject receiving both oral and intravenous pyridostigmine. Subjects were balanced insofar as the order of drug administration was concerned (i.e., first dose oral liquid or intravenous solution). The doses were spaced three days apart. The subjects were fasted and not allowed to smoke for 8 hours before each dose and for 4 hours after each dose, when they were allowed to eat and to smoke if they desired.

As mentioned above in 2.3.1, the study was conducted in two phases, an initial dose-ranging phase followed by a formal bioavailability phase. The outline of the study and the details of how it was to be conducted as initially planned are contained in the Procedural Timetable and Study Flow Charts (Appendix B). The first three subjects received 0.66 mg of pyridostigmine bromide intravenously and 20 mg orally. Assays of pyridostigmine

concentrations revealed barely detectable concentrations of drug after the IV infusion and inhibition of acetylcholinesterase was low. Therefore, the intravenous dose was doubled. Volunteers 04-06 received 1.32 mg of pyridostigmine bromide intravenously and 20 mg orally. Plasma concentrations after both routes of administration were measurable. However, inhibition of acetylcholinesterase after oral administration of 20 mg exceeded 40% at one time point in one volunteer. Accordingly, it was decided to decrease the oral dose for the formal bioavailability trial. In that second phase, eighteen subjects were given pyridostigmine bromide, 1.32 mg intravenously and 16 mg orally.

All subjects were screened outside the hospital. Drug administration, sampling and post-drug toxicology monitoring were performed in the Drug Development Unit of The Johns Hopkins Hospital.

2.4 Clinical Laboratory Examinations

All laboratory examinations except for assay of pyridostigmine were done within The Johns Hopkins Medical Institutions. Hematology and chemistry determinations were performed by the Department of Laboratory Medicine (Clinical Laboratory License number 19-1054). The normal values for these determinations are listed in Appendix C. Erythrocyte acetylcholinesterase assays were performed in the research laboratory of the Division of Clinical Pharmacology (see section 2.5.2).

2.4.1 Hematology

Routine hematologic determinations, including hematocrit, hemoglobin, red blood cell count, white blood cell count with differential count, reticulocyte count and platelet count, were done with a Coulter counter.

2.4.2 Chemistry

Serum was assayed for sodium, potassium, chloride, carbon dioxide, urea nitrogen, creatinine, glucose, uric acid, calcium, phosphate, total protein, albumin, cholesterol, direct and total bilirubin, alanine aminotransferase, aspartate aminotransferase, alkaline phosphatase, lactic dehydrogenase and creatine kinase.

2.4.3 Electrocardiography

Standard 12-lead electrocardiographic tracings were taken on admission to the hospital and on days 3 and 6. Electrocardiograms were interpreted by a physician on the staff of The Johns Hopkins Hospital in the Division of Internal Medicine.

2.4.4 Urine Analysis

Urine analyses were performed in the laboratories of the Division of Clinical Pharmacology. Protein, ketones and bilirubin were measured qualitatively, and pH and specific gravity were determined. A microscopic examination of the

sediment was also performed.

2.4.5 Chest X-ray

Standard PA and lateral chest x-rays were usually taken just prior to admission to the hospital, and in all cases within 6 months of entry.

2.5 Pyridostigmine and Erythrocyte Acetylcholinesterase Determinations

2.5.1 Pyridostigmine Analysis

Blood samples for pyridostigmine assay were placed in heparinized Vacutainer^R tubes and iced immediately. As soon as possible (generally within five minutes) the samples were centrifuged in a refrigerated centrifuge. The plasma was separated and stored at -80° C until it was shipped in dry ice to the assay site.

The assay of pyridostigmine in plasma was performed under contract DAMD 17-85-D-0008, USAMRDC in the laboratory of Dr. Emil Lin at the School of Pharmacy, University of California at San Francisco (14). The assay utilizes protein precipitation with acetonitrile, pre-column purification on a small C8 Bond Elut^R column, and then high performance liquid chromatography on a silica column with ultraviolet detection. The detection limit is 1.5 ng pyridostigmine base/ml with a coefficient of variation between 3.55% and 8.16% (15). The plasma pyridostigmine concentrations of the patients in this study are listed elsewhere (15),

and are discussed below.

Blood for erythrocyte acetylcholinesterase Determinations
Blood for erythrocyte acetylcholinesterase

determination was collected into Vacutainers^R containing
ethylenediaminetetraacetic acid (EDTA), mixed immediately and
brought to the Clinical Pharmacology laboratories for assay -generally within five minutes of collection. Specimens were kept
at ambient temperature until assayed. The assay was performed
according to the Standard Operating Procedure (SOP) for the assay
of the Analytical Chemistry Branch, USAMR, ICD, Aberdeen Proving
Ground, Maryland 21010 dated 18 June 1985 (16). As performed at
Johns Hopkins, the coefficient of variation of the assay
determined from the quality control standard is 2.3%. Details of
the assay and its performance at Johns Hopkins are contained in
Appendix D.

2.6 Pharmacokinetic and Pharmacodynamic Analyses

2.6.1 Pharmacokinetic Analyses

Visual inspection of the pyridostigmine plasma concentration-time curves demonstrated that, following intravenous infusion, a biphasic decline in drug concentrations occurred. This suggested that the body can be represented by a two-compartment system. After oral administration, pyridostigmine concentrations rose gradually, followed by a short plateau and then a monophasic decline. In many subjects the data

were "noisy" both during the infusion and at low plasma concentrations. This "noise" made visual determination of the terminal elimination half-life difficult (and arbitrary) in many subjects. Estimates of the areas under the concentration-time curves by trapezoidal rule were also affected. Curve fitting of the intravenous data to the equation for a two-compartment model with a zero-order (constant infusion) input produced parameter estimates which varied markedly from patient to patient, suggesting that variability in the determination of the plasma concentrations was affecting adversely the curve fitting process. Since the shapes of the intravenous and oral concentration-time curves were consistent with a two-compartment open model with the rapid distribution phase being obscured by absorption during oral dosing, it was decided to fit the data after both intravenous and oral administration to a single model assuming a two-compartment open model with zero-order drug entry during infusion and with first-order absorption after a time lag following oral drug administration. This model assumes that the compartmental volumes and intercompartmental rate constants were the same during both experimental periods. Data from the oral and intravenous doses were used to fit simultaneously the equations for the time course of drug concentrations during the intravenous and oral administrations of pyridostigmine for seven constants: volume of the central compartment, K_{01} , K_{12} , K_{21} , K_{10} , time lag for absorption, and bioavailability. Data points were plotted on semilogarithmic paper. Points which appeared to be grossly

aberrant on visual inspection (far above or far beneath the two adjacent points) were not included in the curve fitting. sample times which were excluded from the analysis are listed in Table 1. PCNONLIN^R, a commercially available curve fitting program for the IBM PC (17), was used to obtain the values for the constants which provided a best fit for the observed data. weighted least squares approach was used to determine the best fit. Weights of the reciprocal of the calculated value of each point were used. In several subjects (08, 12, 14, and 22) peak pyridostigmine concentrations during the infusion did not correspond to the end of the reported infusion time. discrepancy caused large residual errors in the curve fitting process. Accordingly, for these subjects, the duration of infusion used in the curve fitting program was set to the time at which the maximum concentration of drug appeared in the plasma. Once the microconstants were determined, the macroconstants were calculated. Alpha and beta elimination rate constants are thus the same for both the oral and intravenous dosing; only the coefficients vary. The equations and the relationships between the micro- and macroconstants are shown in Table 2.

2.6.2 Pharmacodynamic Modeling

The relationship between pyridostigmine concentrations and pyridostigmine effect (erythrocyte acetylcholinesterase inhibition) was analyzed in two different ways. In both analyses the analytical equations determined from the pharmaco-

kinetic modeling described above were used to represent the plasma concentrations of pyridostigmine after intravenous and oral administration. Inhibition of erythrocyte acetylcholinesterase measured at different times was used as data. Negative and zero values and 24-hour data points were excluded.

Plots of inhibition of erythrocyte acetylcholinesterase versus plasma pyridostigmine concentration produced counterclockwise loops in all patients, suggesting that the drug effect is delayed relative to the concentration of drug. Accordingly, an "effect compartment analysis" as described by Sheiner and associates was performed (18-20). This type of analysis assumes that the delay in drug effect is due to the transfer of drug to a compartment where the drug acts, an "effect compartment." Negligible mass of drug is transported into this compartment so that the pharmacokinetic equations are not altered. assumes that drug enters the effect compartment from the central compartment at a rate K_{IE} and that drug leaves the effect compartment with a rate constant K_{E0} . Furthermore, it was assumed that pyridostigmine behaves as a standard competitive inhibitor of the erythrocyte acetylcholinesterase and that 100% inhibition of the enzyme is possible. Accordingly, inhibition of erythrocyte acetylcholinesterase is given by the concentration of pyridostigmine [Pyr] divided by the concentration of pyridostigmine plus the IC₅₀, where the IC₅₀ is the concentration of pyridostigmine in the plasma which, at steady state, produces

50% inhibition of enzyme activity.

Inhibition =
$$\frac{[Pyr]}{[Pyr] + IC_{50}}$$

Initial analyses also included a Hill coefficient. However, when the data were fit to this model, there was a high correlation between the IC₅₀ and the Hill coefficient, indicating insufficient information for an analysis with three parameters. The Hill coefficient varied around 1.0. Accordingly, it was fixed at 1 and a model including just the IC₅₀ and the K_{E0} was employed. A schematic diagram of the model is depicted in Figure 1a. Inhibition data from both the oral and intravenous dosings were used simultaneously to obtain the best fit estimates of the 50% inhibition concentration of pyridostigmine and the rate constant of elimination from the effect compartment. The curve fitting program, PCNONLIN^R, was used with a weighted least squares regression analysis with weights being equal to the reciprocal of the calculated value of the inhibition.

Erythrocyte acetylcholinesterase inhibition was also related to the concentration of pyridostigmine in the plasma by using a biochemical model that has been used previously to study the in vitro inhibition of acetylcholinesterase (21). Acetylcholinesterase is inhibited by carbamylation of the enzyme. The inactivated enzyme is then reactivated by hydrolysis. The rate constant of inactivation is given by K_{IN} and the rate constant of reactivation is the constant K_{RE} . The differential equations for the two processes were written and solved for the fractional

inhibition of acetylcholinesterase.

PCNONLIN^R was used to fit the differential equation for the values of K_{IN} and K_{RE} which best described the data for each patient. Figure 1b depicts the schematic representation of this model. When inhibition data from the oral and intravenous dosings were used simultaneously to obtain a single value for each constant, the inhibition at late time points after oral inhibition was estimated poorly. Therefore, estimates of the rate constants were performed with the data from the oral dose and separately with the data from the intravenous dose. The estimates were averaged to produce a single estimate for the inactivation and reactivation rate constant for each subject. before, weighted least squares estimates, with weights equal to the reciprocal of the calculated value, were obtained. parameter estimates in Subjects 10 and 19, obtained using the inhibition data achieved after the oral dose, were markedly different from all other estimates. These two estimates have not been included in the summary statistics, having been eliminated as outliers.

2.7 Statistical Methods

Estimates of population parameters, the mean and standard deviation of the calculated variables, have been determined using standard formulae.

3. RESULTS AND DISCUSSION

3.1 Amendments and Compliance

3.1.1 Amendments

The consent form was amended prior to enrolling the first subject to provide information about toxicity in rats. The original consent form and the amended consent form are provided as Appendix E and F, respectively.

The protocol was amended after the first three subjects were completed to increase the intravenous dose of pyridostigmine bromide to 1.3 mg. The protocol was amended after subjects 04, 05, and 06 were completed to reduce the dose of oral pyridostigmine bromide to 16 mg. (Details are provided in section 2.3.2 of this document.) The protocol was amended to collect 24-hour urine specimens for pyridostigmine assay for the last 8 subjects. The institutional review board at Johns Hopkins and the Army's Human Subjects Research Review Board were notified of these changes and granted approval.

3.1.2 Compliance

After the completion of the study it was

discovered that the subjects had signed the originally approved consent form, not the amended version. The institutional review board at Johns Hopkins was notified of this oversight. No attempt was made to contact the subjects who had participated. We believe this had no bearing on the conduct of the study or on the willingness of our subjects to participate.

As mentioned in section 2.2.1, subjects were enrolled into the study even if screening and/or admission CK levels were elevated. We believe these elevations do not reflect a myopathic process, but rather reflect the extremely active physical lifestyle of the subjects. We are satisfied that these subjects were healthy and that those with elevated CK values did not have a pathological process.

Although we had requested and received approval to give 1.3 mg intravenous pyridostigmine bromide, we actually gave 1.32 mg. We believe this deviation is inconsequential.

There were six timed intervals for urine collection within each 24-hour period after dosing, not a single continuous 24-hour collection. This would allow for more precise information regarding urinary pyridostigmine excretion patterns. These data are included in the report prepared by Cheng et al. (22), and are discussed in the Results section (section 3.4.1) of this manuscript.

The baseline (pre-drug) electrocardiogram was done at screening and not on the day of admission to the hospital. This deviation is of no clinical importance.

Screening red blood cell acetylcholinesterase was not done before entry to the hospital. Rather, each subject's baseline red blood cell acetylcholinesterase level was determined before receiving the first dose of pyridostigmine and ascertained to be within the normal range.

Hand grip determination of muscle strength was not done after 09 January 1986 when the electronic device used for this purpose stopped working accurately. Values for this parameter are only complete for the first ten subjects. Neither the sponsor nor the institutional review board at Johns Hopkins was notified. There was no pattern of change in hand grip strength after pyridostigmine in the first ten subjects, and we believe this same pattern would have been similar for the remainder of the subjects. Furthermore, we believe that hand grip testing is a gross, and, in part, subjective measure of changes in voluntary muscle strength after pyridostigmine. It is subjective in that it is effort related and subject to the volunteer's interest and willingness to try hard.

The protocol divided the study into two phases: dose-ranging and bioavailability. The dose-ranging phase required 12 red blood cell acetylcholinesterase determinations for each dose, both intravenous and oral, while the bioavailability phase was designed to have 7 determinations for each dose. For all the intravenous doses, however, 12 samples were collected and assayed from all 24 subjects. No one was notified in advance of this protocol deviation, but we feel confident that these additional

samples were essential to allow the comprehensive pharmacokinetic and pharmacodynamic determinations included in this report. With regard to the oral dose, not only were the 12 sampling times intended just for the dose-ranging phase perpetuated into the bioavailability phase, but two additional sampling times were added, at 8 and 10 hours after the dose, when it became clear that acetylcholinesterase inhibition was still significant (approximately 10%) at 6 hours after the dose. Again, we believe that these additional data points were essential in enabling us to conduct the extensive analysis of pharmacodynamics outlined below.

3.2 Description of Population of Subjects

During the course of the study, a total of two newspaper advertisements were placed. Sixty-six men responded and, of this group, 24 were entered into the study. Reasons for rejecting potential subjects were:

- 66 Scheduled for screening
 - -24 Failed to keep scheduled appointment
- 42 Seen at Johns Hopkins Hospital
 - -12 Abnormal screening laboratory tests
 - 4 Changed mind, moved, etc.
- 26 Examined by physician
 - -2 Failed history or physical exam
- 24 Admitted to Johns Hopkins Hospital and entered the study
- 24 Completed Study

Of the 24 who completed the study, 4 were white and 20 were black. The average age was 26.4 years and ranged from 19 to 35 years. Relevant vital statistics of the volunteers are listed in Table 3. Creatinine clearance was estimated from the Cockcroft-Gault formula (23) using the average of three in-hospital serum creatinine determinations for the serum creatinine concentration.

3.3 Clinical Results

3.3.1 Symptomatic

The subjects tolerated the administration of pyridostigmine very well and none complained of any side effect with either the intravenous or oral administration of the drug. In particular, there were no complaints of nausea, abdominal discomfort, change in bowel habits, muscle aching or tenderness, weakness, tremor, or any other neuromuscular complaint.

3.3.2 Pulse

No patterns of pulse change were consistently or predominantly seen in the subjects with administration of either intravenous or oral pyridostigmine. The pulse results appeared to be completely random and appeared to have no correlation with plasma pyridostigmine concentrations or with the degree of erythrocyte acetylcholinesterase inhibition.

3.3.3 Coordination Testing

There was no change in coordination testing compared to the baseline performance for any of the volunteers during or after administration of either intravenous or oral pyridostigmine. It should be noted that the coordination testing used was limited in its capacity to identify abnormalities. More refined and accurate methods of measuring coordination would need to be employed in order to identify subtle changes in coordination.

3.3.4 Hand Grip

As was the case with the pulse measurements, there was no recognizable or predominant pattern of change in hand grip for the subjects following administration of pyridostigmine. Not only was there little in common from one subject to the next, but there were also occasional discrepancies in the pattern of strength change over the period of observation in one hand compared to the other in the same subject. The hand grip testing was accomplished only for the first 10 subjects and was not done for the remainder because of technical difficulties with the hand grip measuring device.

3.3.5 Laboratory

3.3.5.1 Liver Function Tests

The serum aspartate aminotransferase

(AST) levels increased in two subjects to a substantial degree

and above the upper level of normal (35 units/liter) after their second exposure to pyridostigmine, which in both cases was the intravenous dose. The levels increased from 20 to 44 units per liter in one subject and from 19 to 44 units per liter in the other. In only one of these subjects was the serum alanine aminotransferase (ALT) level found to increase as well, increasing from a predrug level of 12 to a level of 36 units per liter, 24 hours after the intravenous (second) dose. (The upper limit of normal is 30 units/liter). One other subject had a substantial increase in the serum alanine aminotransferase in the absence of a corresponding increase in the serum aspartate aminotransferase to an abnormal level. In this individual the ALT increased from 19 to 36 units per liter, again following the second exposure to pyridostigmine, which was the intravenous dose. No other subject had a noteworthy change in the serum transferases, and no subject had a significant change in alkaline phosphatase, serum cholesterol, total protein, serum albumin, direct or total bilirubin. The serum lactic dehydrogenase (LDH) increased modestly in one subject from a predrug level of 195 to a peak of 224 units per liter after the first (oral) dose of pyridostigmine and was back to normal 24 hours after the second (intravenous) dose of pyridostigmine, to a level of 170 units per liter. (The upper limit of normal is 200 units/liter.) subject with the increase in LDH did not have an increase in either of the transferases. Thus, while the LDH may arise from the liver in some cases, it may originate from other organs and

there was no corroborative evidence suggesting that the liver was the source of this subject's minimal elevation.

3.3.5.2 Creatine Kinase

The creatine kinase was monitored in hopes of using it as a sensitive and fairly specific indicator of muscle cell damage. Interestingly, the creatine kinase level was well above normal in at least half of the subjects prior to receiving either formulation of pyridostigmine, and in most cases gradually fell over the course of the five day admission, consistent with muscle cell injury prior to admission to the hospital and gradual recovery over the course of the hospitalization. It is our impression that these elevations probably represent minor to modest degrees of muscle fiber damage as a result of strenuous exercise, physical exertion during gainful labor, or minor degrees of trauma suffered by our volunteers prior to admission. Only one subject had a remarkable increase in the creatine kinase. This individual had an increase from a level of 94 units per liter 24 hours after the first (oral) dose of pyridostigmine to a peak of 2,060 units per liter 24 hours after the second (intravenous) dose. As was mentioned above, neither this subject nor any others complained of symptoms suggestive of muscle injury, such as soreness, stiffness, or aching. Two other subjects had very modest elevations of creatine kinase following their second administration of pyridostigmine, in one case the administration being oral and in the

other case the administration being intravenous. In neither of these two subjects, however, was the increase nearly so impressive as that of the first subject described above.

Therefore, we believe that the substantial elevation of creatine kinase in the one volunteer may be evidence for subclinical muscle injury in response to intravenous pyridostigmine, but the elevations seen in the other two subjects were clinically insignificant.

3.3.5.3 Other Clinical Chemistry Tests

No subject had an abnormality of any of
the other monitored chemistry tests mentioned above under
"Methods."

3.3.5.4 Hematological Testing

Three of the subjects had very mild reductions in their hemoglobin concentrations and hematocrits over the course of the study. The magnitude of these changes, however, was consistent with the phlebotomies being performed to obtain blood samples for monitoring and outcome measurements. No remarkable changes in white blood cell count occurred. Two subjects developed a very mild relative eosinophilia in the absence of symptoms or signs of hypersensitivity. There were no significant elevations of reticulocyte count over the course of the study.

3.3.5.5 Electrocardiograms

There were no significant changes in electrocardiograms following any dose of pyridostigmine. The only noteworthy changes were mild changes in electrical axis and non-specific ST-T wave changes, both of which might be due to expected intrasubject variability, though a drug effect cannot be excluded.

3.3.6 Clinical Conclusions

It is our view that the subjects tolerated the administration of pyridostigmine in both the oral and intravenous forms very well. The minimal elevations of transferases in 3 of the subjects were not clinically worrisome. The substantial elevation of creatine kinase in one subject is the only observation of clinical importance and could possibly be related to an increased propensity for muscle cell damage in response to intravenous pyridostigmine in this volunteer. Further evaluation of this observation, both in this subject and in others, would be warranted in our view. In our opinion, this single observation does not constitute grounds for any more specific action than to continue our close observation of this parameter.

3.4 Pharmacokinetics and Pharmacodynamics

3.4.1 Pharmacokinetics

Data from the 18 volunteers, 07-24, who received 1.32 mg of pyridostigmine bromide intravenously and 16 mg orally

were analyzed. The times the samples were obtained during and after the intravenous infusion are listed in Table 4. Measured plasma pyridostigmine base concentrations are found in Table 5. Plasma concentrations of pyridostigmine rose rapidly during the intravenous infusion and declined once the infusion had stopped. Plasma concentrations were no longer measurable six hours after beginning the infusion (Figure 2 and Tables 4 and 5). Concentrations during and after the infusion were quite variable. Coefficients of variation at each sample time were greater than 25%. Peak concentrations at the end of the infusion varied as much as fourfold. During the phase of declining concentrations variations were somewhat less but still substantial. Individual plasma concentration curves are depicted in Figures 2a, 2c, and 2e.

Pyridostigmine concentrations following the dose of oral syrup rose slowly, often after a short time delay (Tables 6 and 7). The mean plasma concentrations were nearly constant from one to three and a half hours after dosing and then declined slowly over time. At ten hours after the dose pyridostigmine was still detectable in some subjects (Table 7 and Figures 2b, 2d, 2f). Variations between subjects were greater following oral dosing than following intravenous dosing (Figures 2a-2f). Coefficients of variation were greater than 32% at each time. In some subjects there was rapid absorption with a high peak concentration of pyridostigmine achieved and a relatively rapid decline (e.g., Subject 18) while in other subjects absorption was

delayed, the peak level was later and the decline slower (e.g., Subject 16).

The approach of fitting the oral and intravenous data simultaneously predicted the actual plasma concentration data using a two-compartment open model with different types of drug entry depending upon the route of administration. The sum of the weighted squared residuals and the correlations between the calculated and actual data points for each subject for both routes of administration are shown in Table 8. There was little difference between the means of the sum of the weighted squared residuals for the oral and intravenous routes of administration. Correlation coefficients were generally high, usually above 0.9. The correlation between the calculated and observed data for the intravenous route was higher than that for the oral route (p <0.002, paired t-test). Several factors in the design and analysis of the trial may be responsible for this. First, more data points were collected during intravenous dosing than during the oral dosing and plasma concentrations tended to be higher, reducing the problems created by low measured values approaching the sensitivity of the assay. In this way the weighting procedure may have been biased toward the intravenous dosing. Second, an assumption was required to model the kinetics of the oral dose, i.e., first-order absorption of drug, which was not needed for the intravenous data. The absorption of pyridostigmine may, in fact, not follow strictly first-order kinetics, in which case the actual and estimated concentrations

would be expected to differ. For the intravenous dosing, the infusion was a constant one and modelled as such. In the few subjects where unusual concentration profiles occurred during the infusion, the model was modified (see 2.6.1). Finally, estimates of three parameters were obtained from the data following the oral dose in addition to the estimates derived from both the intravenous and oral data. The requirement of seven as opposed to four parameter estimates may have introduced a greater discrepancy between the actual and estimated plasma concentrations for the oral dosing.

The estimates of the pharmacokinetic microconstants for each subject are detailed in Table 9. Due to the simultaneous fit, there is only a single estimate for each of the microconstants. For each constant a wide variation between subjects exists, the standard deviation of the estimate being greater than 50% of the average estimate in all cases except for the bioavailability. The pharmacokinetic macroconstants, derived from the microconstants, are shown in Table 10. As in the case of the microconstants, there is great variation between subjects. coefficient of variation is greater than 40% for each constant. The variability in the magnitude of these constants is reflected in the calculated areas under the concentration-time curves and the plasma clearances of pyridostigmine as well. The values for each subject are depicted in Table 11. Coefficients of variation of about 30% were present in the areas under the concentrationtime curves following both intravenous and oral dosing. There

was a similar variation in the calculated total plasma clearances. Adjusting the total clearance for body weight or for renal function (estimated creatinine clearance) did not reduce the coefficient of variation.

The results of this study are similar to data collected by others who studied smaller numbers of subjects. Aquilonius et al. studied the intravenous kinetics of 2.5 mg of pyridostigmine bromide given to 2 subjects (24). Total clearance of drug was 806 and 828 ml per minute or 10.6 and 10.9 ml/min/kg. half-life of drug was 1.7 and 1.3 hours. Calvey and coworkers studied 10 patients after administering intravenously 3.65 mg of pyridostigmine bromide per 70 kg body weight (25). Clearance of drug in these patients averaged 16.3 ml/min/kg with a range of 9.3 to 26.5. Since the body weights of the individual subjects were not provided in their report, the values for the total clearances not normalized for weight cannot be calculated from the data. The beta half-life averaged 0.38 hours, ranging from 0.25 to 0.62 hours. The results of the current study (mean total clearance of 779 ml/min with a range of 382 to 1511 ml/min) are similar to the data above. When corrected for body weight, the present data yield a mean clearance of 10.3 ml/min/kg, a value similar to that obtained by the others. Likewise, our mean beta rate constant is 0.768 per hour which translates to a beta half-life of 0.90 hours, a value well within the range of the earlier data. As all these studies were performed with different doses of pyridostigmine, it appears likely that the clearance of

intravenous pyridostigmine is independent of dose.

The variance in the total clearance estimates among our subjects is not reduced by normalizing drug clearance for the estimated creatinine clearance, suggesting that total pyridostigmine clearance is either largely non-renal or that renal pyridostigmine excretion is not highly correlated with the glomerular filtration rate, at least in subjects with normal renal function. For subjects 17 through 24, urine was collected for 24 hours following dosing and assayed for pyridostigmine base using a modification of the assay for plasma (22). Amounts of pyridostigmine base excreted during each collection period are displayed in Table 12. Pyridostigmine was detectable in the urine within one hour after both intravenous and oral dosing. The amount of pyridostigmine excreted declined rapidly with time after intravenous dosing but persisted somewhat longer after oral dosing. The amounts of pyridostigmine excreted were variable among subjects, ranging from 33.8 to 84.9% of the intravenous dose and from 4.8 to 19.8% of the oral dose (Table 13). These urinary recoveries are similar in magnitude and variability to those found by Nowell et al. (26), who studied 5 patients with myasthenia gravis receiving 180 to 3600 mg of pyridostigmine bromide daily. Urinary excretion of intact pyridostigmine ranged from 2 to 16% of the dose with as much as 70% variation on different study days. Somani et al. recovered 9.4% of the pyridostigmine dose in the urine in the one patient they studied (27).

Renal clearances calculated using the fractional excretion of pyridostigmine after intravenous dosing times the total clearance are tabulated in Table 14. Renal clearance estimates range from 215 to 854 ml/min with a mean of 430 ml/min. All exceed the creatinine clearance of the subjects, implying renal secretion of pyridostigmine. The renal clearance estimates in this study are of similar magnitude to those obtained by Chan and Calvey (28) except for two high values of 854 and 694 ml/min. clearance rates, approaching renal plasma flow, are unexplained. The magnitude of the clearance values is consistent with the hypothesis that pyridostigmine is secreted by the secretory system for organic bases which resides in the kidneys. pathway is located in the proximal renal tubule and actively transports organic cations, including primary, secondary, tertiary and quaternary amines, in the direction of excretion (29). Renal clearance of such compounds can approximate renal plasma flow. Within the cationic system, however, there can be competition between compounds for excretion, including endogenous compounds, presumably mediated by competition for the putative carrier protein (29). The observation by Chan and Calvey that the renal secretion of pyridostigmine was reduced in two patients who were being treated concurrently with a basic drug (28) is also consistent with transport by the cationic transport system.

Little prior information exists on the bioavailability of oral pyridostigmine. Aquilonius et al. studied 2 patients to whom they administered 120 mg of pyridostigmine bromide in the

fasting state (24). The dosage form of the drug was not identified in their article. Pyridostigmine was 5.1 and 11.4 percent bioavailable in these subjects. Breyer-Pfaff et al. administered 60 mg of pyridostigmine bromide as a dragee 30 minutes after a standard meal (30). They found the drug to be 14.3% available with a range of 11.5 to 18.9 percent in the 10 patients studied. A much smaller dosage of drug, 16 mg, was to be used in the current study, and assay of the syrup plus determination of the weight of syrup administered revealed the actual dose to be an average of 14.8 mg (Appendix A). Bioavailability was much higher with this dose than the doses mentioned above, ranging from 14.7 to 51.1 percent with a mean of 29.2 percent. These data suggest that the fractional absorption of pyridostigmine decreases as the size of the dose is increased. However, the absolute amount of drug absorbed, calculated as the product of the dose administered times the bioavailability, does increase with increasing dose (with the caveat that different dosage forms and different conditions were present in each study). When the administered dose is calculated in milligrams of pyridostigmine base, the amount of drug given to volunteers in the three trials was 83.3, 41.6 and 10.3 mg, respectively. average amount of drug absorbed was 6.8, 6.0 and 2.9 mg, respectively, with considerable scatter within each study. are depicted in Figure 3. Absorption, therefore, is not directly proportional to the dose. Even excluding the highest dosage, a fourfold dosage increment produced only a twofold increase in

absorbed drug. The curve may also bend (although data are limited) which hints at capacity-limited absorption of drug.

Systemic availability of pyridostigmine also can be estimated from the recovery of intact drug in the urine. Bioavailability calculated in this manner is listed in Table 14 and compared with the values obtained from the pharmacokinetic parameter estimation (Table 9). In general, the bioavailability calculated from the urinary data is lower (7 of 8 subjects), suggesting a systematic The cause of this is not certain. One possibility is that pyridostigmine decomposed in the urine during the collection process. This might be expected to be greatest in the 8-24 hour specimen. As shown in Table 12, urinary excretion after intravenous dosing was rapid, and nearly complete by the end of 8 hours, while urinary excretion after oral dosing showed greater amounts in the later collections compared to intravenous dosing. Thus, if there were decay of pyridostigmine in the urine during the 8-24 hour collection, the end result would be an underestimation of excretion predominantly after oral dosing, thus underestimating bioavailability.

The limited degree of absorption, the hint of capacitylimited absorption, and the positive charge on the pyridostigmine
moiety all suggest that passive diffusion across the gastrointestinal membrane is not the mechanism for absorption of
pyridostigmine. Solubilization of drug is unlikely to be a
problem since pyridostigmine is a charged molecule which is very
soluble in an aqueous environment. Two mechanisms, closely

related, which could account for both the variation in availability between subjects at the same dose and for the apparent saturation in absorption with increasing dose can be postulated. The first invokes a saturable transport mechanism, either active transport or facilitated diffusion. Variations in the capacity or affinity of the transport system for pyridostigmine could result in differences in availability. Second, there might be compounds within the gastrointestinal tract, possibly endogenous, which are transported by the same system. These compounds might then compete with pyridostigmine for uptake; differences in the amounts of competing compounds from subject to subject would produce the variation in availability. Since all bioavailability studies were conducted with the subjects fasted at least 8 hours, food was not a factor in this study.

The small rate constant of absorption (mean equal to 0.373/hr) which is equivalent to a half-time of absorption of 1.86 hours, is also consistent with a capacity-limited transport process. Again, drug solubility is not likely to be an issue. Given in solution, pyridostigmine should be accessible to membranes almost immediately. Arrival at the absorption site and crossing the membrane must be the rate limiting steps. The time lag for absorption to begin, averaging over twenty minutes, represents the time required to arrive at the membrane transfer site, and variations in gastrointestinal motility might be responsible for the large differences observed in the time necessary to initiate absorption. Alternatively, the transport

hypotheses postulated earlier could account for some of the observed differences in the delay of absorption.

3.4.2 Pharmacodynamics

Following pyridostigmine administration inhibition of erythrocyte acetylcholinesterase occurred.

Inhibition increased rapidly during the intravenous infusion, reaching a mean of 30% at the end of the 30-minute infusion (Table 15). Enzyme inhibition declined thereafter, and six hours after dosing the enzyme activity had returned to baseline. The degree of inhibition at the end of the infusion varied almost twofold, from 22% to 39%. Later, more than one and a half hours after the onset of the infusion, inhibition varied by as much as threefold, though the absolute differences between subjects were less, no more than 11%. The mean inhibition as a function of time is shown in Figure 4, with data of individual patients depicted in Figures 4a, 4c, and 4e.

Inhibition of erythrocyte acetylcholinesterase following oral pyridostigmine developed less rapidly (Table 16 and Figure 4). There was a gradual rise in mean inhibition with the peak occurring at three hours after dosing. Inhibition was nearly constant, however, from 1.3 to 4.0 hours. Thereafter, the proportion of enzyme inhibited declined so that 10 hours after dosing the mean inhibition was only 6%. At 24 hours, the next sample time, no inhibition was measured. Variations from subject to subject in the amount of inhibition achieved were greater

during the oral dosing than during the intravenous dosing (Figures 4b, 4d and 4f and Tables 15 and 16). Differences between maximum and minimum amounts of inhibition at each sample time ranged from 10 to 20%. At times later than 4 hours after the oral dose, some subjects had three to four times as much enzyme inhibition as others. Ten hours after dosing, inhibition ranged from 2 to 11% of baseline enzyme activity.

Inhibition of erythrocyte acetylcholinesterase above 20% was more effectively prolonged after oral dosing than after intravenous administration. During the intravenous dose at least 20% inhibition was reached in all subjects. After the infusion was completed, however, the degree of inhibition fell rapidly. By 1.17 hours after the end of the infusion, inhibition was less than 20% in all subjects (Table 15). The mean duration of inhibition greater than 20% was only 0.49 hours (Table 17). The longest duration, observed in Subject 12, was only 1.08 hours. In contrast, after the oral dose, inhibition of erythrocyte acetylcholinesterase remained greater than 20% for an average of 2.62 hours (Table 18). Variability of inhibition was greater with oral than with intravenous administration. After oral dosing, inhibition over 20% was present at only one measurement in Subject 9 (Table 16), while inhibition greater than 20% was present for 5 hours in Subject 14 (Table 18). The maximal inhibition of erythrocyte acetylcholinesterase in each subject for each dosage form is provided in Tables 19 and 20. The mean time to peak inhibition was 2.5 hours after oral dosing (Table

20), while in all cases but one (Subject 9) the peak inhibition with intravenous dosing was seen at 0.5 hours, corresponding to the end of the infusion (Table 15).

For both the intravenous and oral dosing, the fraction of red blood cell acetylcholinesterase inhibition was plotted against the concentration of pyridostigmine in the plasma (Figure 5a and 5b). When the data points were connected in order of time after dosing, an anticlockwise loop was obtained. A loop in this direction suggests that the effect of the drug is delayed relative to the plasma concentration of the drug. The area within the hysteresis loop for both doses for each patient was calculated. The mean area in the loop following the intravenous dose and the mean area in the loop following the oral dose were each significantly greater than zero. In addition, the area of the loop during the intravenous dose was greater than that from the oral dose. Data of individual subjects are detailed in Table 21.

Since the above plots suggested a delay in the effect of the drug, we employed an effect compartment model to explain the delay. The pharmacokinetic equations previously described were used to estimate pyridostigmine in the effect compartment. Using the oral and intravenous inhibition data simultaneously, we sought the K_{E0} and the IC_{50} concentration of pyridostigmine which best described the data.

The ability to fit the data to this model was quite good (Table 22). Correlations between the predicted inhibition and

observed inhibition were above 0.8 in all instances and in the majority of cases above 0.9. The estimates had slightly higher correlations with the intravenous data than the oral data but the difference was not significant. The sum of the weighted squared residuals was small in all instances of fitting the intravenous data and in most instances for fitting the oral data. The estimate was particularly high for the oral dosing of patient 10 for reasons not apparent. Curve fittings of the oral data were not as good as those for the intravenous data. This may be due in part to the fact that the equations for the pyridostigmine concentrations did not describe the actual oral data as well as they did the intravenous data.

The mean K_{E0} , or rate constant of elimination from the effect compartment, was 3.42 per hour, equivalent to a half-time at steady state in the effect compartment of 0.2 hours (Table 23). There was a greater than fourfold variation in the magnitude of the estimates of this constant, from 1.22 to 5.76 per hour. These rate constants are equivalent to a half-time of effect of 0.57 and 0.12 hours, respectively. The concentration of pyridostigmine in the plasma which would produce 50% inhibition of effect at steady state, the IC_{50} , also varied. The mean of the estimates was 29.5 ng/ml. Individual estimates, however, ranged from 17.3 to 41.1 ng/ml, a 2.5-fold difference.

This analysis suggests that there are significant interindividual differences in the sensitivity of erythrocyte acetylcholinesterase to pyridostigmine. These differences are

manifest both in the time it would take, at a constant pyridostigmine concentration, to achieve steady state inhibition of enzyme and also in the amount of pyridostigmine required in the plasma to produce a given amount of inhibition of the erythrocyte acetylcholinesterase. In the first instance, since three to five half-lives are required to reach steady state, one would expect that it would take from 20 minutes to 2.5 hours to achieve steady state inhibition of enzyme, depending on the rate comstant of elimination (K_{E0}) from the effect compartment. Regarding the IC_{50} variability, if the pharmacokinetic parameters were identical in a group of subjects, some subjects would still require 2.5 times as much drug to produce a given effect as others. Those with low IC_{50} 's would be sensitive to the drug and would require lower doses, while those with elevated IC_{50} 's would require larger amounts of drug.

The effect compartment analysis, while useful, is not a completely accurate representation of the interaction between pyridostigmine and the erythrocyte acetylcholinesterase. The imteraction is not really reversible, at least in the classical sense of almost instantaneous association and dissociation of emzyme with inhibitor. After carbamylation the hydrolysis, equivalent to dissociation, is delayed. Furthermore, since the red blood cell is circulating in the plasma, the interaction between drug and enzyme should occur in the central, highly perfused, kinetic compartment. Given these conditions, it seemed reasonable to model the interaction between enzyme and drug

producing enzyme inhibition using available information about the biochemistry of the enzyme. It was assumed that separate rate constants describe the inactivation of enzyme by pyridostigmine and the reactivation of inactivated enzyme by hydrolysis. two constants were denoted K inactivation (KIN) and K reactivation (KRE), respectively. Attempts at fitting the observed inhibition data to the model using the data from both the intravenous and oral doses simultaneously were unsuccessful; the inhibition which occurred at late time points following oral dosing could not be accurately estimated. However, when separate estimates were produced from the oral dosing and from the intravenous dosing, reasonable estimations of acetylcholinesterase inhibition could be produced. As in the case of the effect model, estimates of the effect following the intravenous dose were marginally better than the estimates for the effects after the oral dose. Sums of the weighted squared residuals were small for both routes of administrations and in all patients. Correlation coefficients for the relationship between the observed and predicted inhibition were greater than 0.9 in all cases but one. Correlation coefficents were higher for the estimates of the intravenous effects than for the oral (p <0.02 by paired t-test). The data for the "goodness of fit" to this model for each subject are detailed in Table 24.

Two estimates of each biochemical rate constant, one from the intravenous dosing and one from the oral dosing, were obtained for every subject and averaged (Table 25). The two estimates for

each constant in each patient were generally similar. Nevertheless, the estimates in two subjects (10 and 19) using the data obtained after oral administration of drug were clearly at variance with the 34 other estimates. The reasons for this are not apparent. Subject 10 had unusual pharmacokinetics with a much greater plasma clearance of pyridostigmine than the other volunteers. The parameter estimate for K_{IN} generated from Subject 10's intravenous data is the largest estimate from the intravenous data, but the estimate for K_{RE} is not substantially different from those generated in the other subjects. estimate of the constants in Subject 19 from the data following intravenous dosing of pyridostigmine and the pharmacokinetics of pyridostigmine appear similar to the estimates in the other subjects. The two aberrant results have been excluded from the computation of the estimates of the mean and standard deviation of K_{IN} and K_{RE} . Whether the two unusual values are a consequence of the curve fitting process and thereby artifactual, or whether they represent some unusual occurrence during the oral dosing of these two subjects, is not known.

Excluding the aberrant results of Subjects 10 and 19 as described above, there was a coefficient of variation of 25 to 33% in the estimates of the biochemical rate constants. The mean inactivation rate constant was 0.087 ml/ng/hr and the mean reactivation rate constant was 2.736/hr. The latter represents the rate at which inactive enzyme is reactivated. This mean rate constant is equivalent to a half-time of reactivation of 0.25

hours, indicating that, on the average, it would take three to five half lives, or 0.75-1.25 hours, for inactivated enzyme to be completely reactivated, even in the absence of pyridostigmine. The smallest reactivation rate constant was 1.65/hour. In this subject it would take up to two hours to completely reactivate the enzyme. The heterogeneity in these two rate constants is similar in magnitude to the variation in the magnitude of the effect compartment elimination rate constant and the IC₅₀ determined in the preceding analyses.

The values for the two estimated constants can be expressed in the units used to describe biochemical reaction rate constants. These are, in the case of K inactivation, liters per mole per minute, and in the case of the reactivation rate constant, per minute. The estimate of the inactivation rate constant of pyridostigmine for acetylcholinesterase in intact erythrocytes is thus $26.3 \pm 8.2 \times 10^4$ L/mole/min. This compares quite favorably to an estimate of 19.8 x 10^4 L/mole/min for k_i , the reaction rate of pyridostigmine with purified bovine erythrocyte acetylcholinesterase in vitro at 37°C measured by Watts and Wilkinson in 1977 (21), and an estimate of 1.6 x 104 L/mole/min for the inactivation rate of electric eel acetylcholinesterase performed in vitro by Wilson et al. (31). The latter experiments were performed at 25 degrees Celsius; the reaction rate would be expected to be higher at body temperature (37°C), in part explaining the tenfold difference. The mean estimate obtained in the present study for the rate of enzyme reactivation (hydrolysis

ke three to i enzyme to be ostigmine.

In this eactivate cants is of the

%pressed

ers per

ite act

pares

k_i,
ythro-

in e

ents be

in

restored immediately after drug is removed. Rather, enzyme inhibition is gradually dissipated as the carbamylated, inhibited enzyme is reactivated by hydrolysis. The intersubject variations in the estimates for the rate constants may represent true differences in the interactions between the drug and the enzyme, possibly related to molecular heterogeneity of the enzyme. Another possibility is that these differences are artifacts of the experimental, analytical, and mathematical processes which were required to obtain the estimates. Small errors generated in each of these processes might have contributed sufficiently in the aggregate to produce the scatter in the estimates.

4. CONCLUSIONS

Analysis of the experimental data from this study indicates that significant interindividual variation in erythrocyte acetylcholinesterase inhibition will result from a uniform dose of pyridostigmine. This interindividual variation in drug effect is a consequence of two major factors: (1) the variation in the pharmacokinetic parameters in any group of individuals, and (2) the variability of the concentration-inhibition relationship (pharmacodynamics) between pyridostigmine and acetylcholinesterase in different subjects.

Pharmacokinetic variations appear to be at two major sites.

The first major factor in the variability of pyridostigmine pharmacokinetics is in the bioavailability of the oral

formulation of the drug. There were large differences in both the extent and rate of absorption of pyridostigmine in the volunteers in this trial. The amount of drug absorbed from the liquid preparation given varied fourfold within these 18 subjects. This is presumably a situation where the pharmaceutics are optimal for drug absorption, i.e., a freely soluble drug adminstered in solution. Other workers using different doses and different dosage forms have also seen significant variation in the extent of pyridostigmine absorption. Differences in the time of onset of absorption were also observed. Nevertheless, while these might be fairly significant for single dose administration, for chronic dosing the time lag for absorption would not be a major factor in the systemic availability of pyridostigmine. Second, there are large differences in the plasma clearance of pyridostigmine between healthy volunteers. These differences cannot be explained on the basis of the estimated glomerular filtration rates of these individuals, even though the drug is apparently excreted primarily by the kidney. This suggests that some other variable (as yet unidentified) is a major determinant of pyridostigmine elimination. The large differences in the amount of drug entering the systemic circulation and in the variability in the rate at which drug leaves the systemic circulation would therefore be expected to result in large individual differences in the steady state plasma concentrations of pyridostigmine with any standard dosage regimen if it were applied to a large population.

The analyses of the concentration-inhibition (pharmacodynamic) relationship between pyridostigmine and erythrocyte acetylcholinesterase from this study also indicate that there is significant interindividual variation in enzyme inhibition at any given concentration of pyridostigmine. analyses performed, i.e., the effect compartment model and the biochemical model, suggest that within the small number of subjects studied there is a twofold difference in the sensitivity of erythrocyte acetylcholinesterase to pyridostigmine. even if a dosage form and dosage regimen could be devised to achieve a target pyridostigmine concentration in the plasma, there would be nearly a twofold difference in the amount of erythrocyte acetylcholinesterase inhibition in the population because of differences in the sensitivity of the enzyme to the drug. To the extent that the red cell enzyme is a marker for acetylcholinesterase at other sites, protection against organophosphate poisoning would also be expected to be highly variable. Therefore, a strategy to provide optimum protection against organophosphate poisoning may require individualized dosage regimens using the effect of the drug as the end point.

REFERENCES

- 1. Berry WK and Davies DR. The use of carbamates and atropine in the protection of animals against poisoning by 1,2,2trimethylpropylmethylphosphonofluoridate. Biochem Pharmacol 1970; 19: 927-934.
- Gordon JJ, Leadbeater L, Maidment MP. The protection of animals against organophosphate poisoning by pretreatment with carbamate. Toxicol Appl Pharmacol 1978; 43: 207-216.
- 3. Dirnhuber P and Green DM. Effectiveness of pyridostigmine in reversing neuromuscular blockade produced by soman. J
 Pharm Pharmacol 1978; 30: 419-425.
- 4. Adler M, Maxwell D, Foster RE, et al. <u>In vivo</u> and <u>in vitro</u> pathophysiology of mammalian skeletal muscle following acute and subacute exposure to pyridostigmine: Studies on muscle contractility and cellular mechanisms. Proc 4th Annual Chemical Defense Bioscience Review, U. S. Army Medical Research and Development Command. Aberdeen Proving Ground, Maryland, 30 May 1 June 1984, pgs 173ff.
- 5. Hudson CS and Foster RE. Ultrastructural pathology in mammalian skeletal muscle following acute and subacute exposure to pyridostigmine: Studies of dose-response and

- recovery. Proc 4th Annual Chemical Defense Bioscience
 Review, U. S. Army Medical Research and Development
 Command. Aberdeen Proving Ground, Maryland, 30 May 1 June
 1984, pgs 131ff.
- 6. Wolf PL, Lott JA, Nitti GJ, and Bookstein R. Changes in serum enzymes, lactate, and haptoglobin following acute physical stress in international-class athletes. Clin Biochem 1987; 20:73-77.
- 7. Occhi G, Gemma S, Buselli P, and Miserocchi G. Effects of repeated endurance. J Sports Med 1987; 27:184-190.
- Clarkson PM, Apple FS, Byrnes WC, et al. Creatine kinase isoforms following isometric exercise. Muscle Nerve 1987;
 10:41-44.
- 9. Stendig-Lindberg G, Shapiro Y, Epstein Y, et al. Changes in serum magnesium concentration after strenuous exercise. J

 Am Coll Nutr 1987; 6:35-40.
- 10. Nicholson GA, Morgan GJ, Meerkin M, et al. The effect of aerobic exercise on serum creatine kinase. Muscle Nerve 1986; 9:820-824.

- 11. Evans WJ, Meredith CN, Cannon JG, et al. Metabolic changes following eccentric exercise in trained and untrained men.
 J Appl Physiol 1986; 61:1864-1868.
- 12. Stansbie D, Aston JP, Dallimore NS, et al. Effect of exercise in plasma pyruvate kinase and creatine kinase activity. Clin Chim Acta 1983; 132:127-132.
- 13. Newham DJ, Jones DA, and Edwards RHT. Large delayed plasma creatine kinase changes after stepping exercise. Muscle Nerve 1983; 6:380-385.
- 14. Lin ET, Benet LZ, Upton RA, Gee WL. High pressure liquid chromatography (HPLC) of pyridostigmine in plasma using silica gel column and an aqueous mobile phase. Study Report No. 5, U. S. Army Medical Research and Development Command, July 21, 1986.
- 15. Lin ET, Gee WL, Yturralde O. Routine analysis of pyridostigmine plasma samples from Johns Hopkins. Analysis Report
 No. PY-85-6-3, U. S. Army Medical Research and Development
 Command, June 4, 1986.
- 16. Kaminskis A. A determination of erythrocyte acetylcholinesterase activity in pyridostigmine-inhibited human blood. Technical Report, Analytical Chemistry Branch, U. S. Army

- Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, Maryland.
- 17. Metzler CM and Weiner DL. PCNONLIN. Version 01.A.

 Statistical Consultants, Inc., Lexington, Kentucky, 1985.
- 18. Sheiner LB, Stanski DR, Vozeh S, et al. Simultaneous modeling of pharmacokinetics and pharmacodynamics: Application to d-tubocurarine. Clin Pharmacol Ther 1979; 25: 358-371.
- 19. Holford NHG and Sheiner LB. Understanding the dose-effect relationship. Clin Pharmacokin 1981; 6: 429-453.
- 20. Sheiner LB. Modeling pharmacodynamics: Parametric and nonparametric approaches, in <u>Variability in Drug Therapy</u>:

 <u>Description</u>, <u>Estimation</u>, <u>and Control</u>, Rowland M, et al., eds., Raven Press, New York, 1985, pp. 139.
- 21. Watts P and Wilkinson RG. The interaction of carbamates with acetylcholinesterase. Biochem Pharmacol 1977;
 26: 757-761.
- 22. Cheng J, Lin ET, et al. Routine analysis of pyridostigmine urine samples obtained from clinical protocol titled "Bioavailability of Oral Pyridostigmine and Inhibition of

- Red Blood Cell Acetylcholinesterase by Oral and Intravenous Pyridostigmine." Report PYR/U 86-3B, 1986.
- 23. Cockcroft DW and Gault MH. Prediction of creatinine clearance from serum creatinine. Nephron 1976; 16: 31-41.
- 24. Aquilonius S-M, Eckernas S-A, Hartvig P, et al. Pharmaco-kinetics and oral bioavailability of pyridostigmine in man.

 Eur J Clin Pharmacol 1980; 18: 423-428.
- 25. Calvey TN, Chan K, Dehghan A, Williams NE. Kinetics of intravenous pyridostigmine in man. Br J Clin Pharmacol 1981; 11: 11-13.
- 26. Nowell PT, Scott CA and Wilson A. Determination of neostigmine and pyridostigmine in the urine of patients with myasthenia gravis. Br J Pharmacol 1962; 18: 617-624.
- 27. Somani SM, Roberts JB and Wilson A. Pyridostigmine metabolism in man. Clin Pharmacol Ther 1972; 13: 393-399.
- 28. Chan K and Calvey TN. Renal clearance of pyridostigmine in patients with myasthenia gravis. Eur Neurol 1977; 16: 69-72.

- 29. Rennick BR. Renal tubule transport of organic cations. Am J Physiol 1981; 240: F83-89.
- 30. Breyer-Pfaff U, Maier U, Brinkman AM, Schumm F. Pyridostigmine kinetics in healthy subjects and patients with myasthenia gravis. Clin Pharmacol Ther 1985; 37: 495-501.
- 31. Wilson IB, Harrison MA, Ginsburg S. Carbamyl derivatives of acetylcholinesterase. J Biol Chem 1961; 236: 1498-1500.

Table 1

Sample Times with Measured Plasma Concentrations
Below the Minimum Detectable Level Which were
Not Included as Data Points for the Pharmacokinetic
Curve Fitting¹

Subject	IV Dosing	PO Dosing
8	2.0	
9	0.33, 0.917	
11	0.167, 0.25	
12		3.0
13	0.417	5.0
14	0.33	10.0
16	2.5	2.0, 3.5
17	0.167	
19	0.417, 2.5	0.5, 6.0
	3.0, 3.5, 4.0	
21	0.83, 1.33	
22		7.0
23	0.083, 0.417 2.0	
24	2.5	

¹All sample times are in hours after dosing.

Table 2

Equations To Which the Plasma Pyridostigmine Concentrations Were Fit

Constants:

IVD intravenous dose

OD oral dose

TI duration of intravenous infusion

Parameters to be estimated:

V volume of the central compartment

K10 rate constant from central compartment to outside

K12 rate constant from central to peripheral compartment

K21 rate constant from peripheral to central compartment

K01 absorption rate constant, from outside to central compartment

AVA availability of oral dose

TL time lag for absorption to begin

C(T) is the concentration of pyridostigmine at time T

For the intravenous dose:

$$C(T) = A(e^{-alpha*T} - e^{-alpha*Ts}) + B(e^{-beta*T} - e^{-beta*Ts})$$
where

$$Ts = T - TI$$
 for $T > TI$ and $Ts = 0$ for $T < or = TI$

alpha and beta (alpha > beta) are the positive and negative roots of the quadratic:

$$r*r + (K12 + K21 + K10)*r + K21*K10 = 0$$

For the oral dose:

$$C(T) = Ae^{-alpha*To} + Be^{-beta*To} + Ce^{-K01*To}$$

where

$$To = T - TL$$
 for $T > TL$ and $To = 0$ for $T < or = TL$

A = (AVA*OD/V)*KO1*(K21 - alpha)/(alpha - beta)/(alpha - K01)

B = -1*(AVA*OD/V)*KO1*(K21 - beta)/(alpha - beta)/(beta - KO1)

C = (AVA*OD/V)*KO1*(K21 - KO1)/(beta - KO1)/(alpha - KO1)

alpha and beta (alpha > beta) are the positive and negative roots of the same quadratic as used for the intravenous dose

Table 3

Vital Statistics of Subjects Receiving Pyridostigmine

SUB #	WT kg	Scr* mg/ml	AGE yr	EST CLCr** ml/min
1 2 3 4 5 6	65.7 64.4 89.9 63.5 60.3 52.2	.97 1.13 .87 1.00 .93	35 25 27 25 22 26	92.94 116.23 122.75 101.42 91.91 74.39
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	80.5 75.9 76.5 77.0 77.1 79.2 62.7 86.5 79.2 104.8 65.0 78.8 61.6 76.2 72.3 70.1 83.0 68.0	.97 .97 1.30 .97 1.23 1.15 .97 .97 1.10 1.03 .97 1.10 1.13 1.10	28 35 30 31 28 29 30 24 21 27 22 22 24 19 24 20 32	121.47 107.37 151.94 113.07 147.52 140.41 92.92 135.18 143.99 164.48 109.72 125.27 111.05 138.73 133.65 116.33 145.25 96.90
AVG 07-2 STD DEV 07-2		1.05	26.00 4.61	127.51 20.06

^{*} Scr = serum creatinine

^{**} Calculated from the equation of Cockroft and Gault $Cl_{Cr} = ((140 - age)*(wt in kg))/72*S_{Cr}$

Table 4
Sampling times During and Following Intravenous Pyridostigmine

						SCHEDUL	ED TIME	OF SAME	PLE					
	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
SUBJ#						ACTUAL	TIME OF	SAMPLE	(HRS)					
7	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
8	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.867	0.750	0.833	0.867	1.000	1.333
9	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.600	0.667	0.750	0.833	0.917	1.000	1.333
10	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
11	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
12	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
13	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
14	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.887	0.750	0.833	0.917	1.000	1.333
15	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.867	0.750	0.833	0.917	1.000	1.333
16	0.000	0.083	0.167	0.250	0.333	0.417	0.483	0.583	0.667	0.750	0.833	0.917	1.000	1.333
17	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1,000	1.333
18	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
19	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
20	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
21	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
22	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
23	0.000	0.083	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
24	0.000	0.150	0.167	0.250	0.333	0.417	0.500	0.583	0.667	0.750	0.833	0.917	1.000	1.333
			-,									-,		1.000
#	18	18	18	18	18	18	18	18	18	18	18	18	18	18
MEAN	0.000	0.087	0.167	0.250	0.333	0.417	0.499	0.584	0.667	0.750	0.833	0.914	1.000	1.333
SD	0.000	0.016	0.000	0.000	0.000	0.000	0.004	0.004	0.000	0.000	0.000	0.012	0.000	0.000
CV(Z)		18,210	0.000	0.000	0.000	0.000	0.803	0.686	0.000	0.000	0.000	1.289	0.000	0.000
MAX	0.000	0.150	0.167	0.250	0.333	0.417	0.500	0.600	0.667	0.750	0.833	0.917	1.000	1.333
MIN	0.000		0.167	0.250	0.333	0.417	0.483	0.583	0.667	0.750	0.833	0.867	1.000	1.333
	0.000													
	0.000													
	0.000						ED TIME		PLE					
	1.667		2.500	3.000						8.000	10.000	12.000	24.000	
SUBJ#						SCHEDUI	ED TIME	OF SAM	7.000	8.000		12.000	24.000	
SUBJ#		2.000				SCHEDUI 4.000 ACTUAL	ED TIME 5.000	OF SAM	7.000					
	1.667	2.000	2.500	3.000	3.500	SCHEDUI 4.000 ACTUAL	ED TIME 5.000 TIME OF	OF SAM 6.000 SAMPLE	7.000 (HRS)	8.000	10.000	12.000	24.000	
7	1.667 1.667	2.000 2.000 2.000	2.500 2.500	3.000	3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000	ED TIME 5.000 TIME OF 5.000	OF SAM 6.000 SAMPLE 6.000	7.000 (HRS) 7.000	8.000 8.000	10.000	12.000 12.000	24.000 24.000	
7 8	1.667 1.667 1.667	2.000 2.000 2.000 2.000	2.500 2.500 2.500	3.000 3.000 3.000	3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000	ED TIME 5.000 TIME OF 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000	7.000 (HRS) 7.000 7.000	8.000 8.000 8.000	10.000 10.000 10.000	12.000 12.000 12.000	24.000 24.000 24.000	
7 8 9	1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000	8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000	
7 8 9 10 11	1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12	1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13	1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000	OF SAMPLE 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 5.967	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13	1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 5.967 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	.ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAMPLE 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 5.967	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAMM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 5.967 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883	
7 8 9 10 11 12 13 14 15 16 17 18 19	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 SAMPLE 6.000 6.000 6.000 6.000 5.967 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950	
7 8 9 10 11 12 13 14 15 16 17 18 19 20	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.3883 23.883 23.895 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.850 24.000 24.000 23.4000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.850 24.000 24.000 23.4000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000	7.000 (HRS) 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6,000 SAMPLE 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 6,000 8,000 8,000 8,000 18	7.000 (HRS) 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950 24.000 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 #MEAN SD	1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000	ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 8.000 8.000 8.000 8.000 8.000 8.000	7.000 (HRS) 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000	12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000 12.000	24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 24.000 23.883 23.950 24.000 24.000 24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 #MEAN SD CV(Z)	1.667 1.667	2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000	2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 4.000 6.000 6.000 6.000 6.000	ED TIME 5.000 TIME OF 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000 5.000	OF SAME 6.000 5.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 8.000 8.000 8.000 8.000 8.000	7.000 (HRS) 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.000	12.000 12.000	24.000 24.000	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 #MEAN SD	1.667 1.667	2.000 2.000	2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500 2.500	3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000 3.000	3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500 3.500	SCHEDUI 4.000 ACTUAL 4.000	ED TIME 5.000 TIME OF 5.000	OF SAM 6.000 SAMPLE 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 6.000 8.000 8.000 8.000 8.000 8.000 8.000	7.000 (HRS) 7.000	8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000 8.000	10.000 10.000	12.000 12.000	24.000 24.000	

Table 5 $\hbox{ Concentration of Pyridostigmine During and Following Intrevenous Pyridostigmine} {\color{blue}1,2} \\$

					SCHEDUL	ED TIME	OF SAM	PLE						
	0.00	0.08	0.17	0.25	0.33	0.42	0.50	0.58	0.67	0.75	0.83	0.92	1.00	1.33
SUBJ#					PYRIDOS	TIGMINE		TRATION						
7	0.00	8.51	13.80	18.40	15.60	16.30	17.90	13.00	10.10	9.95	7.65	6.16	6.34	4.14
8	0.00	9.69	20.90	18.90	14.50	13.50	11.70	10.80	8.15	5.82	5.26	4.34	3.51	3.37
9	0.00	4.75	8.73	13.60	25.90	18.30	20.30	14.00	9.24	9.19	6.29	2.63	6.27	4.04
10	0.00	2.27	5.23	3.94	6,06	9.81	12.40	7.46	5.32	4.78	3.68	2.99	2.81	1.62
11 12	0.00	5.11 3.37	10.40	3.37	19.20 30.90	23.30 28.10	27.80 22.10	25.70 16.30	19.50	13.90 17.30	12.70 8.30	9.38	8.65	6.10
13	0.00	8.78	11.70	25.00	32.50	19.70	46.80	15.20	12.80 14.10	11.20	8.34	8.30 6.12	9.06 7.06	6.17
14	0.00	11.60	17.70	23.10	19.00	33.30	24.50	20.20	11.30	6.59	6.70	4.07	4.42	3.04
15	0.00	8.11	13,90	18.20	16.40	14.00	18.00	12.50	8.67	8.96	5.37	3.86	5.30	2.25
16	0.00	7.83	10.90	14.20	14.60	19.40	21.00	11.00	8.64	10.70	7.20	5.83	5.85	2.63
17	0.00	8.87	6.17	12.90	14.20	16.20	20.70	10.10	8.92	6.16	5.22	3.82	3.10	2.29
18	0.00	6.90	10.50	12.30	12.90	20.60	18.90	14.50	10.50	5.67	5.50	4.61	4.37	3.19
19	0.00	10.80	17.10	17.50	18.60	14.20	24.20	17.60	10.10	8.42	6.93	6.80	7.12	5.07
20	0.00	9.09	12.60	20.50	19.90	26.80	29.50	18.10	16.80	10.30	6.98	6.26	5.18	5.79
21	0.00	19.60	16.30	20.70	21.70	19.00	22.70	13.70	10.50	7.54	7.02	5.66	3.86	5.84
22	0.00	7.80	14.90	22.40	21.80	24.80	20.60	11.60	11.30	8.55	6.59	4.80	6.36	4.09
23	0.00	12.30	8.39	16.70	22.60	14.90	20.50	12.70	11.00	14.80	9.26	7.74	3.86	4.24
24	0.00	9.91	11.20	13.00	19.20	22.30	20.80	17.00	12.40	9.04	6.98	6.77	5.37	3.19
*	18	18	17	18	18	18	18	18	18	18	18	18	18	18
MEAN	0.00	8.63	12.38	16.39	19.20	19.68	22.24	14.53	11.07	9.38	7.00	5.56	5.47	3.86
SD	0.00	3.84	4.18	5.93	6.35	5.96	7.54	4.23	3.27	3.34	1.95	1.84	1.79	1.44
CV(I)	0.00	44.50	33.78	36.20	33.08	30.30	33.91	29.12	29.52	35.62	27.82	33.03	32.79	37.14
MAX	0.00	19.60	20.90	25.00	32.50	33.30	46.80	25.70	19.50	17.30	12.70	9.38	9.06	6.17
MIN	0.00	2.27	5.23	3.37	6.06	9.61	11.70	7.46	5.32	4.78	3,68	2.63	2.81	1.62
						SCHEDUI	ED TIME	OF SAM	PLE					
	1.67	2.00	2.50	3.00	3.50	SCHEDUI 4.00	ED TIME 5.00	OF SAM	PLE 7.00	8.00	10.00	12.00	24.00	
SUBJ#	1.67	2.00	2.50	3.00			5.00	6.00	7.00	8.00	10.00	12.00	24.00	
7	2.86	2.00	2.71	3.00 1.95	PYRIDO: 2.15	4.00 STIGMINE 0.00	5.00 CONCEN 0.00	6.00	7.00 ng/ml 0.00	8.00 0.00	10.00	12.00	24.00	
7 8	2.86 2.75	2.58	2.71 1.59	1.95 0.00	2.15 0.00	4.00 STIGMINE 0.00 0.00	5.00 CONCEN 0.00 0.00	6.00 TRATION 0.00 0.00	7.00 ng/ml 0.00 0.00	0.00	0.00	0.00	0.00	
7 8 9	2.86 2.75 1.81	2.58 0.00 2.38	2.71 1.59 2.21	1.95 0.00 0.00	2.15 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00	0.00 0.00 0.00	0.00 0.00 0.00	0.00 0.00 0.00	0.00	
7 8 9 10	2.86 2.75 1.81 0.00	2.58 0.00 2.38 0.00	2.71 1.59 2.21 0.00	1.95 0.00 0.00 0.00	PYRIDO: 2.15 0.00 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00	
7 8 9 10 11	2.86 2.75 1.81 0.00 6.07	2.58 0.00 2.38 0.00 4.07	2.71 1.59 2.21 0.00 3.31	1.95 0.00 0.00 0.00 3.86	PYRIDOS 2.15 0.00 0.00 0.00 3.23	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01	6.00 TRATION 0.00 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	
7 8 9 10 11	2.86 2.75 1.81 0.00 6.07 4.66	2.58 0.00 2.38 0.00 4.07 3.31	2.71 1.59 2.21 0.00 3.31 2.53	1.95 0.00 0.00 0.00 3.86 1.74	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01	6.00 TRATION 0.00 0.00 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	
7 8 9 10 11 12	2.86 2.75 1.81 0.00 6.07 4.66 2.34	2.58 0.00 2.38 0.00 4.07 3.31 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00	1.95 0.00 0.00 0.00 3.86 1.74 0.00	PYRIDO: 2.15 0.00 0.00 0.00 3.23 1.57 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62	1.95 0.00 0.00 0.00 3.86 1.74 0.00	PYRIDO: 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07	PYRIDO: 2.15 0.00 0.00 3.23 1.57 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14 15	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07	PYRIDOS 2.15 0.00 0.00 3.23 1.57 0.00 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14 15 16	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07	PYRIDO: 2.15 0.00 0.00 3.23 1.57 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14 15	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05	1.95 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00	PYRIDO: 2.15 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14 15 16 17 18	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00	1.95 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59	1.95 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 0.00	PYRIDOS 2.15 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 3.71	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 2.01 0.00 0.00 0.00	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00 1.81 3.87 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 0.00 3.66 0.00 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00 1.81 3.87 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 0.00 3.66 0.00 0.00	PYRIDO: 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00 1.81 3.87 0.00	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 0.00 3.66 0.00 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 3.66 0.00 0.00 1.73 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 3.86 1.74 0.00 0.00 0.00 0.00 3.66 0.00 0.00 1.73 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 1.93	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 3.66 0.00 0.00 1.73 0.00	PYRIDOS 2.15 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 2.70 0.00 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 3.66 0.00 0.00 1.73 0.00 1.57	PYRIDOS 2.15 0.00 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 MEAN SD	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 3.66 0.00 0.00 1.73 0.00 1.57	PYRIDOS 2.15 0.00 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 MEAN SD CV(I)	2.86 2.75 1.81 0.00 6.07 4.66 2.34 2.92 1.97 1.67 1.71 2.34 3.16 3.99 2.41 3.66 3.80 3.77	2.58 0.00 2.38 0.00 4.07 3.31 0.00 2.04 2.63 0.00 0.00 1.81 3.87 0.00 1.75 0.00 4.40	2.71 1.59 2.21 0.00 3.31 2.53 0.00 1.62 1.83 2.05 0.00 0.00 8.59 1.70 0.00 2.76 2.92 4.64	1.95 0.00 0.00 0.00 3.86 1.74 0.00 0.00 2.07 0.00 0.00 3.66 0.00 0.00 1.73 0.00 1.57	PYRIDOS 2.15 0.00 0.00 0.00 0.00 3.23 1.57 0.00 0.00 0.00 0.00 0.00 0.00 0.00 1.93	4.00 STIGMINE 0.00 0.00 0.00 0.00 2.70 0.00 0.00 0.00	5.00 CONCEN 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	6.00 TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	7.00 ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	

Sample was not obtained at the scheduled time. See Table 4 for the time of sample.

Concentrations are nanograms of pyridostigmine base/ml

^{1 2} Concentrations less than 1.5 ng/ml, the lower limit of dectability of the assey are denoted 0.00

Table 6
Sampling Times Following Oral Pyridostigmine

				SCHEDUL	ED TIME	OF SAME	LE			
	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
SUBJ#				ACTUAL	TIME OF	SAMPLE	(HRS)			
7	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
8	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
9	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
10	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
11	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
12	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
13	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
14	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
15	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
16	0.00		0.50	0.75						
17	0.00				1.00	1.33	1.67	2.00	2.50	3.00
			0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
18	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
19	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
20	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
21	0.00		0.50	0.73	1.00	1.33	1.67	2.00	2.50	3.00
22	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
23	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
24	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
140	5.50									
#	18		18	18	18	18	18	18	18	18
MEAN	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
SD	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CV(%)	0.00		0.00	0.53	0.00	0.00	0.00	0.00	0.00	0.00
MAX	0.00		0.50	0.75	1.00	1.33	1.67	2.00	2.50	3.00
MIN										3.00
	0.00	0.25	0.50	0.73	1.00	1.33	1.67	2.00	2.50	0.00
	0.00	0.25	0.50	0.73	1.00	1.33	1.6/	2.00	2.50	0.00
	0.00	0.25	0.50					2.00	2.50	0.00
				SCHEDUI	ED TIME	OF SAME	PLE			0.00
	3.50		5.00	SCHEDUT 6.00	ED TIME 7.00	OF SAM	PLE 10.00	12.00	24.00	3.33
SUBJ#	3.50	4.00	5.00	SCHEDUI 6.00 ACTUAL	ED TIME 7.00 TIME OF	OF SAM 8.00 SAMPLE	PLE 10.00 (HRS)	12.00	24.00	3.33
SUBJ#	3.50	4.00	5.00	SCHEDUI 6.00 ACTUAL 6.00	ED TIME 7.00 TIME OF 7.00	OF SAM 8.00 SAMPLE 8.00	PLE 10.00 (HRS) 10.00	12.00 12.00	24.00	3.33
SUBJ# 7 8	3.50 3.50 3.50	4.00 4.00 4.00	5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00	7.00 TIME OF 7.00 7.00	OF SAME 8.00 SAMPLE 8.00 8.00	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00	24.00 24.00 24.00	3.00
SUBJ# 7 8 9	3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00	.ED TIME 7.00 TIME OF 7.00 7.00 7.13	OF SAM 8.00 SAMPLE 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00	12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00	3.00
SUBJ# 7 8 9 10	3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00	JED TIME 7.00 TIME OF 7.00 7.00 7.13 7.00	OF SAM 8.00 SAMPLE 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37	0.00
SUBJ# 7 8 9 10 11	3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00	7.00 TIME OF 7.00 7.00 7.00 7.13 7.00 7.00	OF SAM 8.00 SAMPLE 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00	3.00
SUBJ# 7 8 9 10 11 12	3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00	.ED TIME 7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00	OF SAM 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37	0.00
SUBJ# 7 8 9 10 11 12 13	3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00	.ED TIME 7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00 7.00	OF SAMPLE 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12	3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00	.ED TIME 7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00	OF SAM 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13	3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00	.ED TIME 7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00 7.00	OF SAMPLE 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14	3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUT 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00	7.00 TIME OF 7.00 7.00 7.00 7.13 7.00 7.00 7.00 7.00 7.00	OF SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAM 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAM 8.00 SAMPLE 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME 0F 7.00 7.00 7.13 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME 0F 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.37 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME 0F 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.00	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.00	PLE 10.00 (HRS) 10.00	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 4.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	JED TIME 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 #MEAN	3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	4.00 6.00 4.00 6.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 MEAN SD	3.50 3.50	4.00 6.00 6.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.13 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	
SUBJ# 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 MEAN SD CV(%)	3.50 3.50	4.00 6.00 6.00	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00	SCHEDUI 6.00 ACTUAL 6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.0	7.00 TIME OF 7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.0	OF SAME 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	PLE 10.00 (HRS) 10.00 10	12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00	24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00 24.00	

Table 7 Plasma Concentration of Pyridostigmina Following Oral Pyridostigmina 1,2

				SCHED	ULED II	ME OF S	AMPLE			
	0.00	0.25	0.50	0.75	1.00	1.33	1.67		2.50	3.00
SUBJ#							TRATION	ng/ml		
7	0.00	0.00	6.43	8.24	8.33	8.69	12.30	9.72	9.63	13.80
8	0.00	0.00	2.21	6.44	5.14	4.72	6.56	8.03	7.15	12.20
9	0.00	0.00	2.27	3.95	4.01	5.46	6.01	4.91	4.76	4.21
10	0.00	0.00	0.00	2.33	2.66	3.28	3.56	4.57	4.54	7.26
11	0.00	5.11	12.70	17.20	19.30	23.70	19.90	16.30	14.60	
12	0.00	0.00	5.67	4.49	6.28	7.04	11.70	13.90	13.20	8.75
13	0.00	0.00	7.61		12.80	16.80	14.40	14.30	16.70	13.50
14	0.00	3.51	4.98		14.50	10.00	9.07	16.20	20.90	
15	0.00	0.00	0.30	5.14	7.83		14.60	12.30	10.90	13.50
16	0.00	0.00	0.00	0.00	0.00	1.67	3.75	10.20	5.90	7.96
17	0.00	0.00	7.75		11.10	13.80	10.20	10.50	9.42	8.37
18	0.00	0.00	8.04			11.80	11.90	7.74	6.01	5.26
19	0.00	4.16		8.17	13.30	11.00	8.45	8.39	6.22	5.04
20	0.00	0.00		8.17	6.30	9.86		13.60	10.30	12.10
21	0.00	1.93			11.50	9.74	10.10	10.90	5.56	7.35
22	0.00	2.81	6.13		11.40	9.23	10.10	7.57	8.17	8.17
23	0.00	4.80	7.77	10.90	14.00	16.10	15.30	13.90	13.40	12.30
24	0.00	2.00	10,60	10.90	11.60	10.00	13.00	10.60	9.38	9.98
#	18	18	18	18	18	18	18	18	18	18
MEAN	0.00	1.35	6.01	8.96	9.71	10.21	10.83	10.76		9.92
SD	0.00	1.90	4.24		4.97	5.22		3.51		3.42
CV(I)		140.82			51.13	51.17	38.97	32.62	45.77	34.51
MAX	0.00	5.11			19.30	23.70	19.90		20.90	14.80
MIN	0.00	0.00	0.00	0.00	0.00	1.67	3.56	4.57	4.54	4.21
				COURDIN	ED ED#	OF 544	··· -			
	3 50	4 00	5 00		ED TIME			12.00	27, 00	
CITO 14	3.50	4.00	5.00	6.00	7.00	8.00	10.00	12.00	24.00	
SUBJ#				6.00 PYRIDOS	7.00 TIGMINE	8.00 CONCEN	10.00 TRATION	ng/ml		
7	10.90	8.48	5.08	6.00 PYRIDOS 2.69	7.00 TIGMINE 4.25	8,00 CONCEN 0,00	10.00 TRATION 0.00	ng/ml 0.00	0.00	
7 8	10.90 9.30	8.48 9.24	5.08 4.96	6.00 PYRIDOS 2.69 3.22	7.00 TIGMINE 4.25 2.24	8.00 CONCEN 0.00 1.80	10.00 TRATION 0.00 0.00	ng/ml 0.00 0.00	0.00	
7 8 9	10.90 9.30 5.00	8.48 9.24 6.19	5.08 4.96 6.65	6.00 PYRIDOS 2.69 3.22 2.11	7.00 TIGMINE 4.25 2.24 1.75	8.00 CONCEN 0.00 1.80 0.00	10.00 TRATION 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00	0.00	
7 8 9 10	10.90 9.30 5.00 6.62	8.48 9.24 6.19 5.67	5.08 4.96 6.65 5.76	6.00 PYRIDOS 2.69 3.22 2.11 2.01	7.00 TIGMINE 4.25 2.24 1.75* 2.21	8.00 CONCEN 0.00 1.80 0.00 0.00	10.00 TRATION 0.00 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00	0.00 0.00 0.00	
7 8 9 10 11	10.90 9.30 5.00 6.62 9.84	8.48 9.24 6.19 5.67 11.90	5.08 4.96 6.65 5.76 9.18	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99	7.00 TIGMINE 4.25 2.24 1.75 2.21 6.88	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08	10.00 TRATION 0.00 0.00 0.00 0.00 2.04	ng/ml 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	
7 8 9 10 11 12	10.90 9.30 5.00 6.62 9.84 12.20	8.48 9.24 6.19 5.67 11.90 10.20	5.08 4.96 6.65 5.76 9.18 6.62	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10	7.00 STIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50	10.00 ITRATION 0.00 0.00 0.00 0.00 2.04 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13	10.90 9.30 5.00 6.62 9.84 12.20	8.48 9.24 6.19 5.67 11.90 10.20 11.10	5.08 4.96 6.65 5.76 9.18 6.62 16.20	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55	7.00 STIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70 3.17	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08	10.00 ITRATION 0.00 0.00 0.00 0.00 2.04 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13	10.90 9.30 5.00 6.62 9.84 12.20 13.40	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21	10.00 TRATION 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21	10.00 ITRATION 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	
7 8 9 10 11 12 13 14 15	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28	7.00 TIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70 3.17 6.45 2.56 4.61	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57	10.00 ITRATION 0.00 0.00 0.00 2.04 0.00 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 1.92 0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00	8.00 CONCEN 0.00 1.80 0.00 5.08 2.50 4.21 0.00 3.57 0.00	10.00 ITRATION 0.00 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00	7.00 TIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00	10.00 ITRATION 0.00 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38	5.08 4.96 6.65 5.76 9.18 6.62 16.20 11.50 7.02 4.69 3.06 3.99	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02	7.00 TIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00	10.00 ITRATION 0.00 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14	5.08 4.96 6.65 5.76 9.18 6.62 12.70 11.50 7.02 4.69 3.06 3.99 5.45	6.00 PYRIDOS 2.69 3.22 2.11 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02	7.00 TIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50	10.00 ITRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76	8.48 9.24 6.19 5.67 11.90 10.20 11.10 13.20 7.53 6.53 3.45 5.38 8.14 5.70	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33	6.00 PYRIDOS 2.69 3.22 2.11 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00	10.00 (TRATION 0.00 0.00 0.00 0.00 2.04 0.00 0.00 0.00	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33	6.00 PYRIDOS 2.69 3.22 2.11 7.99 4.10 5.55 7.59 6.22 0.00 7.02 4.07 3.90 3.56	7.00 STIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34	8.00 CONCEN 0.00 1.80 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.233 4.77 7.46	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89	7.00 TIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60	8.00 CONCEN 0.00 1.80 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33	6.00 PYRIDOS 2.69 3.22 2.11 7.99 4.10 5.55 7.59 6.22 0.00 7.02 4.07 3.90 3.56	7.00 STIGMINE 4.25 2.24 1.75* 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34	8.00 CONCEN 0.00 1.80 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41	7.00 CTIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 0.1.93 5.29 2.13	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46 4.62	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41	7.00 CTIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29 2.13	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46 4.62	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41	7.00 FIGHINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29 2.13	10.00 ITRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46 4.62	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41 18 4.34 2.11	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29 2.13	10.00 ITRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 ******************************	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46 4.62 18 6.78 3.56 52.56	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41 18 4.34 2.11 48.65	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29 2.13	10.00 (TRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.90 9.30 5.00 6.62 9.84 12.20 13.40 13.90 12.00 5.30 6.28 4.28 3.74 12.10 3.76 7.67 13.10 9.04	8.48 9.24 6.19 5.67 11.90 10.20 11.10 11.60 13.20 7.53 6.53 3.45 5.38 8.14 5.70 5.91 8.57 9.45	5.08 4.96 6.65 5.76 9.18 6.62 16.20 12.70 11.50 7.02 4.69 3.06 3.99 5.45 2.33 4.77 7.46 4.62	6.00 PYRIDOS 2.69 3.22 2.11 2.01 7.99 4.10 5.55 7.59 6.22 6.28 3.52 0.00 7.02 4.07 3.90 3.56 4.89 3.41 18 4.34 2.11	7.00 STIGMINE 4.25 2.24 1.75 2.21 6.88 2.70 3.17 6.45 2.56 4.61 0.00 0.00 2.59 5.90 3.34 0.00 4.60 2.63	8.00 CONCEN 0.00 1.80 0.00 0.00 5.08 2.50 4.21 0.00 3.57 0.00 0.00 1.78 5.50 0.00 1.93 5.29 2.13	10.00 ITRATION 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	ng/ml 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	

Sampla was not obtained at the acheduled time. Same Table 6 for the time of sampla. Concentrations are nanograms of pyridostigmine base/ml

¹ Concentrations less than 1.5 ng/ml, the lower limit of dectability of the assay are denoted 0.00

Table 8

Indicators of "Goodness of Fit" for the Pharmacokinetic Curve Fitting

	IV		PO	
SUB #	SUM of WEIGHTED	CORREL	SUM of WEIGHTED	CORREL
	SQUARED RESIDUALS	5	SQUARED RESIDUAL	s
7	3.134	.976	4.221	.924
8	2.192	.985	7.625	.846
9.	1.664	.989	3.591	.878
10	1.803	.961	3.990	.822
11	2.425	.989	2.718	.973
12	7.669	.967	7.514	.888
13	13.916	.928	2.402	.971
14	3.366	.987	8.474	.891
15	4.277	.967	4.609	.948
16	1.604	.983	2.253	.959
17	1.375	.984	.652	.984
18	1.987	.974	.648	.990
19	1.551	.987	4.088	.908
20	3.677	.981	6.431	.911
21	1.348	.988	4.971	.904
22	2.351	.988	2.633	.934
23	5.649	.933	2.745	.971
24	2.098	.988	2.294	.971
AVG	3.449	.975	3.992	.926
STD DEV	3.091	.018	2.288	.049

Table 9
Estimates of the Pharmacokinetic Microconstants for Pyridostigmine

SUB	#	VOL L	K10 hr ⁻¹	K12 hr ⁻¹	K21 hr ⁻¹	K01 hr-1	BIOAVA	TLAG hr
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24			3.150 1.535 2.077 2.109 1.106 1.819 3.707 3.064 3.394 9.994 5.649 1.985 4.804 2.915 5.959 2.081 2.180 2.000	1.483 1.655 4.141 24.961 4.655 2.849 5.261 2.618 4.995 2.201	3.046 1.543 3.263 .731	.367 .284 .236 .193 .580 .292 .268 .188 .306 .149 .310 1.441 .334 .202 .272 .503 .298 .495	.265 .194 .393 .250 .248 .341 .511 .371 .387 .354 .147 .213 .333 .244 .186	.347 .391 .359 .596 .174 .341 .388 .202 .673 1.305 .332 .422 .151 .346 .223 .229 .145 .213
AVG		17.465	3.307			.373		.380
STD I	E.	V 8.930	2.159	5.405	1.690	.290	.094	.272

Table 10 Estimates of the Pharmacokinetic Macroconstants for Pyridostigmine

SUB #	ALPHA hr ⁻¹	BETA hr ⁻¹	A(IV) ng/ml	B(IV) ng/ml	A(PO) ng/ml	B(PO) ng/ml	C(PO) ng/ml
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10.223 3.208 4.444 5.386 2.950 4.528 5.618 5.128 8.360 40.848 12.072 5.210 11.880 6.370 12.683 4.688 5.855 5.042	.488 .596 .219 .687 .830 .607 .564 1.909 1.555 .231 1.232 .706 1.533 .324 .739	-34.992 -15.358 -6.399 -26.573 -18.174 -27.758 -29.098 -13.484 -5.909 -9.726 -16.513 -12.689 -18.533 -12.953 -24.370 -12.401	-22.867 -13.867 -55.226 -43.529 -18.440 -24.435 -23.667 -22.550 -15.260 -30.374 -25.133 -28.984 -19.682 -36.178 -31.626	-5.645 -4.161 -2.812 -26.227 -6.169 -14.972 -12.759 -9.336 -1.812 -6.757 -32.303 -5.154 -7.083 -5.417 -12.111 -7.523	-24.042 -13.974 -14.951 -34.505 -7.473 -12.900 8.219 -13.611 -15.037 -9.791 29.152	15.198 -16.965 37.962
						,	
AVG	8.583 8.610		-17.443 8.116	-30.104	-9.918 7.940		17.530

Table 11

Estimates of the Area Under the Concentration-Time Curve and Clearance of Pyridostigmine

SUB	#	AUC(IV) ng-hr/ml		DOSE(IV)	DOSE(PO) ug	CL ml/min	CL ml/min/kg	CL ml/ml GFR
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24		22.05 16.77 19.11 10.13 40.90 25.73 23.10 22.32 18.58 14.23 12.49 23.44 18.91 23.76 16.32 25.25 22.01 24.80	62.14 52.02 41.54 44.45 11.81 67.06 88.70 120.34 80.98 58.36 54.63 46.02 44.70 86.86 49.13 53.39 84.95 62.50	900.73 860.66 922.14 918.63 938.26 980.98 915.16 962.67 866.09 950.27 838.53 776.69 927.57 936.95 840.21 910.13 958.80 823.07	9995.15 10095.00 10328.41 10248.01 10257.15 10300.24 10309.44 10158.16 10187.92 10065.18 10361.55 10391.74 10272.74 10272.74 10291.64 10355.23 10320.92 10412.92 10231.68	855.61 804.11 1510.95 382.34 635.43 660.32 718.74 777.07 1113.07 1118.67 552.16 817.49 657.29 858.22 600.77	8.46 11.27 10.51 19.62 4.96 8.02 10.53 8.31 9.81 10.62 17.21 7.01 13.27 8.63 11.87 8.57 8.75 8.14	
AVO		21.11 V 6.70	67.20 23.43	901.53 55.50	10254.62 115.06	779.01 258.29	10.31	6.27 2.42

Table 12

Amounts of Pyridostigmine Base Recovered in the Urine After Intravenous and Oral Administration of Pyridostigmine Bromide (micrograms)

MIN	158 0 0 4 4 0 0	NEW	18 161 111 15 4	501
MAX	403 272 102 85 23 279 788	MAX	235 420 602 382 183 395	2056
SD CV(%)	36 79 71 101 114 157 33		68 33 55 81 75	43
SD (89 82 36 27 8 94	SD (73 84 154 129 63	471
MEAN	248 104 50 27 7 60	MEAN	108 255 280 159 84 206	1001
24	NA 403 67 50 27 9 0	24	NA 32 319 242 155 117 340	1207
23	NA 158 78 36 19 10 25	23	NA 18 265 372 254 52 229	1189
22	NA 188 69 52 14 2 0	22	NA 235 267 199 97 58 88	943
SUBJECT 20 21	NA 187 68 30 4 1 90 379	SUBJECT 20 21	NA 85 220 111 15 4	501
SUI 20	NA 232 232 116 0 85 23 53 53	20	NA 62 207 305 261 183 245	1263
19	NA 366 0 101 42 0 279 788	19	NA 168 161 256 68 125 96	873
18	NA 241 162 29 4 4 1 0	18	NA 126 185 149 37 9	869
17	NA 211 272 102 16 7 32 640	17	NA 134 420 602 382 123 395	2056
COTTE	STED 17	COTTE	222222	STED
SCHEDULED ROUTE INTERVAL	BLANK 0.00-1.00 1.00-2.00 2.00-4.00 1.00-6.00 6.00-8.00 1.00-24.0 IOTAL EXCRETED	SCHEDULED ROUTE	BIANK 0.00-1.00 1.00-2.00 2.00-4.00 4.00-6.00 6.00-8.00 8.00-24.0	TOTAL EXCRETED

^{1 1.32} mg of pyridostigmine bromide infused over 30 minutes

^{2 16} mg of pyridostigmine bromide administered as Mestinon^R syrup

Table 13

Recovery of Pyridostigmine in the Urine after Intravenous and Oral Administration of Pyridostigmine Bromide

	INTRAVENOUS			ORAL ADMI		
	URINARY	DOSE	8	URINARY	DOSE	8
SUBJ#	RECOVERY-ug	ug	EXCRETED	RECOVERY-ug	ug	EXCRETED
	640	0.20	76.2	2056	10262	10.0
17	640	839	76.3	2056	10362	19.8
18	438	777	56.4	698	10392	6.7
19	788	928	84.9	873	10273	8.5
20	510	937	54.4	1263	10292	12.3
21	379	840	45.1	501	10355	4.8
22	326	910	35.8	943	10321	9.1
23	324	959	33.8	1189	10413	11.4
24	556	823	67.6	1207	10232	11.8
MEAN	495	877	56.8	1091	10330	10.6
SD	163	65	18.5	471	62	4.6
CV(%)	33	7	32.6	43	1	43.1
MAX	788	959	84.9	2056	10413	19.8
MIN	324	777	. 33.8	501	10232	4.8

Table 14

Pyridostigmine Renal Clearance and Bioavailability
Calculated Using Urinary Excretion Data

SUBJ#	RENAL CLEARANCE ¹ ml/min	BIOAVAILABILITY ²
17	854	26.0 (35.4) ³
18	311	11.9 (14.7)
19	694	10.0 (21.3)
20	357	22.5 (33.3)
21	387	10.7 (24.4)
22	215	25.5 (18.6)
23	245	33.8 (35.5)
24	374	17.5 (20.3)
MEAN	430	19.7 (29.2)
SD	225	8.6 (9.4)
CV(%)	52	43.7 (32.2)
MAX	854	33.8 (35.5)
MIN	215	10.0 (14.7)

Calculated from the total clearance times the fractional excretion of unchanged pyridostigmine into the urine.

Calculated by dividing the fractional excretion of pyridostigmine into the urine after oral dosing by the fractional excretion of urine into the urine after intravenous dosing.

Values in () represent the parameter estimate of bioavailability determined from the curve fitting process (Table 9).

Table 15

Inhibition of Erythrocyte Acetylcholinesterase During and Following Intravenous Pyridostigmine

	SCHEDULED TIME OF SAMPLE											
	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	3.00	4.00	6.00	24.00
SUBJ#				INHIBIT	ION OF	ERYTHRO	CYTE AC	ETYLCHO	LINESTE	RASE		
7	0.00	0.18	0.29	0.26	0.23	0.18	0.15	0.12	0.08	0.05	0.03	-0.02
8	0.00	0.22	0.28	0.22	0.18	0.12	0.09	0.08	0.06	0.00	-0.03	-0.03
9	0.00	0.14	0.27	0.29	0.23	0.21	0.17	0.13	0.10	0.06	0.01	0.04
10	0.00	0.14	0.26	0.22	0.17	0.16	0.14	0.10	0.09	0.04	-0.02	-0.04
11	0.00	0.13	0.29	0.28	0.25	0.21	0.14	0.14	0.06	0.02	0.02	-0.03
12	0.00	0.20	0.33	0.29	0.24	0.24	0.11	0.13	0.11	0.05	0.03	-0.03
13	0.00	0.19	0.39	0.24	0.20	0.13	0.09	0.06	0.06	0.02	0.00	-0.01
14	0.00	0.20	0.32	0.25	0.18	0.14	0.10	0.06	0.06	0.01	-0.01	0.02
15	0.00	0.18	0.28	0.24	0.22	0.16	0.15	0.10	0.06	0.01	0.01	0.02
16	0.00	0.17	0.26	0.24	0.19	0.15	0.12	0.11	0.11	0.01	0.01	0.03
17	0.00	0.16	0.29	0.24	0.19	0.12	0.09	0.07	0.04	0.03	0.00	0.03
18	0.00	0.12	0.22	0.19	0.14	0.10	0.06	0.05	0.05	0.00	0.00	0.01
19	0.00	0.19	0.36	0.29	0.28	0.23	0.16	0.13	0.07	0.05	0.03	0.02
20	0.00	0.19	0.32	0.27	0.23	0.19	0.15	0.12	0.06	0.03	0.02	0.00
21	0.00	0.17	0.29	0.22	0.16	0.12	0.10	0.06	0.06	0.04	0.01	0.00
22	0.00	0.21	0.30	0.23	0.18	0.14	0.10	0.09	0.05	0.03	0.01	0.00
23	0.00	0.15	0.29	0.27	0.19	0.17	0.15	0.10	0.07	0.03	0.05	-0.01
24	0.00	0.14	0.31	0.23	0.19	0.16	0.12	0.12	0.07	0.03	0.02	-0.04
#	18	18	18	18	18	18	18	18	18	18	18	18
MEAN	0.00	0.17	0.30	0.25	0.20	0.16	0.12	0.10	0.07	0.03	0.01	0.00
SD	0.00	0.03	0.04	0.03	0.04	0.04	0.03	0.03	0.02	0.02	0.02	0.03
CV(I)	0.00	17.11	12.94	11.58	17.40	24.83	25.25	29.44	28.99	63.19	182.29	1150.45
MAX	0.00	0.22	0.39	0.29	0.28	0.24	0.17	0.14	0.11	0.06	0.05	0.04
MIN	0.00	0.12	0.22	0.19	0.14	0.10	0.06	0.05	0.04	0.00	-0.03	-0.04

Sample was not obtained at the scheduled time. See Table 4 for the time of sample.

Table 16

Inhibition of Erythrocyte Acetylcholinesterase Following Oral Pyridostigmine

						SCHEDUL	ED TIME	OF SAM	PLE					
	0.00	0.25	0.50	0.75	1.00	1.33	1.67	2.00	3.00	4.00	6.00	8.00	10.00	24.00
SUBJ#				INHIBIT	ION OF	ERYTHRO	CYTE AC	ETYLCHO	LINESTE	RASE (I)			
7	0.00	0.02	0.11	0.14	0.17	0.19	0.27	0.29	0.31	0.26	0.16	0.11	0.06	0.02
8	0.00	0.01	0.04	0.10	0.11	0.10	0.19	0.20	0.26	0.25	0.11	0.07	0.05	0.00
9	0.00	0.03	0.06	0.10	0.11	0.14	0.14	0.17	0.19	0.22	0.14	0.08	0.08	0.01
10	0.00	0.03	0.05	0.08	0.10	0.14	0.17	0.21	0.25	0.22	0.11	0.06	0.04	0.02
11	0.00	-0.01	0.12	0.22	0.31	0.34	0.34	0.29	0.29	0.25	0.17	0.10	0.05	-0.01
12	0.00	-0.02	0.07	0.11	0.14	0.18	0.23	0.27	0.30	0.25	0.11	0.07	0.02	-0.03
13	0.00	0.00	0.12	0.20	0.28	0.33	0.33	0.32	0.35	0.34	0.18	0.13	0.08	-0.02
14	0.00	-0.01	0.04	0.17	0.25	0.27	0.28	0.32	0.31	0.30	0.21	0.12	0.09	-0.04
15	0.00	0.02	0.01	0.06	0.14	0.22	0.27	0.33	0.39	0.36	0.22	0.11	0.07	-0.04
16	0.00	0.01	0.00	-0.01	-0.01	0.02	0.07	0.14	0.20	0.19	0.20	0.14	0.08	-0.01
17	0.00	-0.01	0.09	0.18	0.23	0.29	0.31	0.30	0.28	0.26	0.13	0.09	0.04	0.00
18	0.00	0.01	0.09	0.18	0.20	0.22	0.21	0.20	0.17	0.15	0.05	0.03	0.02	-0.04
19	0.00	0.10	0.20	0.26	0.31	0.30	0.27	0.25	0.21	0.17	0.12	0.07	0.08	0.01
20	0.00	0.00	0.07	0.11	0.15	0.17	0.21	0.25	0.29	0.23	0.16	0.07	0.09	0.01
21	0.00	-0.01	0.05	0.10	0.16	0.18	0.20	0.22	0.17	0.14	0.06	0.05	0.05	0.00
22	0.00	0.04	0.11	0.18	0.27	0.26	0.23	0.25	0.27	0.22	0.10	0.06	0.04	-0.01
23	0.00	0.04	0.11	0.15	0.21	0.24	0.26	0.25	0.27	0.23	0.15	0.13	0.08	0.01
24	0.00	-0.01	0.15	0.17	0.25	0.25	0.30	0.31	0.26	0.21	0.12	0.08	0.11	0.04
#	18	18	18	18	18	18	18	18	18	18	18	18	18	18
MEAN	0.00	0.01	0.08	0.14	0.19	0.21	0.24	0.25	0.27	0.24	0.14	0.09	0.06	0.00
SD	0.00	0.03	0.05	0.06	0.08	0.08	0.07	0.06	0.06	0.06	0.05	0.03	0.03	0.02
CV(Z)	0.00	213.69	60.16	46.26	44.99	38.75	29.05	21,71	22.58	24.44	34.03	35.35	40.45	513.21
MAX	0.00	0.10	0.20	0.26	0.31	0.34	0.34	0.33	0.39	0.36	0.22	0.14	0.11	0.04
MIN	0.00	-0.02	0.00	-0.01	-0.01	0.02	0.07	0.14	0.17	0.14	0.05	0.03	0.02	-0.04

Sample was not obtained at the scheduled time. See Table 6 for the time of sample.

Table 17

Duration of Inhibition of Erythrocyte Acetylcholinesterase After 1.3 mg of Pyridostigmine Bromide Intravenously

SUBJ#	EARLIEST TIME OF ACHE INHIBITION ABOVE 20%	LATEST TIME OF AChE INHIBITION ABOVE 20%	DURATION OF ACHE INHIBITION ABOVE 20%1
7	0.50	1.00	0.50
8	0.25	0.75	0.50
9	0.50	1.33	0.83
10	0.50	0.75	0.25
11	0.50	1.33	0.83
12	0.25	1.33	1.08
13	0.50	1.00	0.50
14	0.25	0.75	0.50
15	0.50	1.00	0.50
16	0.48	0.75	0.27
17	0.50	0.75	0.25
18	0.50	0.50	0.222
19	0.50	1.33	0.83
20	0.50	1.00	0.50
21	0.50	0.75	0.25
22	0.25	0.75	0.50
23	0.50	0.75	0.25
24	0.50	0.75	0.25
MEAN	0.44	0.92	0.49
SD	0.11	0.26	0.25
CV(%)	24.01	28.06	52.09
MAX	0.50	1.33	1.08
MIN	0.25	0.50	0.22

The method of determining duration for this table (except for Subject 18, see below) was by subtracting the time of the first measured value from the time of the last measured value over 20%. This is conservative in defining the actual duration of time above 20% and should be considered at least likely to be an underestimate of the true duration.

As mentioned in the text and shown in Table 15, Subject 18 had only one value in excess of 20%. Applying the method used elsewhere in this table would give a duration of 0 hours of inhibition in excess of 20% of baseline, which is clearly incorrect. In this subject's case only, the degree of inhibition was determined by linear interpolation using the values bounding the single measured inhibition of more than 20%.

Duration of Inhibition of Erythrocyte Acetylcholinesterase
After 16 mg of Pyridostigmine Bromide Orally

Table 18

SUBJ#	EARLIEST TIME OF ACHE INHIBITION ABOVE 20%	LATEST TIME OF ACHE INHIBITION ABOVE 20%	DURATION OF ACHE INHIBITION ABOVE 20% ¹
7	1.67	4.00	2.33
8	2.00	4.00	2.00
9	4.00	4.00	1.172
10	2.00	4.00	2.00
11	0.75	4.00	3.25
12	1.67	4.00	2.33
13	0.75	4.00	3.25
14	1.00	6.00	5.00
15	1.33	6.00 ₃	4.67
16	3.00	0.00	3.00
17	1.00	4.00	3.00
18	1.00	2.00	1.00
19	0.50	3.00	2.50
20	1.67	4.00	2.33
21	1.67	2.00	0.33
22	1.00	4.00	3.00
23	1.00	4.00	3.00
24	1.00	4.00	3.00
MEAN	1.50	4.06	2.62
SD	0.87	1.11	1.15
CV(%)	57.76	27.37	43.80
MAX	4.00	6.00	5.00
MIN	0.50	2.00	0.33

The method of determining duration for this table (except for Subject 9, see below) was by subtracting the time of the first measured value from the time of the last measured value over 20%. This is conservative in defining the actual duration of time above 20% and should be considered at least likely to be an underestimate of the true duration.

² As mentioned in the text and shown in Table 16, Subject 9 had only one value in excess of 20%. Applying the method used elsewhere in this table would give a duration of 0 hours of inhibition in excess of 20% of baseline, which is clearly incorrect. <u>In this subject's case only</u>, the degree of inhibition was determined by linear interpolation using the values bounding the single measured inhibition of more than 20%.

³ Inhibition at 4 hours was 19%.

Table 19

Maximal Inhibition of Erythrocyte Acetylcholinesterase
With 1.3 mg of Pyridostigmine Bromide Intravenously

SUBJECT	#		MAXIMAL INHIBITION	(%)
7			29	
8			28	
9			29	
10			26	
11			29	
12			33	
13			39	
14			32	
15			28	
16			26	
17			29	
18			22	
19			36	
20			32	
21			29	
22			30	
23			29	
24			31	
		Mean	30	
		SD	3.8	
		CV (%)		
		MAX	39	
		MIN	22	

Table 20

Maximal Inhibition of Erythrocyte Acetylcholinesterase
With 16 mg of Pyridostigmine Bromide Orally

SUBJECT		MAXIMAL IBITION (%)	TIME AFTER DOSE TO PEAK INHIBITION (hrs)
7		31	3.0
8		26	3.0
9		22	4.0
10		25	3.0
11		34	1.67
12		30	3.0
13		35	3.0
14		32	2.0
15		39	3.0
16		20	3.0
17		31	1.67
18		22	1.33
19		31	1.0
20		29	3.0
21		22	2.0
22		27	3.0
23		27	3.0
24		31	2.0
	Mean	29	2.5
	SD	5.1	0.8
	CV (%)	17.1	30.9
	MAX	39	4.0
	MIN	20	1.0

Table 21

Area within the Hysteresis Loop

SUB #	IV CURVE	PO CURVE
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21	242.195 197.285 264.740 68.450 96.955 261.205 437.330 332.490 225.785 181.780 204.725 140.495 308.875 324.390 263.840	104.040 39.090 35.600 3.885 198.335 53.730 156.225 146.970 151.300 61.760 157.790 141.060 85.010 91.130
21 22 23 24	263.840 241.605 250.070 185.295	70.655 111.635 115.860 114.140
AVG STD DEV	234.862 87.470	101.956 51.459
p vs zero	7.25×10^{-3}	0.048

Paired t, p = 1.65×10^{-5}

Table 22

Indicators of "Goodness of Fit" for the Curve Fitting to the Effect Model

SUB #		IV WEIGHTED RESIDUALS	CORREL	PO SUM of WEIGHTED SQUARED RESIDUALS	CORREL
7		.0367	.998	4.0510	.938
8		.0197	.988	1.1140	.872
9		.0388	.987	9.0830	.920
10		.0691	.970	34.0136	.860
11		.0582	.979	.0234	.988
12		.1130	.964	.1220	.930
13		.1530	.965	.0982	.936
14		.0440	.972	.0698	.985
15		.0363	.982	5.0580	.962
16		.2340	.868	.2740	.958
17		.1580	.933	.0271	.988
18		.0182	.993	1.1620	.927
19		.9070	.987	.2280	.925
20		.0441	.988	.0853	.909
21		.2590	.817	.0530	.924
22		.0645	.985	.1160	.945
23		.3150	.969	.0631	.981
24		.0490	.967	.1280	.936
AVG		.1454	.9618	3.0983	.9380
STD DE	V	.2097	.046	7 8.0857	.0362

Table 23

Estimates of the Rate Constant of Exit from the Effect Compartment and Pyridostigmine Concentration that Produces 50% Inhibition of Erythrocyte Acetylcholinesterase

SUBJ#	KEO /hr	IC50 ng/ml
7	3.756	27.563
8 9	4.624	30.165
	2.621	22.066
10	4.530	18.345
11	3.399	41.063
12	3.330	31.484
13	5.761	30.934
14	2.759	37.632
15	3.205	26.721
16	1.278	23.126
17	1.890	22.308
18	4.571	39.738
19	3.067	17.307
20	3.237	33.916
21	1.222	33.386
22	5.728	31.013
23	3.186	33.398
24	3.395	30.116
2.		001220
AVG	3.420	29.460
STD DEV	1.284	6.842
OID DEV	1.207	0.042

Table 24

Indicators of "Goodness of Fit" for the Curve Fitting to the Biochemical Model

SUB #	IV SUM of WEIGHTED SQUARED RESIDUALS	CORREL	ORAL SUM of WEIGHTED SQUARED RESIDUALS	CORREL
7	.00163	.995	.00890	.957
8	.00111	.991	.00688	.910
9	.00169	.986	.00905	.888
10	.00442	.960	.01416	.868
11	.00736	.969	.00394	.989
12	.00740	.968	.01050	.911
13	.00399	.967	.01390	.946
14	.00161	.987	.01040	.978
15	.00262	.975	.00603	.964
16	.00471	.907	.00212	.965
17	.00388	.988	.00255	.985
18	.00048	.993	.00547	.935
19	.00697	.979	.00789	.973
20	.00201	.984	.00954	.912
21	.00496	.972	.00245	.977
22	.00158	.981	.00600	.960
23	.00510	.984	.00483	.990
24	.00205	.971	.01120	.939
AVG	.00353	.97539	.00754	.94706
STD DEV	.00222	.01978	.00370	.03622


Table 25

Estimates of the Biochemical Rate Constants for Erythrocyte Acetylcholinesterase Inactivation and Reactivation

SUB #		K(inact)K (po-fit) ml			K(react) Property (po-fit)	
7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	.092 .102 .077 .164 .073 .095 .076 .071 .093 .053 .083 .079 .067 .073 .064 .096	.073 .063 .177 9.651* .067 .130 .118 .045 .089 .077 .093 .059 .589* .075 .051 .115 .060	.083 .083 .127 .164 .070 .113 .097 .058 .091 .065 .088 .069 .067 .074 .058 .105 .077	2.729 3.278 1.860 2.860 3.570 3.202 3.294 2.968 2.909 1.488 2.371 3.596 1.652 2.495 2.489 3.799 3.339 3.160	1.933 1.938 4.455 245.280* 2.726 5.012 3.426 1.634 2.398 1.948 2.272 2.394 13.984* 2.912 1.914 3.335 2.184 2.440	2.331 2.608 3.158 2.860 3.148 4.107 3.360 2.301 2.653 1.718 2.321 2.995 1.652 2.704 2.201 3.567 2.762 2.800
AVG STD DEV	.086	.076	.087 .027	2.837 .669	2.385 1.246	2.736 .616


^{*} Not included in the calculations of the average and standard deviation

Figure la




Schematic of the model used for the analysis of pyridostigmine effects assuming an effect compartment

Figure 1b



Schematic of the model used for the analysis of pyridostigmine effects using a biochemical model






TIME

Average pyridostigmine base concentration in nanograms per milliliter for each sampling time after oral and intravenous administration of drug






IINE

+ 10

Time in hours after dosing Subjects 7 - 12: Concentrations of pyridostigmine base in nanograms per milliliter at each sampling time after intravenous administration of drug







TIME -X-9 -+ 10

Time in hours after dosing
Subjects 7 - 12: Concentrations of pyridostigmine base in nanograms per milliliter at each sampling time after oral administration of drug






Time in hours after dosing
Subjects 13 - 18: Concentrations of pyridostigmine base in nanograms per milliliter at each sampling time after intravenous administration of drug








5 0 0 1 2 3 4 5 6 7 8 9 10 11

Time in hours after dosing

Subjects 13 - 18: Concentrations of pyridostigmine base in nanograms per milliliter at each sampling time after oral administration of drug






IIIE


Time in hours after dosing
Subjects 19 - 24: Concentrations of pyridostigmine base in nanograms per milliliter at each sampling time after intravenous administration of drug






IIE


Time in hours after dosing
Subjects 19 - 24: Concentrations of pyridostigmine base in
nanograms per milliliter at each sampling time after oral
administration of drug



Amount of pyridostigmine base absorbed plotted as a function of the size of the oral dose. X's are individual data points from subjects in whom pyridostigmine bioavailability was determined. The amount absorbed was calculated from the fraction available times the size of the oral dose. The 10.3 mg doses are from this study, the 41.6 mg doses are data from Breyer-Pfaff et al. Clin Pharmacol Ther 37:495,1985 and the 83.3 mg doses are calculated from the data of Aquilonius et al. Eur J Clin Pharmacol 18:423,1980.




Average fractional inhibition of erythrocyte acetylcholinesterase for each sampling time after oral and after intravenous administration of drug







Time in hours after dosing
Subjects 7 - 12: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after intravenous administration of drug



Time in hours after dosing
Subjects 7 - 12: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after oral administration of
drug




Time in hours after dosing Subjects 13 - 18: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after intravenous administration of drug






Time in hours after dosing
Subjects 13 - 18: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after oral administration of
drug




Time in hours after dosing Subjects 19 - 24: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after intravenous administration of drug






Time in hours after dosing
Subjects 19 - 24: Fractional inhibition of erythrocyte acetylcholinesterase at each sampling time after oral administration of
drug





Fractional inhibition of erythrocyte acetylcholinesterase after an intravenous dose of pyridostigmine plotted against the plasma concentration of pyridostigmine base in nanograms per milliliter. At each sampling time after dosing the mean fractional inhibition is plotted against the mean pyridostigmine concentration. The data points are connected in sequence beginning with the baseline sample.

Figure 5b



Fractional inhibition of erythrocyte acetylcholinesterase after an oral dose of pyridostigmine plotted against the plasma concentration of pyridostigmine base in nanograms per milliliter. At each sampling time after dosing the mean fractional inhibition is plotted against the mean pyridostigmine concentration. The data points are connected in sequence beginning with the baseline sample.

APPENDIX A

DETERMINATIONS OF PYRIDOSTIGMINE DOSE ADMINISTERED

The amount of pyridostigmine in each intravenous and oral dose was determined. Syringes containing the intravenous pyridostigmine solution were intentionally overfilled by the pharmacy so that a small aliquot of each solution could be expelled and saved for pyridostigmine assay in the process of preparing the correct volume of solution to administer. Just before the syringe containing the intravenous pyridostigmine was placed in the electrical infusion pump for infusion, it was weighed on a Mettler balance. The infusion was administered, and upon removing the syringe from the pump it was promptly weighed again on the Mettler balance. The specific gravity of the pyridostigmine solution was determined by delivering exactly 5 ml of the solution using a Rainin pipette into a pre-weighed beaker, and dividing by 5 to determine the weight of the solution per milliliter.

A similar procedure was followed for oral dosing. The appropriate volume of syrup was drawn up into the syringe and weighed on a Mettler balance. The dose was administered by expelling the contents of the syringe onto the volunteer's tongue, and the syringe was promptly weighed again. The specific gravity of the syrup was determined by placing exactly 10 ml of syrup in a pre-weighed volumetric flask, weighing the flask with its syrup contents, subtracting to find the weight of the syrup, and dividing that weight by 10 to determine the weight of syrup per milliliter.

The weights of the syringe before and after dosing, the weight of the dose, the specific gravity of the solution, the concentration of pyridostigmine by assay after the solution, and the total dose administered, is provided in the following table:

		Pre-Dose	Post-Dose	Weight			Concent, of	
		Syringe	Syringe	Administered	Specific	Volume	Administered	Dose of Drug
		Weight	Weight	Drug	Gavity	Administered	Drug	Administered
Subject	Route	(grams)	(grams)	(grams)	(g/m1)	(nf)	(ug/m])	(mg base)
7	ΛI	42.74206	22.68077	20.06129	1.00225	20016.25	45	0.90073
80	ΙΛ	2.5	22.48720	20.06036	1.00225	20015.33	43	0.86066
6	ΙN	42.74748	22.65576	20.09172	1.00225	20046.62	94	0.92214
10	IV	42.44876	22.43366	20.01510	1.00225	19970.17	97	0.91863
11	IV	42.35694	22.34901	20.00793	1.00225	19963.01	47	0.93826
12	IV	42.55140	22,48629	20.06511	1.00225	20020.06	67	86086.0
13	ΙN	41.77491	21.83540	19.93951	1.00225	19894.75	97	0.91516
14	IV	41.94086	21.84021	20.10065	1.00225	20055.53	87	0.96267
15	IV	42.46026	22.27323	20.18703	1.00225	20141.71	43	0.86609
16	IV	43.42423	23.16032	20.26391	1.00225	20218.42	47	0.95027
17	IV	7	22.68	20.01	1.00225	19965.08	42	0.83853
18	IV	2.4	22.47	19.96	1.00225	19915.19	39	0.77669
19	ΙV	.7	22.95	19.78	1.00225	19735.59	47	0.92757
20	ΛI	2.3	22.38	19.98	1.00225	19935.15	47	0.93695
21	IV	41.82	21.77	20.05	1.00225	20004.99	42	0.84021
22	ΛI	1.4	21.58	19.83	1.00225	19785.48	97	0,91013
23	11	~	23.13	20.02	1.00225	19975.06	87	0.95880
2.7	110	91 17	23.06	20 12	1 00225	20074 83	41	0 82307
† 7	• 1		7	71.07	7700.1		4	
							Mean	0.90153
							Std. Dev.	0.05393
7	0ra1	7.62439	5.99580	1.62859	1.23235	1321.53	7563.33	9.99515
80	Oral	7.62874	5,98388	1.64486	1.23235	1334.73	7563.33	10.09500
6	Oral	7.63517	5.95228	1.68289	1,23235	1365.59	7563.33	10.32841
10	Oral	7.62954	5.95975	1.66979	1.23235	1354.96	7563.33	10,24801
11	Oral	7.61223		1.67128	1.23235	1356.17	7563.33	10.25715
12	Oral	7.65615	5.97785	1.67830	1.23235	1361.87	7563.33	10.30024
13	Oral	7.65328	5.97348	1.67980	1.23235	1363.08	7563.33	10.30944
14	Oral	7.60178	5.94663	1.65515	1.23235	1343.08	7563.33	10.15816
15	Oral	7.60	5.94	1.66	1.23235	1347.02	7563.33	10.18792
16	Oral	7.54	5.90	1.64	1.23235	1330.79	7563.33	10.06518
17	Oral	7.63947		1.68829	1.23235	1369.97	7563.33	10.36155
18	Oral	7.63564	•	1.69321	1.23235	1373.96	7563.33	10.39174
19	Oral	7.63020	5.95638	1.67382	1.23235	1358.23	7563.33	10.27274
20	Oral	7.63334		1.67690	1.23235	1360.73	7563.33	10.29164
21	Oral	7.65638	5.96912	1.68726	1.23235	1369.14	7563.33	10.35523
22	Oral	7.65638		1.68167	1.23235	1364.60	7563.33	10.32092
23	Oral	7.68977	5.99311	1.69666	1.23235	1376.76	7563.33	10.41292
24	Oral	7.63559	5.96846	1.66713	1.23235	1352.80	7563.33	10.23168

The weights reported above were obtained using a Mettler analytical balance, except where the weights are reported only to the nearest hundredth gram. In those case, a table top Mettler balance was used because the analytical Mettler balance was not functional.

It should be noted that the mean dose of drug administered intravenously, expressed as milligrams of base, was 0.90153 mg, and corresponds to a dose of 1.3 mg pyridostigmine bromide. For the oral dose, the mean pyridostigmine base of 10.25462 mg administered corresponds to a dose of 14.8 mg pyridostigmine bromide.

APPENDIX B

PROCEDURAL TIMETABLE

				DA	JN YA	JMBE	3	
	Screening*	1	2	3	4	5	6	13**
Medical history	X							
Physical examination	X							
CBC with differential	X	X		X			х	
Reticulocyte count		X		X			х	
SMA-6	X	X		X			X	
SMA-12	X	X		X			X	
LDH	Х	X		X			X	
CPK***	Х	X		X			X	
Chest x-ray (within 6 months)	х							
Electrocardiogram	Х			X			X	
Urinalysis	Х	X		X			X	
Red blood cell acetylcholinesterase		X	Х	X		Х	X	
Pyridostigmine levels			х	X		Х	х	
Drug administration			Х			Х		
Discharge from hospital							Х	

- * Screening is conducted within two weeks of the subject's entry into the hospital.
- ** If the subject's clinical labs on Day 6 are abnormal, he will be invited to return for follow-up clinical testing on Day 13 and weekly thereafter until tests become normal or an alternative explanation is determined.
- *** CPK (creatine phosphokinase) is equivalent to creatine kinase (CK), the nomenclature used elsewhere in this report except here in Appendix B.

STUDY FLOW CHART - DOSE-RANGING PHASE

DAY	HOUR	APPROXIMATE TIME	PROCEDURE
1	-21	11:00 AM	Subjects admitted to Unit Sign Consent Form Blood: CBC with differential Reticulocyte count SMA-6 SMA-12 LDH CPK Red blood cell acetylcholinesterase (RBC AChE) EKG Urinalysis
2	-8	12:00 MN	NPO No smoking
	-1	7:00 AM	Insert heparin lock for blood sampling
	0	8:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip, coordination testing (clinical examination) ADMINISTER TEST DRUG
			Pyridostigmine 0.66 mg IV over 30 minutes by infusion pump
	0.08	8:05 AM	Blood: Pyridostigmine level
	0.16	8:10 AM	Blood: Pyridostigmine level
	0.25	8:15 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	0.33	8:20 AM	Blood: Pyridostigmine level

DAY	HOUR	APPROXIMATE TIME	PROCEDURE
2	0.42	8:25 AM	Blood: Pyridostigmine level
	0.50	8:30 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip, coordination testing (clinical examination)
	0.58	8:35 AM	Blood: Pyridostigmine level
	0.66	8:40 AM	Blood: Pyridostigmine level
	0.75	8:45 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	0.83	8:50 AM	Blood: Pyridostigmine level
	0.92	8:55 AM	Blood: Pyridostigmine level
	1.0	9:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	1.33	9:20 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	1.66	level RBC A	Blood: Pyridostigmine level RBC AChE level Pulse rate
	2.00	10:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	2.5	10:30 AM	Blood: Pyridostigmine level

DAY	HOUR	APPROXIMATE	TIME	PROCEDURE
2	3.0	11:00	AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	3.5	11:30	AM	Blood: Pyridostigmine level
	4.0	12:00	PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	5.0	1:00	PM	Blood: Pyridostigmine level
	6.0	2:00	PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	7.0	3:00	PM	Blood: Pyridostigmine level
	8.0	4:00	PM	Blood: Pyridostigmine level and CPK
	10.0	6:00	PM	Blood: Pyridostigmine level
	12.0	8:00	PM	Blood: Pyridostigmine level
3	0	8:00	AM	Blood: Pyridostigmine level RBC AChE level CBC with differential Reticulocyte Count SMA-6 SMA-12 LDH CPK Urinalysis EKG Pulse rate, hand grip Heparin lock may be
				discontinued if the patient desires

DAY	HOUR	APPROXIMATE TIM	PROCEDURE PROCEDURE
4			No procedures scheduled
5	-8	12:00 MN	NPO No smoking
	-1	7:00 AM	Insert heparin lock for blood sampling
	0	8:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
			ADMINISTER TEST DRUG
			Pyridostigmine 20 mg syrup orally
	0.25	8:15 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	0.50	8:30 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	0.75	8:45 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	1.0	9:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	1.33	9:20 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	1.66	9:40 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	2.0	10:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	2.5	10:30 AM	Blood: Pyridostigmine level

DAY	HOUR	APPROXIMATE TIME	PROCEDURE
5	3.0	11:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	3.5	11:30 AM	Blood: Pyridostigmine level
	4.0	12:00 PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	5.0	1:00 PM	Blood: Pyridostigmine level
	6.0	2:00 PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	7.0	3:00 PM	Blood: Pyridostigmine level
	8.0	4:00 PM	Blood: Pyridostigmine level and CPK
	10.0	6:00 PM	Blood: Pyridostigmine level
	12.0	8:00 PM	Blood: Pyridostigmine level
6	0	. 8:00 AM .	Blood: CBC with differential Reticulocyte count SMA-6 SMA-12 LDH CPK RBC AChE level Pyridostigmine level Pulse rate, hand grip, coordination testing (clinical examination) EKG Urinalysis Discharge after breakfast

NOTE: There will be no timed urine collections for this protocol.

STUDY FLOW CHART - BIOAVAILABILITY PHASE

DAY	HOUR	APPROXIMATE TIME	PROCEDURE
1	-21	11:00 AM	Subjects admitted to Unit Sign Consent Form Blood: CBC with differential Reticulocyte count SMA-6 SMA-12 LDH CPK Red blood cell acetylcholinesterase (RBC AChE) EKG Urinalysis
2	-8	12:00 MN	NPO No smoking
	-1	7:00 AM	Insert heparin lock for blood sampling
	0	8:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip, coordination testing (clinical examination)
			ADMINISTER TEST DRUG
			Pyridostigmine intravenous solution
		190 - 1	Dose will be based upon the data obtained in the Dose Ranging Phase. The doses administered will be such that RBC AChE inhibition will be between 20% and 40%.
	0.08	8:05 AM	Blood: Pyridostigmine level
	0.16	8.10 AM	Blood: Pyridostigmine level
	0.25	8:15 AM	Blood: Pyridostigmine level
	0.33	8:20 AM	Blood: Pyridostigmine level
	0.42	8:25 AM	Blood: Pyridostigmine level

DAY	HOUR	APPROXIMATE	TIME	PROCEDURE	<u> </u>	
2	0.50	8:30 /	AM		Pyridostigmine ip, coordination (clinical exami	l
	0.58	8:35	AM	Blood:	Pyridostigmine	level
	0.66	8:40	AM ·	Blood:	Pyridostigmine	level
	0.75	8:45	AM	Blood:	Pyridostigmine	leve1
	0.83	8:50	AM	Blood:	Pyridostigmine	level
	0.92	8:55	AM	Blood:	Pyridostigmine	level
	1.0	9:00	AM		Pyridostigmine E level ate, hand grip	level
	1.33	9:20	AM	Blood:	Pyridostigmine	level
	1.66	9:40	AM	Blood:	Pyridostigmine	level
	2.0	10:00	AM	Blood: RBC AChi Pulse ra	Pyridostigmine E level ate, hand grip	level
	2.5	10:30	AM	Blood:	Pyridostigmine	level
	3.0	11:00	AM	Blood: RBC AChi Pulse ra		level
	3.5	11:30	AM	Blood:	Pyridostigmine	level
	4.0	12:00	PM	RBC ACh	Pyridostigmine E level ate, hand grip	level
	5.0	1:00	PM	Blood:	Pyridostigmine	level
	6.0	2:00	PM .	Blood: RBC AChi Pulse ra	Pyridostigmine E level ate, hand grip	level
	7.0	3:00	PM	Blood:	Pyridostigmine	level
	8.0	4:00	PM	Blood: and CPK	Pyridostigmine	level

DAY	HOUR	APPROXIMATE TIME	PROCEDURE
2	10.0	6:00 PM	Blood: Pyridostigmine level
	12.0	8:00 PM	Blood: Pyridostigmine level
3	0	8:00 AM	Blood: Pyridostigmine level RBC AChE level, CBC with differential Reticulocyte count SMA-6 SMA-12 LDH CPK Pulse rate, hand grip EKG Urinalysis Heparin lock may be discontinued if the patient
			desires.
4			NO procedures scheduled
5	-8	12:00 MN	NPO No smoking
	-1	7:00 AM	Insert heparin lock for blood sampling
5	0	8:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
			ADMINISTER TEST DRUG
			Pyridostigmine syrup
	0.25	8:15 AM	Blood: Pyridostigmine level
	0.50	8:30 AM	Blood: Pyridostigmine level Hand grip
	0.75	8:45 AM	Blood: Pyridostigmine level
	1.0	9:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	1.33	9:20 AM	Blood: Pyridostigmine level

DAY	HOUR	APPROXIMATE TIM	ME PROCEDURE
	1.66	9:40 AM	Blood: Pyridostigmine level
	2.0	10:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	2.5	10:30 AM	Blood: Pyridostigmine level
	3.0	11:00 AM	Blood: Pyridostigmine level RBC AChE level Pulse rate
	3.5	11:30 AM	Blood: Pyridostigmine level
	4.0	12:00 PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	5.0	1:00 PM	Blood: Pyridostigmine level
	6.0	2:00 PM	Blood: Pyridostigmine level RBC AChE level Pulse rate, hand grip
	7.0	3:00 PM	Blood: Pyridostigmine level
	8.0	4:00 PM	Blood: Pyridostigmine level and CPK
	10.0	6.00 PM	Blood: Pyridostigmine level
	12.0	8:00 PM	Blood: Pyridostigmine level
6		8:00 AM	Blood: CBC with differential Reticulocyte count SMA-6 SMA-12 LDH CPK RBC AChE level Pyridostigmine level Pulse rate, hand grip, coordination testing (clinical examination) EKG Urinalysis Discharge after breakfast

NOTE: There will be no timed urine collections for this protocol.

APPENDIX C
CLINICAL LABORATORY NORMAL VALUES

The ranges of these values have been determined and are utilized by the Department of Laboratory Medicine of The Johns Hopkins Hospital.

Serum Chemistry	Normal Limits		<u>Units</u>
<u>Tests</u>	Lower	Upper	
Sodium	135	148	mEq/1
Potassium	3.5	5.0	mEq/1
Chloride	96	109	mEq/1
Carbon dioxide	24	30	mEq/1
Serum urea nitrogen	12	25	mg/dl
Creatinine	0.4	1.5	mg/dl
Glucose	70	115	mg/dl
Calcium	9.0	11.0	mg/dl
Total bilirubin	0.3	1.2	mg/dl
Direct bilirubin	0.1	0.4	mg/dl
Total protein	6.0	8.5	g/dl
Albumin	3.2	5.3	g/dl
Aspartate aminotransferase	0	35	IU/1
Alanine aminotransferase	0	30	IU/l
Alkaline phosphatase	0	95	IU/l
Phosphate, inorganic	3.0	4.5	mg/dl
Lactic dehydrogenase	0	200	IU/l
Creatine kinase	0	160	IU/l
Uric Acid	4.2	8.8	mg/dl
Cholesterol	146	270	mg/dl

	Normal Limits		<u>Units</u>
	Lower	Upper	
Hematology Tests			
White blood cells	4,500	11,000	#/mm³
Red blood cells	4.50	5.90	million/mm ³
Hemoglobin	13.9	16.3	g/dl
Hematocrit	41.0	53.0	*
Platelets	150	350	thousand/mm ³
Reticulocytes	0.5	1.5	8
White Blood Cell Differentials:			
Bands	2	6	%
Segmental Neutrophils	31	76	8
Lymphocytes	24	44	8
Monocytes	2	11	ર
Eosinophils	1	4	8
Basophils	0	1	%

APPENDIX D

ASSAY OF ERYTHROCYTE ACETYLCHOLINESTERASE AT THE JOHNS HOPKINS UNIVERSITY DIVISION OF CLINICAL PHARMACOLOGY

The determination of erythrocyte acetylcholinesterase (AChE) follows the protocol established by the US Army Medical Research Institute for Chemical Defense for the determination of erythrocyte AChE.

Principle of Method, Chemicals, Equipment and Materials, Preparation of Reagents

The principles of the methods, the chemicals, materials, and equipment needed, the preparation of reagents and standards are detailed in the Standard Operating Procedure (SOP) of 18 June 1985 of the Analytical Chemistry Branch, USAMR, ICD, Aberdeen Proving Ground, Maryland 21010. These are followed exactly except for preparation of the stock glutathione solution, page 3, number 6 under A. Reagents. The typographical error is corrected and 1.844 g of glutathione, reduced form, (GSH) are dissolved in a total volume of 100ml of EDTA diluent.

Collection and Preparation of Specimen for Sampling

Blood is collected into 3 ml purple topped Vacutainer^R

(EDTA) through a catheter inserted into an arm vein. After the sample is drawn, it is mixed and brought promptly to the lab at room temperature. One and one half ml of the blood is centrifuged for 2 minutes at 15,000 RPM in an Eppendorf model

5414 centrifuge. After centrifugation all plasma is removed from

the top of the packed erythrocytes with a Pasteur pipet. The erythrocytes are then picked up in a clean Pasteur pipet, starting from the bottom of the centrifuge tube, avoiding drawing air after all the cells have been pipetted. The packed cells are transferred to a 0.5 ml sample cup and are ready for AChE analyses. Sample preparation time is kept to 3 minutes.

Preparation of Standards

Standards are prepared according to the SOP of the ICD.

Preparation of AChE Control Material

A quality control enzyme standard is used to measure assay precision. The enzyme used is electric eel AChE which is diluted in a large volume to a specific activity and frozen in aliquots. An aliquot is used each day the assay is performed. Electric eel samples were assayed in accordance with ICD SOP. See below for specific details.

Analysis Start-Up

ICD SOP is followed. The heating circulator is turned on. We allow temperature to reach 37 degrees C and verify constancy periodically.

Proportioning Pump

ICD SOP is followed. Water with Brij 35 precedes the reagents. A good bubble pattern is verified.

Colorimeter and Recorder

ICD SOP is followed. Lamp warmup is at least 10 minutes.

Once reagents are running, the recorder is zeroed for no signal and full scale. Speed is set to 1.0 cm/min.

Sampler

ICD SOP is followed. Proper operation of sampler is verified prior to running any samples.

Analyses

Recorder baseline is set to zero.

Manifold has been modified so that the substrate blank side pumps saline continuously. Erythrocytes enter only one side and one photo cell. At the end of the day, a substrate blank is run by placing the substrate line into saline and measuring the activity of the erythrocytes without a substrate. ICD SOP is followed.

Incubation time is measured. The length of time it takes erythrocytes to travel from the point where substrate is added to the sample stream until exit from the dialyzer is defined as the incubation time.

GSH standards and electric eel AChE are assayed first. A 60 micromoles/ml (u moles/ml) GSH standard is assayed followed by 15, 30, and 45 u moles/ml GSH standards, and a 1:1 dilution of the electric eel quality control. Machine sensitivity is adjusted using the STD CAL knob so that the 60 u moles/ml standard reads 80 to 90 on the recorder scale. Two standard curves and a minimum 3 electric eel samples are assayed prior to assay of erythrocyte samples. Erythrocyte samples are separated by at least two saline cups. Machine performance is standardized periodically throughout the day by running standard curves and electric eel samples. Gain is decreased if the 60 u moles/ml GSH rises above 90. Where possible, standards are assayed hourly and

erythrocyte samples measured in duplicate.

Shut Down SOP

ICD is followed. Care is taken to dry the tubes and to insure that they are in a relaxed position.

Data Reduction

AChE activity is determined as follows:

The concentration of each GSH standard is divided by the peak height of the deflection produced. This value is then divided by the time of incubation in minutes to produce a factor of u moles/ml/min/chart unit. The individual factors from each standard in the curve are averaged. The average factor, u moles/ml/min/chart unit, is used to convert the deflection produced by samples to AChE activity, u moles/ml/min.

ASSAY PERFORMANCE AT JOHNS HOPKINS HOSPITAL Quality Control

A standard solution of electric eel acetylcholinesterase is measured several times on the days that assays are performed.

The current batch of JHH electric eel was prepared on 14 October 1985 using Sigma Co, Type VI-S Cholinesterase, Acetyl (catalog #C3389) from electric eel, Lot #83F8100. One 10,000 unit vial was diluted in Tris buffer containing 1% bovine serum albumin. The dilution target was such that a one to one dilution of the solution should contain 15.0 u moles/ml/min of cholinesterase activity.

After the dilution, aliquotting and freezing of the cholinesterase solution was performed, it was discovered that the

colorimeter was not functioning correctly and that the GSH standards were in error. New standards were made; the colorimeter was repaired. The instrument was calibrated with GSH standards and the Aberdeen quality control standard. The latter's activity was 7.46 u moles/ml/min, within the range at ICD, Aberdeen.

The assay was used on 13 different days. On each day an aliquot of the electric eel quality control standard was thawed and assayed several times. The overall statistics of 127 samples from the 13 aliquots show mean activity 9.88 u moles/ml/min with a standard deviation of 0.27 u moles/ml/min. The coefficient of variation is 2.7%. Looking only at the means of the assays in a single day, the activity of the quality control is 9.91 \pm 0.23 u moles/ml/min (C.V. = 2.3%).

Precision

Blood samples were drawn from 6 study subjects prior to drug administration. Each sample was assayed on several occasions.

The results are tabulated in the Table.

The intra-assay coefficient of variation ranged from 0.5 to 3.5%, with a mean of 2.5%. Day to day variations in the measurement of a subject's drug free erythrocyte AChE ranged from 0.14 to 0.67 u moles/ml/min or 1.08 to 5.48% of the value on a single day. This variation is about twice the intra-assay variation and suggests that either the inter-assay variation is greater than that of the intra-assay and/or the activity of erythrocyte AChE drawn from a single subject on different days varies to a small degree.

With a coefficient of variation of 2.5% in the assay, inhibition of enzyme activity of 5% or more probably represents real inhibition by pyridostigmine, not random variation due to the assay.

TABLE - APPENDIX D

ERYTHROCYTE ACETYLCHOLINESTERASE ACTIVITY IN 6 NORMAL VOLUNTEERS

Intra-Assay and Inter-Assay Results

Subject	Day	n	AChE u moles/ml/min (± S.D.)	C.V.	b-a u moles/ml/min	(b-a)/b %
A	a b	7 9	$12.90 \pm 0.31 \\ 13.04 \pm 0.31$	2.4	0.14	1.08
В	a b	10 8	$13.82 \pm 0.37 \\ 14.48 \pm 0.55$	2.7	0.66	4.78
С	a b	8 8	$12.59 \pm 0.36 \\ 13.28 \pm 0.31$	2.9	0.69	5.48
D	a b	4 5	$13.20 \pm 0.52 \\ 13.87 \pm 0.07$	3.9 0.5	0.67	5.08
Е	a b	4 6	$\begin{array}{c} 13.35 \pm 0.43 \\ 13.86 \pm 0.33 \end{array}$	3.2 2.4	0.51	3.82
F	a b	4 6	$\begin{array}{c} 14.26 \pm 0.36 \\ 14.86 \pm 0.20 \end{array}$	2.5 1.3	0.60	4.21

Form C (Revised 4/84) J.H.U.M.S.

CLINICAL INVESTIGATION CONSENT FORM The Johns Hopkins Medical Institutions

Title of Research Project:

BIOAVAILABILITY OF ORAL PYRIDOSTIGMINE AND INHIBITION OF ACETYLCHOLINESTERASE BY ORAL AND INTRAVENOUS PYRIDOSTIGMINE

Patient I.	. .	

Explanation of Research Project to Subject:

You are invited to participate in a study of an approved and marketed drug, pyridostigmine. This drug is currently in use for treating patients with a disease called myasthenia gravis. The drug, based on studies in animals, may also be an effective pre-treatment for accidental poinoning with certain insecticides which work in ways similar to nerve gases. This study is designed to see how much of the drug gets into the blood stream when given by mouth and how effective the levels are in changing a blood test that may relate to the degree of protection from poisoning.

If you agree to join this study, you will be hospitalized for five days. During the course of the hospitalization you will have a number of blood samples drawn by means of a "heparin lock" which allows us to take repeated blood samples without sticking a new needle into your vein each time. The total blood taken will be about one pint, which is the amount taken if you donated blood at a blood bank. You should not donate blood for six to eight weeks after the conclusion of the study.

On the second day, after the first blood sample is taken, you will be given the drug pyridostigmine, either in syrup form or into your vein. On the fifth day, you will be given pyridostigmine in the manner not previously given (i.e., syrup or into the vein). Wehther you get the syrup or dose by vein first will be determined "randomly" in a process like flipping a coin.

We believe that the risks of participation in this study are small. Pyridostigmine is not an experimental drug. The dose of the drug you will receive is much smaller than the dose usually used in treating patients. In a study similar to this one where doses about three times larger than the ones you will receive were used, the only side effects noted were temporary fatigue, muscle twitching, and gastric distress. Patients who take more than 20 times as much of the drug each day than you will receive in this study sometimes develop nausea, vomiting, diarrhea, abdominal cramps, and increased body secretions. Because the dose you will receive is so much small than what doctors use in patients, it is not likely that you will develop these symptoms. Treatment is available if any such symptoms occur and become severe.

You are under no obligation to participate in this project. Should you decide not to participate or should you decide to withdraw during the course of the project, your future medical care at Hopkins will not be affected. Benefits to you for participation in this study are primarily financial, but another potential asset is the comprehensive evaluation which accompanies this project, the records of which will be available in the future. You will be paid by check for whatever proportion of the study you complete. Successful completion of the entire study will pay \$340. You will be paid by check at the time you leave the hospital.

Expenses for medical treatment to treat any injury that is directly a result of your participation in accordance with this protocol will be provided by the U.S. Army. The medical treatment provided might include, if necessary, laboratory tests, x-rays, and other procedures used in diagnosis and treatment. No other compensation for injury is offered.

If you sign this form, you are willing to join the research project described on the other side of this page. Your doctors did explain the other kinds of treatment that are available to you and to others. You should ask any questions you have about this research study. You may ask questions in the future if you do not understand something that is being done.


The records from this research study will not be given to anyone who is not helping on this study unless you agree to have the records given out. If the study uses a drug that is under the jurisdiction of the Food and Drug Administration (FDA), the FDA government officials may look at the relevant part of your

medical records as part of their job to review new drug studies.

If you want to talk to anyone about this research study because you think you have not been treated fairly or you think you have been hurt by joining the study, you should call <u>Dr. Brent G. Petty</u> (Principal Investigator) at <u>955 - 8181</u>, or call the Office of the Joint Committee on Clinical Investigation at 955-3008. Either the investigator or the people in the Committee office will help to find medical care for the injury you feel you have suffered. You should understand that The Johns Hopkins University, The Johns Hopkins Hospital, and the Federal Government do not have any program to provide compensation for you if you experience injury or other bad effects which are not the fault of the investigators.

You may withdraw from the research study at any time. Even if you do not want to join the study, or if you withdraw from it, you will still have the same quality of medical care available to you at Johns Hopkins.

If you agree to join this study, please sign your name below.



Subject's signature (Including children, when ap	plicable	
Signature of Parent or Guard	dian (when applicable)	
Witness to Consent Procedu	res*	
Signature of Investigator		

NOTE: Signed copies of this consent form must be a) retained on file by the Principal Investigator; b) deposited in the patient's medical record; and c) given to the patient.

CLINICAL INVESTIGATION CONSENT FORM The Johns Hopkins Medical Institutions

Title of Research Project:

Bioavailablity of Oral

Pyridostigmine and Inhibition of Acetylcholinesterase by Oral and Intravenous Pyridostigmine

Patient I.D. Plate	

Explanation of Research Project to Subject:

You are invited to participate in a study of an approved and marketed drug, pyridostigmine. Pyridostigmine is a medicine that has been prescribed for patients, using daily doses for months or years, for a period of over 20 years. This drug is used for treating patients with a disease called myasthenia gravis. The drug, based on studies in animals, may also be effective pre-treatment for accidental poisoning with certain insecticides which work in ways similar to nerve gases. This use of the drug is considered investigational by the Food and Drug Administration. This study is designed to see how high the drug level is in the blood stream when given by mouth or into your vein and how effective the levels are in changing a blood test that may relate to the degree of protection from poisoning.

If you agree to join this study, you will be hospitalized for five days. During the course of the hospitalization you will have a number of blood samples drawn by means of a "heparin lock" which allows us to take repeated blood samples without sticking a new needle into your vein each time. The total blood taken will be about one pint which is the amount taken if you donated blood at a blood bank. You should not donate blood for six to eight weeks after the conclusion of the study.

On the second day, after the first blood sample, you will be given the drug pyridostigmine, either in syrup form or into your vein. On the fifth day, you will be given pyridostigmine in the manner not previously given (i.e., syrup or into the vein.) Whether you get the syrup or intravenous dose first will be determined "randomly" in a process like flipping a ccin.

We believe that the risks of participation in this study are small. Pyridostigmine is considered an experimental drug in this project because of the new use for which it is intended. The dose of the drug you will receive is much smaller than the dose usually used in treating patients. In a study similar to this one where doses about three times larger than the ones you will receive were used, the only side effects noted were temporary fatigue, muscle twitching, and gastric distress. Patients who take more than 20 times as much of the drug each day than you will in this study sometimes develop nausea, vomiting, diarrhea, abdominal cramps, and increased body secretions. Pecause the dose you will receive is so much smaller than what doctors use in patients, it is not likely that you will develop these symptoms. Treatment is available if any such symptoms occur and become severe.

Recently, a possible new toxicity in rats was discovered which may be related to the drug. The rats were given pyridostigmine in a higher dosage than the one planned for this study. A change in the muscle tissues in the area where the nerves and muscles meet was seen using the electron microscope. The significance of this finding is unknown, but it is generally accepted in the medical field that pyridostigmine is safe for man at the doses to be used in this study. Further testing in other animals is being done to see if this nerve-muscle effect occurs in animals other than the rat. Additional tests in rats are in progress to find out the dose at which the effect occurs and whether the tissue returns to normal when the pyridostigmine is stopped. If results of these tests become available during this study, you will be informed of them. Until these tests are completed, a small risk of developing these changes cannot be ruled out for participants in this study. However, all

THIS CONSENT FORM CONTINUES ON THE REVERSE SIDE


If you sign this form, you are willing to join the research project described of the other side of this page. Your doctors did explain the other kinds of treatment that are available to you and to others. You should ask any questions you have about this research study. You may ask questions in the future if you do not understand something that is being done.

The records from this research study will not be given to anyone who is not helping on this study unless you agree to have the records given out. Since the study uses a drug that is under the jurisdiction of the Food and Drug Administration (FDA), the FDA government officials may look at the relevant part of your medical records as part of their job to review new drug studies. Officials from the U.S. Army involved with drug development may also review the records for routine monitoring of the study.

If you want to talk to anyone about this research study because you think you have not been treated fairly or you think you have been hurt by joining the study, you should call __Dr. Brent G. Petty __ (Principal Investigator) at __x8181 _, or call the Office of the Joint Committee on Clinical Investigation at 955-3008. Either the investigator or the people in the Committee office will help to find medical care for the injury you feel you have suffered. You should understand that The Johns Hopkins Unversity, The Johns Hopkins Hospital, and the Federal Government do not have any program to provide compensation for you if you experience injury or other bad effects which are not the fault of the investigator.

You may withdraw from the research study at any time. Even if you do not want to join the study, or if you withdraw from it, you will still have the same quality of medical care available to you at Johns Hopkins.

If you agree to join this study, please sign your name below.



Subject's s	ignature		
Witness to	Consent Pro	cedures	
Signature o	f Investiga	tor	
Date			

NOTE: Signed copies of this consent form must be a) retained on file by the Principal Investigator; b) deposited in the patient's medical record; and c) given to the patient.

experience with the clinical use of pyridostigmine in man over the past 20 years has not led to any reports of nerve-muscle difficulties attributed to or resulting from the use of this drug.

You are under no obligation to participate in this project. Should you decide not to participate or should you decide to withdraw during the course of the project, your future medical care at Hopkins will not be affected. Benefits to you for participation in this study are primarily financial, but another potential asset is the comprehensive evaluation which accompanies this project, the records of which will be available in the future. You will be paid by check for whatever proportion of the study you complete. Successful completion of the entire study will pay \$375. You will be paid by check at the time you leave the hospital.

Under Army regulation, you are authorized all necessary care which is the direct result of your participation in the research and in accordance with this protocol. The medical treatment provided might include, if necessary, laboratory tests, x-rays and other procedures used in diagnosis and treatment. No other compensation for injury is offered.

APPENDIX G

Publication Supported by this Contract

Kornhauser D, Petty B, Lin E, Schuster B, Pamplin C, Lietman P: In vivo relationship of the plasma concentration of pyridostigmine to the inhibition of red blood cell acetylcholinesterase. III World Conference on Clinical Pharmacology and Therapeutics, Stockholm, Sweden, July 27 - August 1, 1986. (Abstract)

APPENDIX H

Personnel Receiving Contract Support

Paul S. Lietman, M.D., Ph.D., Principal Investigator
Brent G. Petty, M.D., Assistant Investigator
David M. Kornhauser, M.D., Investigator
Laura E. Rocco, R.N., M.S., Research Coordinator
Joan Baumgardner, R.N., Research Nurse
Sherry Sigelman, R.N., Research Nurse
Amina Woods, M.S., Laboratory Administrator
Evelyn Fleckenstein, Secretary
Michelle Janouris, Secretary

All of the personnel above are members of the:

Division of Clinical Pharmacology

Department of Medicine

The Johns Hopkins University School of Medicine

Baltimore, Maryland 21205

APPENDIX I Distribution List

5 copies: Director, Walter Reed Army Institute of

Research

ATTN: SGRD-UWI-F/COL Brian Schuster Washington, D.C. 20307-5100

1 copy: Commandant

Academy of Health Sciences

ATTN: AHS-CMD

Fort Sam Houston, TX 78234

Commander 1 copy:

U.S. Army Medical Materiel Development

Activity Fort Detrick

Frederick, MD 21701-5009

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	P C F M L	01	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	10 Oct \$5	Screening laboratory
_	1700785	History, Physical Exam
0	21007 85	Admission
2	22 OCT 85	I.V.
5	2500785	P.O.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of ______ pages, for subject #_____.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/y)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RCF	01	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY Date of evaluation 10/17/85 In dd mmm yy
15 Nov
In dd mmm yy Date of birth 35yrs Age \overline{w} Sex B

Race

	No	Yes	Comments
Allergy	/		
Tobacco Use		1	1/2 ped
Alcohol Use		/	6 beers /wk
Recreational Drug Use		/	Marijuano + horain 1-1/2 yr
Medications past 2 weeks	1		
Experimental Drug Exposure		~	tobramquin study
Blood or plasma donor		/	lest ~ + year ap
Prior Surgery		/	Surgery Dzygomatic
Eye, ear, nose, throat	1		
Endocrine(diabetes,thyroid)	/		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	1		
Genito-urinary	/		
Musculoskeletal			
Neuropsychiatric			
Pesticide/herbicide use			Not asked
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	R CF F M L	01	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 17/001/85 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height	(cm)	Wt. (kg)
3 6.7 c	_ <i>§ A</i> min	_1 3 /min	7 0	8/72	736	<u>5. 0</u>	<u>65.7</u>

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	/		
EENT	/		
Chest, lungs		/	inspiratory rhocki (1) base that clear
Heart	/		•
Abdomen	/		
Genitalia			N.D.
Rectal			N.D.
Extremities		~	SCAT dorsum @ foot from childhood
Skin		V	scretches both valor forcerms from work
Neurologic	/		

CHEST X-RAY
Date 10/17/85

NORMAL	ABNORMAL	Describe	abnormalities:	
/	•			
ł				

Examiner

Brent G. Petty

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	P C F	01	PROTOCOL DAMD 17-85-C-5133-02

MEDICATION RECORD

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	55 09 82	0900	0930	IV	2 A.

· lole mg. DOSAGE (total)

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
BO1, B47	0	22 02 85	0845	0845	B01: *	B47:12.90
B02	0.08		0905	0905	B05: *	
воз	0.16	٠,	0910	0910	B03: 1,86	
B04, B48	o . 25	••	0915	0915	B04:5,/7	B48: 11:53
B05	0.33	.,	0920	0923	B05:4.77	
B06	0.42		0925	<u> </u>	BO6: 6.48	
B07, B49	0.50		0930	0930	B07: 6.88	B49: 10.73
BOB	0.58	*4	0935	0935	BOB: 3.08	
B 09	0.66	• •	0940	0940	B09: 3.05	
B10, B5 0	0.75	11	0945	0945	B10: 1.86	B50:/0.79
B11	0.83	1.	0950	0950	B11: 🐇	
B12	0.92	11	0955	0955	B12: *	
B13, B51	1.0	٠,	1000	1000	B13: *	B51://.00
B14, B 5 2	1.33	×4	1020	1025	B14: *	B52: //.89
B15, B53	1.66	11	1040	1040	B15: ¥	B53: //. 77
B16, B54	2.0	11	1100	1101	B16: *	B54: 12,06
B17	2.5	13	1130	1130	B17: 💥	

* below away servitivity 137

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	2 C F F M L	Ol	PROTOCOL DAMD 17-85-C-5133-02

MEDICATION RECORD

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
Ω	ಶಾಂಕ್ಷಣ	0 9 0	0930	IV	NA

.66 mg. DOSAGE (total)

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	22 ot 85	1500	1200	B18: *	B55:/2.44
B19	3.5	2202 88		1230	B19: 🔆	
B20, B56	4.0	٠,	1300	1300	B20: #	B56:/2.98
B21	5.0	6	1400	1400	B21: *	
B22, B57	6.0	, V)	1500	1500	B22: *	B57: 12.36
B23	7.0	4.	1600	1600	B23: ★	
B24	8.0	٠,	1700	1700	B24: 1	
B25	10.0	4 •	1900	1900	B25: *	
B26	12.0	11	2100	2100	B26: *	
B27, B58	24.0	230088	0900	১৭০১	B27:*	B58:/3.4/
* le	tow assa	y				
Ser	cow assa	I	138			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	Ref F M L	01	PROTOCOL _DAMD_17085_C-5133-02

MEDICATION RECORD

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	220188	0900	0935	IV	NA

DOSAGE (total) .66 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	220285	0845	0845	#60	#41/45	# C K
	0.8	٠,	0905	0905			
	0.16		0910	0910			
	0.25	٠,	0915	0915	#60		
	0.33	١.	0920	0920			
	0.42		0925	0925			
	0.50	1.	0930	0930	# 55	# 46/49	# ck
	0.58	"	0935	0935		,	
	0.66	• •	0940	0940			
	0.75	٠.	0945	0945	# 58		
	0.83	٤.	0950	೦950			
	0.92		0955	0955			
	1.0		1000	1000	* 64	#46/51	
	1.33	٠,	1020	1050	#62		
	1.66	(.	1040	1040	* 56		
	2.0	٠,	1100	1100	#72	#44/52	
	2.5	1.	1130	1130			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	P C F	0	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	220485	0900	0930	IV	Nti

DOSAGE (total) , 66 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	2202485	1200	1200	# 64		
	3.5		1230	1230			
	4.0		1300	1330	# 84	#19/SL	
	5.0		1400	1400			
	6.0		1500	1500	#80	#57/55	
	7.0		1600	1600			
	8.0		1700	1700			
	10.0		1900	1900.			
	12.0		2100	2100			
	24.0		0900	0920	* 76	# 48/54	# ck
					*/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RCF FML	01	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 02185	0800	2 8	PO	NA

DOSAGE (total) _____ 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	POMC NG/MC	RBC ACHe uM/ml/min	
B28, B59	٥	25 0 tt 85	not recorded	not recorded	B28: *	B59: 13.04	1
B29, B60	0.25	2500+85			B29: ≭	B60: 12.84	
B30, B61	0.50	1.4	0830		B30:4.97	B61: \\. 67	
B31,B62	0.75	١.	0845		B31:5,59	B62: 10.77	
B32, B63	1.0	1.	0900		B32: 9, 8	B63: (□.€)	
B33, B64	1.33	1.	0920		B33:6,79	B64: 9.54	
B34, B65	1.66	9.1	0940		B34: 8,/4	B65: q. eq	
B35, B66	2.0		1000		B35:8,09	B66: 9.25	
B36	2.5	4 (1030		B36: 7,65		
B37, B67	3.0	ι,	1100		B37: 6.85	B67: 9.97	
B38	3.5		1130		B38: 5.66		
B39, B68	4.0	٠,	1200		B39:5,//	B68: 10.44	
B40	5.0		1300		B40:5,99		
B41, B69	6.0	٠,	1400		B41: 2.37	B69:11.39	
B42	7.0	ţ ·	1500		B42: 💥		
B43	8.0	٠,	1600		B43: 💥	B69A: N V.	
B44	10.0	1 *	1800		B44:*		

141

* below assay sensetivity

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	RCE FML	01	PROTOCOL _ DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	250t85	0800	NA	PO	26

DOSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO)	RBC ACHe uM/ml/min
B45	12.0	250265	3000	2000	B45:	
B46, B70	24.0	३६८५५६		0800	B46:*	B70:13 145
			•			
-						
: - - · · · ·						
			,			
	ud amaili					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	RCF FML	٥١	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	25at85	0800	NA	PO	24

DOSAGE (total) 20mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	25 0885	recorded	not receided	# 60	#45/45	# ch.
	0.25	٠,	0815	0815	#64		
	0.50	*1	0830	0830	#64	#44/4	# c L
	0.75	٠,	0845	0845	#72		
	1.0	٠,	0900	0900	#64	#38/56	
	1.33	11	0920	0920	#84		
	1.66	.	0945	0940	# 80		
	2.00	٠,	1000	1000	#92	#48/58	
	2.5	٠,	1030	1030			
	3.0	٠,	1100	0011	#84		
	3.5	وم	1130	1130		,	
	4.0	٠,	1200	1200	#92	#45/53	
	5.0	٧.	1300	1300			
	6.0	٠.	. 1400	1400	*84	# 44/49	
	7.0	v,	1500	1200			
	8.0	٧.	1600	1600			
	10.0	11	1800	1800			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RUF FM L	٥١	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
S	25 at 85	0800	NA	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	25085	2.000	2,000	# 82		
	24.0	2600 85	08 <i>0</i> 7	0800	# _{&} 2	#49/m	#
						,	
	-						
	-						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R C F F M L	01	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopkins Hoppits

	Screen	Predrug				Stuc
TEST:NORMAL			gqwwwyh Sroq82		25at85	Date
NA:135-148 MEQ/L	146	142		142		
K:3.5-5.0 MEQ/L	4.5	4.4		4.7		
CL:96-109 MEQ/L	107	106		154		
CO2:24-30 MEQ/L	28	28		30		
SUN:12-25 MG/DL	12	15-		1)		
CREAT: 0.4-1.5 MG/DL	1.0	1.2		0.9		
GLU:70-115 MG/DL	87	71		85		
T.BILI:0.3-1.2MG/DL	0.5	0.2		0.3		
D. BILI:0.1-0.4MG/DL	0.0	0.1		0.0		
CA:9.0-11.0 MG/DL	9.7	9.7		9.9		
PO4:3.0-4.5 MG/DL	2.5	1.9		27		
URIC A:4.2-8.8MG/DL	5.0	5-7		4.0		
T. PROT:6.0-8.5G/DL	6.9	69		6.9		
ALB.:3.2-5.3 G/DL	4.3	4.4	٠	4.4		
AST:0-35 IU/L	19	17		25		
ALT:0-30 IU/L	23	17		28		
ALK PHOS:0-95 IU/L	66	רר		69		
CHOL:151-268 MG/DL	ND	C4		234		
LDH:0-200 IU/L	ND	ND 136		141		
	+	0.0	1			

75

83

61

ND

CPK:0-160 U/L(male)

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R C F F M L	01	PROTOCOL DAMD 17-85-C-5133-02

CHEMIBTRY VALUEB
Laboratory Johns Hopetal

Screen Predrug

Study

TEST:NORMAL	ddmmmyy			Date
NA:135-148 MEQ/L		142		
K:3.5-5.0 MEQ/L		4.2		
CL:96-109 MEQ/L		104		
CO2:24-30 MEQ/L		Jo		
SUN:12-25 MG/DL		14		
CREAT:0.4-1.5 MG/DL		0.8		
GLU:70-115 MG/DL		93		
T.BILI:0.3-1.2MG/DL		0.3		
D. BILI:0.1-0.4MG/DL		0.1		
CA:9.0-11.0 MG/DL		9.7		
PO4:3.0-4.5 MG/DL		3.1		
URIC A:4.2-8.8MG/DL		4.3		
T. PROT:6.0-8.5G/DL		7.2		
ALB.:3.2-5.3 G/DL		4.2		
AST:0-35 IU/L		19		
ALT:0-30 IU/L		30		
ALK PHOS:0-95 IU/L		ND		
CHOL:151-268 MG/DL		220		
LDH:0-200 IU/L		20		
CPK:0-160 U/L(male)		63		

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	RCF FML	01	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

		Screen	Predrug				Study
TEST	NORMAL	14 octss ddmmmyy	21 Oct 85	230285 ddmmmyy	260285°	ddmmmyy	Date
Ндь	13.9-16.3	15.8	15.5	16.0	15.7		
PCV	41.0-53.0	47.6	45.7	47.7	47.9		
Plt	150-350 K	189	196	198	219		
RBC	4.50~5.90	5.49	5 49	5 58	5.61		
WBC	4500-11000	6000	7600	6700	7600		
Bands	2-6%	S	4	3	7		
Polys	31-76%	45	5 2	27	53		
Eos	1-4%	ίλ	1	2	2		
Bas		1	0	0	0		
Lymphs	24-44%	39	36	33	34		
Atyp Lym		0	0	B	[
Monos	2-11%	11	6	5	3		
Other		0	0	0	0		
Retics	0.5-1.5%	N.D.	0.7	0.7	1.3		
		*					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R C F	01	PROTOCOL DAMD 17-85-C-5133-02

URINALYBIB VOLUEB Laboratory J. Chineal Pharmacology

ruo	edrug
r	edr

Study

TEST	NORMAL	ddmmmyy		240ct 85 ddmmmyy		ddmmmyy	Date
Color/Ap		ND	Je110-	ND.	ND.		
Sp. Gr.		٠,	1.009	1.008	1.018		
рH		•,	6.5	6.0	6.0		
Protein		•	neg	trace	nex		
Ketones		1,	neg	neg	nex		
Occ Bld		٠.	nex	neg	nex		
Bili.			nei	near	rex		
RBC		۲.	١	1	0		
WBC		1.	0	*1	0.2		
Casts		٠.	0	6	0		
Epi. Cel		ι,	١	0	0		
Crystals		ι.	0	0	0		
Bacteria		٠.	Few	0	0		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
21 Oct 85	V		
21 Oct 85 23 Oct 85 26 Oct 85	V		•
26 Oct 85	✓		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R C F F M L	٥١	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

chec

-k	if	none	occurred	X
				1

					
ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	Mild	DEF.	None
	(0-2400)	(0-2400)	☐ Mod	PROB.	Treatment
	(0-2400)	(0-2400)	MOG	DEF. NOT	ireatment
			Sev	UNKNOWN	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
				PROB.	
	(0-2400)	(0-2400)	Mod	POSS. DEF. NOT	Treatment
			Sev	UNKNOWN	Stop test

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E B	02	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	230185	Screening laboratory
-	250ct 85	History, Physical Exam
0	2800785	Admission
2	2900785	P.O.
5	01 Novi65	I.V.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 1/2 pages, for subject 1/2.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	K E L	02	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	25/	10ct	28.
			dd	mmm	VV

Exami

Brent G. Pott

Date of birth

2 Aug 60 2 Aug 60 dd mmm yy

Age

25 yrs

Sex

 \overline{M}

Race

B

	No	Yes	Comments
Allergy		/	Penicillin > hives include
Tobacco Use	V		
Alcohol Use		/	3 cans
Recreational Drug Use		V	3 joints/lut norths.
Medications past 2 weeks	~		3
Experimental Drug Exposure	1		
Blood or plasma donor	~		
Prior Surgery		V	adenoidectory as
Eye, ear, nose, throat	/		
Endocrine(diabetes, thyroid)			
C-V (heart murmur, HBP)			
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal			
Genito-urinary		V	G.C. 10 years ago
Musculoskeletal	V		
Neuropsychiatric	/		
Pesticide/herbicide use		1	not asked.
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E O	02	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 25/00/1/85 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt. (kg)
37.00	_76/min	22/min	7 3	4178	182.5	64.4

	GENERAL EXAMINATION:						
(check)	Nor.	Abn.	Provide details of abnormalities				
Head/Neck	~						
EENT		V	minimal tonus palatinus				
Chest, lungs	>						
Heart	/						
Abdomen	1						
Genitalia			ND.				
Rectal			N.D				
Extremities							
Skin		/	scars on hands from injury atent				
Neurologic	~						

CHEST X-RAY Date 25/QCI/85

NORMAL	X ABNORMAL	Describe	abnormalities:

Examiner

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	KE A F M L	02	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	D1 N1182	0800	0830	IV	24

Syringe + pyride = 41,97817 8 DOSAGE (total) .66 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ UG/ML	RBC ACHe uM/ml/min	
BO1, B47	0	01 00185	0755	0750	B01: *	B47: 14.48	5
B02	0.08	1.	2805	0805	B02: *		
B03	0.16	٧,	0180	0810	воз:4,38		
B04, B48	0.25	٠,	0815	0815	B04: 8.66	B48: 13 53	
B05	0.33	¢,	0820	0820	B05: 6.66		
B06	0.42	•,	0825	0825	BO6: 8,52		
B07, B49	0.50	4	0830	0830	B07: 9.05	B49: 12 27	
BOB	0.58	ε,	0835	0835	BOB: 1.64		
B09	0.66	٠,	0840	08 40	809: ≭		
B10, B50	0.75	٠,	0845	0845	B10:4.73	B50: 12 56	
B11	0.83	τ,	0850	0850	B11:5.92		
B12	0.92	t.	0855	25-5-	B12:2.34		
B13, B51	1.0	١,	0900	0900	B13: 1,42	B51: 13 34	20
B14, B52	1.33	۲.	0920	0920	B14: *	B52:1313	
B15, B53	1.66	٠.	0940	0940	B15: *	B53: ; 3 4.7	
B16, B54	2.0	۲.	1000	1000	B16:*	B54: 13 76	
B17	2.5	l.	1030	1030	B17:*		

* below assay sensitivity 153

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E O	07	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	01 NN85	0830	0830	IV	NA

DOSAGE (total) . 46 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	LC Mr	RBC ACHe uM/ml/min
B18, B55	3.0	01 N=085	1160	1100	B18: *	B55:/5:7/
B19	3.5	١.	1130	1130	B19: 🚜	
B20, B 5 6	4.0	1,	1200	1200	B20: *	B56:
B21	5.0	C ₁	1355	1300	B21: *	
B22 , B57	6.0	٧,	1400	1400	B22: 🚁	B57: 14.80
B23	7.0	٠,	1500	1500	B23: *	
B24	8.0	.,	1600	1600	B24: *	
B25	10.0	41	1800	1800	B25: *	
B26	12.0	11	2000	2000	B26: *	
B27, B 5 8	24.0	02 NIVES	0802	0800	B27: 놡	B58: 15 00
			·			
		l pensit				

			COMPOUND Pyridostigmine
Lietman	K M LD	02	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	0110085	0800	0835	IV	NA

DOSAGE (total) .66 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	01 200 85	0750	0750	#84	#48/52	# OK
	0.8	٠,	0805	0805		,	
	0.16	٠,	0810	0130			
	0.25	٠,	2180	0815	# 68		
	0.33	1.	O825	C870			
	0.42	t,	0825	0825			
	0.50	٠,	0830	0830	#64	# 47/5/	# 200
	0.58	1,	2830	0837			
	0.66	٠,	0840	0840			
	0.75	٠,	0845	0845	#72		
	0.83	14	c 280	0850			
	0.92	••	0855	0855			
	1.0	4.	0900	0900	* 68	*38/47	
	1.33	l.	0920	0920	* 68		
	1.66	٠,	0940	0940	# 68		
	2.0	• •	1000	1000	# 72	#37/46	
	2.5	0	1030	1030			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	K M L	22	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
S	01 NO185	c080	0835	IV	NA

DOSAGE (total) ______ (6 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	0100185	1100		* 75		
	3.5	١,	1130				
	4.0	١,	1200		* 76	#40 /19	
	5.0		1300				
	6.0	30	1400		*100	#49/52	
	7.0	N	1500				
	8.0	11	1600				
	10.0	l i	1800				
	12.0	1 1	2000				•
	24.0	02 Nov 85			* 112	#49/50	# c.kc.
			·				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E A	02	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	29 Oct 8:	0805	2030	PO	NA

syme + gyride: 6.68746 g.

DOSAGE (total)

20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (9-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B28, B59	0	29 0485	w755	0755	B28: *	B59: 13 93
B29, B60	0.25	٠,	0820	0820	B29: 🗶	B60:14.46
B30, B61	0.50	١,	0835	0835	B30: *	B61:13.83
B31, B62	0.75	٤,	0855	0850	B31: *	B62:14.38
B32, B63	1.0	1	0905	0,905	B32: 🙏	B63: 13.69
B33, B64	1.33	١,	0925	0925	B33:5.75	B64: [∂, 3 ∈
B34, B65	1.66	4.4	0945	0945	B34: 7,22	B65:1132
B35, B66	2.0	1,	1005	1005	B35:4,24	B66: 11.0 Ł
B36	2.5). Ja	1035	1035	B36: 5.50	
B37, B67	3.0	14	1105	1105	B37:5.5	B67: 10.31
B38	3.5	C,	1135	1135	B3B:3,92	
B39, B68	4.0	15	1205	1205	B39:6.85	B68:10.20
B40	5.0	45	1305	1305	B40: 2.76	
B41,B69	6.0	<i>'</i>	1405	1405	B41: 5.36	B69: / / 3
B42	7.0	٠.	1505	1505	B42: 2,34	
B43	8.0	^	1605	1605	B43: *	B69A: N. D.
B44	10.0	4,	1805	1805	B44: *	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	F M L	02	PROTOCOL _ DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	29 cct85	0805	0805	PO	NA

DOSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO)	RBC ACHe uM/ml/mir
B45	12.0	29 oct 85	2005	2005	B45: *	
B46, B70	24.0	30 citas	0805	0806	B46: *	B70: 14.4 0
	in the second					
	4					
			·			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	K M L	02	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	296488	0805	2080	PO	2

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	290285	0755	0755	# 84	#41/45	# GK
	0.25	1,	0820	೧೯೨೦	03*	/	
	0.50	' 、	0835	0832	* 72	35/46	# c.k
	0.75	ι,	0.530	0850	#64		
	1.0	4 ,	0905	0905	* 68	39/50	
	1.33	k.	0925	0925	#64		
	1.66	1,	0945	0945	#60		
	2.00	٠,	1005	1005	# 60	#3 7/50	
	2.5	l,	1035	1035			
	3.0	٠.	1105	1105	# 60		
	3.5	L _s	1135	1135			
	4.0	٠,	1205	1205	#74	#31 X4	
	5.0	t,	1305	1305			
	6.0	1,	1405	1405-	* 82	#42/49	
	7.0		1505	1505		7	
	8.0	١.	1605	1605			
	10.0	١.	1805	1805			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	نك	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	29 octs	0805	0805	PO	NAI

DOSAGE (total) 20mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	29 Cct 85	2005	2005	# N.D.		
	24.0	30 Oct 85	7000	0806	* 85	#41 /45	#
						,	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K M L	02	PROTOCOL DAMD 17-85-C-5133-02

CHEMIBTRY VALUEB
Laboratory Johns Hopkins Hopkins

	Screen	Predrug			
TEST:NORMAL	23 Oct 8 o		29 0885	ddmmmyy	ddmmmyy
NA:135-148 MEQ/L	20	140	_	142	
K:3.5-5.0 MEQ/L	ND	4.0		4.5	
CL:96-109 MEQ/L	20	106		108	
CO2:24-30 MEQ/L	20	27		28	
SUN:12-25 MG/DL	13	16		14	•
CREAT:0.4-1.5 MG/DL	1.1	1-1		1.1	
GLU:70-115 MG/DL	72	71		83	
T.BILI:0.3-1.2MG/DL	0.4	1-2		0.7	
D. BILI: 0. 1-0. 4MG/DL	0.6	0.6		00	
CA:9.0-11.0 MG/DL	10 2	9.8		9.8	
PO4:3.0-4.5 MG/DL	3.6	3,7		4.0	
URIC A:4.2-8.8MG/DL	5.6	5.7		5.1	
T. PROT:6.0-8.5G/DL	7.6	7.3		69	
ALB.:3.2-5.3 G/DL	5.0	4.8		4.6	
AST:0-35 IU/L	28	23		21	
ALT:0-30 IU/L	26	19		14	
ALK PHOS:0-95 IU/L	61	62		55	
CHOL:151-268 MG/DL	199	175		174	
LDH:0-200 IU/L	ND.	134		118	
CPK:0-160 U/L(male)	NO	228	140	120	116

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E D	02	PROTOCOL DAMD 17-85-C-5133-02

CHEMIBTRY VALUES
Laboratory Johns Hopkins Hospital

Savage	Dyadsun

Study

TEST: NORMAL	adwwwyy - NB	ddmminyy	OSNN82	ddmmmyy	ddmmmyy
NA:135-148 MEQ/L			145		
K:3.5-5.0 MEQ/L			4.1		
CL:96-109 MEQ/L			101		
CO2:24-30 MEQ/L			24		
SUN:12-25 MG/DL			19		
CREAT: 0.4-1.5 MG/DL			1.2		
GLU:70-115 MG/DL			78		
T. BILI: 0. 3-1. 2MG/DL			1.2		
D. BILI:0.1-0.4MG/DL			00.		
CA:9.0-11.0 MG/DL			N.D		
PD4:3.0-4.5 MG/DL			3.5		
URIC A:4.2-8.8MG/DL			5.6		
T. PROT:6.0-8.5G/DL			8.1		
ALB.:3.2-5.3 G/DL			S.D		
AST:0-35 IU/L			22		
ALT:0-30 IU/L			18		
ALK PHOS:0-95 IU/L			57		
CHOL:151-268 MG/DL			205		
LDH:0-200 IU/L			131		
CPK:0-160 U/L(male)			128		

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	K E B	07	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory __ Johns Hopking Hospital

Screen	Predrug

Study

Date

TEST	NORMAL	230485	280ct 85 ddmmmyy		OZNW82 OZNW82	ddmmmyy	
Hgb	13.9-16.3	13.9	13.6	14.1	15.0		
PCV	41.0-53.0	41.0	41.2	41.7	44.6		
Plt	150-350	362	391	375	410		
RBC	4.50-5.90	4.63	4.81	4.82	5.22		
WBC	4500-11000	6100	6200	6600	5700		
Bands	2-6%	1	1	8	11		
Polys	31-76%	42	47	35	38		
Eos	1-4%	7	8	10	3		
Bas		0	0	(ì		
Lymphs	24-44%	44	38	41	37		
Atyp Lym		0	0	l	0		
Monos	2-11%	6	6	٩	0		
Other		0	0	0	9		
Retics	0.5-1.5%	ND	0.9	0.9	1.0		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	02	PROTOCOL DAMD 17-85-C-5133-02

URINALYBIS VALUES
Laboratory Johns Hopking Hospital on
Clinical Pharmacology

Screen Predrug

Study

TEST	NORMAL	250tss ddmmmyy	280365 ddmmmyy			ddmmmyy	Date
Color/Ap		ND	2.0	0.0	ND.		
Sp. Gr.		1.023	1.027	1.023	NJ		
pН		6.5	6.0	6.0	6.0		
Protein		Troce	tioce	ncey	200		
Ketones		nes	nea	new	mex		
Occ Bld		2	ner	201	~~~		
Bili.		nev	nes	2 ×	nex		
RBC		0	0	9	nane		
MBC		0	\	9	nane		
Casts		6	0	0	0		
Epi. Cel		0	0	-	0		
Crystals		0	9	0	0		
Bacteria		0	<u>ي</u>	0	0		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
25 Oct 85		/	non-specific ST-T changes
28 Oct 85		/	non- specific ST-Tchrogs, nin. change between
30 Oct 85		V	non specific ST-Tchanges min change between
2 NOV 95		$\sqrt{}$	non-specific IT-Tchanges, min change between

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	K E A	02	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

ADVERSE EXPERIENCES

check if none occurred

/	/	ı
7		ı
	1	V
	>	X

ORUS DOS	- ANDET	SESSOTION	GENERATY	DELIN TO	OCTION
ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		TEST DRUG	(Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
				PROB.	
	(0-2400)	(0-2400)	Mod	Poss.	Treatment
	(0-2400)	(0-2400)	Mod		reatment
				DEF. NOT	
			Sev	UNKNOWN	Stop test drug
#					
	dd mmm yy	dd mmm yy	Mild	DEF.	None
				PROB.	
	(0-2400)	(0-2400)	Mod	POSS.	Treatment
				DEF. NOT	
			Sev	UNKNOWN	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	24007 85	Screening laboratory
_	25 OCT 85	History, Physical Exam
0	28 00185	Admission
2	29 00.785	I.V.
5	01 NOV 85	OPAL

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AIG	03	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation 25/OCT/85
		dd mmm yy

Examiner _

breut Potterty
Brent G. Petty

Date of birth

26/JUN58

Age

27yrs

Sex

 \underline{M}

Race

B

	No	Yes	Comments
Allergy		/	nay fever, dust conjection of AM resolver sponteneously into 15
Tobacco Use		V	1/2 782
Alcohol Use		~	rarely
Recreational Drug Use		V	marijauna tried 714 yrago
Medications past 2 weeks	V		
Experimental Drug Exposure		V	"Pharma-Kinetics" 2-3 months ago was last; 2 beforethet
Blood or plasma donor		/	12001 4-3 grs. cfs
Prior Surgery		/	R) achilles tender repair 10/84 Skin creft to both hand afterly
Eye, ear, nose, throat		/	skin grafts to both hands after to hay fever / congestion on above
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	~		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	/		
Genito-urinary		/	60~ 10 yrs ego-Px 8CN
Musculoskeletal	/		
Neuropsychiatric	/		
Pesticide/herbicide use			Not asked
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AILG	03	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 10/0ct/85

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt.(kg)
<u>37.2</u> c	_ 9 Q/min	20min	_1 3	0/70	175.2	89.9

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck			
EENT			
Chest, lungs			Not done
Heart	/		
Abdomen	V		
Genitalia			Not done
Rectal			Not done
Extremities	/		
Skin	~		Scar (Dlowerleg, on hands
Neurologic			

CHEST X-RAY
Date 25/007/85

NORMAL	X	ABNORMAL	Describe	abnormalities:
			•	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	A T G F M L	03	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	29 0285	0800	0830	IV	NA

sognige + pyride = 41.98 DOSAGE (total) . (46 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	EPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B01,B47	0	29 oct 85	0755	0755	B01: *	B47:12.59
Bos	0.08	••	0805	0805	B02:5,59	
B03	0.16	4.	0810	2810	воз: 5,90	
B04, B48	0.25	ā ,	0815	0815	B04:8,81	B48: 12.06
B05	0.33	L _c	0820	0820	B05:7,89	
B06	0.42	45	0825	0825	BOE: 9.85	
B07, B49	0.50	ł c	0830	0830	B07:/0,7	B49:11.28
BOB	0.58	4.	0835	0835	BOB: 5.96	
B09	0.66	٠,	08 40	0845	809:4,21	
B10, B50	0.75	t q	0845	0845	B10: 2,74	B50:10.49
B11	0.83	14	0850	0850	B11: 2.52	
B12	0.92	41	3855	0855	B12:2.49	
B13, B51	1.0	٠,	0900	0900	B13: 🐥	B51:11.74
B14, B52	1.33	• •	0920	0920	B14: 💥	B52:11.66
B15, B53	1.66		0940	0940	B15: 🙏	B53:12.22
B16, B54	2.0	4,	1000	1000	B16: *	B54:12.60
B17	2.5	3,	1030	1030	B17:	

* below assay sensitivity 169

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	29088	0800	0830	IV	NA

DOSAGE (total) . 66 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ UC/ML	RBC ACHe uM/ml/min
B18, B55	3.0	290285	1100	1100	B18: *	B55: 12.71
B19	3.5	٤.	1130	1130	B19: *	
B20, B56	4.0	ι,	1200	1200	B20: *	B56: 12.26
B21	5.0	ŧξ	1300	1300	B21: *	
B22, B57	6.0	١,	1400	1400	B22: *	B57:
B23	7.0	Ly	1500	1506	B23: *	
B24	8.0	I,	1600	1600	B24: *	
B25	10.0	11	1800	1800	B25: 🟃	
B26	12.0	11	2000	2000	B26: ★	
B27, B58	24.0	30 Oct 85		0804	B27: *	B58: 13.34
			13			

* below assay sensitivity 170

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	29 00 85	0800	0830	IV	NA

DOSAGE (total) ____. 46 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	2900485	0755	0755	#70	#30/33	# ok
	0.8	٠,	0805	0805			
	0.16	4	0810	0810			
	0.25	4,	0815	0815	# 68		
	0.33	4(0820	0820			
	0.42	e _q	0825	0825			
	0.50	t _d	0830	0830	#64	#37/42	#ok
	0.58	81	9832	0835			
	0.66	٠,	0840	0840			
	0.75	ų	0845	0845	# 68		
	0.83	44	0850	0850			
	0.92	14	2822	0855			
	1.0	4,	0900	copo	# 68	#33/46	
	1.33	٠,	0920	0920	#66		
	1.66	٧.	0940	0940	#62		
	2.0	٠,	1000	1000	#70	#30/41	
	2.5	۱.	1030	1030			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	290085	0800	0830	IV	NA

DOSAGE (total) .66 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	290285	1100	1100	# 68		
	3.5	l s	1130	1130			
	4.0	1,	1500		#66	#44/39	
	5.0	٠.	1300	1300			
	6.0	11	1400	1400	# 75	# 46/44	
	7.0	V (1500	1506			
	8.0	14	1600	1600			
	10.0	فر	1800	1800			
	12.0	()	2000	2000			
	24.0	78£00E	080	0804	#92	#33/27	#ok
			,				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	0111185	0805	0805	PO	NA.

DUSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	LPYRIDOJ UG/mL	RBC ACHe uM/ml/min	
B28, B59	0	010015	0745	0745	B28: *	B59: 3.35	35
B29, B60	0.25	ć q	0830	0820	829: ★	B60: 13.26	
B30, B61	0.50	45	2835	0835	B30:-★	B61: 13.08	
B31,B62	0.75	١,	0856	0850	B31:6,80	B62:12.90	
B32, B63	1.0	•(0905	0905	вз2:6.07	B63: 11.81	
B33, B64	1.33	4	0925	0925	взз: 725	B64: 11.06	
B34, B65	1.66	4.6	0945	0945	B34:7.49	B65: 10.60	55
B35, B66	2.0	١,	1005	1005	вз5:9,82	B66: 10.09	
B36	2.5	4	1035	1035	B36:10.7		
B37, B67	3.0	٤,	1105	1105	B37:12.0	B67: 9.64	
B38	3.5	14	1135	1135	B38:10.7		
B39, B68	4.0		1205	1205	B39:/0,5	B68: 10.10	
B40	5.0	t.	1305	1305	B40: ¥		
B41,B69	6.0	٠,	1405	1405	B41:3.36	B69:11.12	
B42	7.0	4	1505	1505	B42: 2.13		
B43	8.0	٠.	1605	1608	B43:255	B69A: N.D.	
B44	10.0	(1805	1805	B44: *		

* Velow assay sensitivity 173

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	ATG	03	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	01228	0805	0805	PO	28

DOSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO)	RBC ACHe uM/ml/min
B45	12.0	280010	2005	2005	B45: *	
B46, B70	24.0	OZ NOVES		0805.	B46:*	B70: 13.20
		٠				
		_				
			•			
* bei	ow awa	ry Deno	litivity 170	4		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	ATG	03	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	01 NW85	0805	0802	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	0110085	0745	0745	#64	#40/37	# ok
	0.25	٠,	0820	0820	#78		
	0.50	4.4	0835	835	#64	#37/45	#0E
	0.75	٠,	0850	0850	# 68	,	
	. 1.0	44	0905	0905	# 7 ₀	#36/S3	
	1.33	t,	0925	0925	# 64		
	1.66	۸,	0945	0945	# 64	*	
	2.00	Ĺţ	1005	1005	# 60	#40/44	
	2.5	C,	1035	1035		,	
	3.0	i,	1105	1105	#60		
	3.5		1135	1135			
	4.0	٠,	1205	1205	#56	#42 /40	
	5.0	4	1305	1305			
	6.0	4.	1405	1405	#7 0	#44/44	
	7.0	۲.	1505	1505			
	8.0	<i>t</i> .	1605	1608			
	10.0	• • •	1805	1805			·

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG FML	03	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	OINOY &S	0805	0805	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	011185	2005	2005	#80		
	24.0	02 N~85		0805	#-75	#40/40	ok#
			,				
				-			

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ATG	03	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hypkins Horpital

Screen Predrug

Study

TEST:NORMAL	24 oct 85	280d85	290782	30 Otes	01 Nn85	Da
NA:135-148 MEQ/L	141	139		137		
K:3.5-5.0 MEQ/L	4.5	4.3		4.3		
CL:96-109 MEQ/L	105	100		103		
CO2:24-30 MEQ/L	24	27		26		
SUN:12-25 MG/DL	9	12		14		
CREAT:0.4-1.5 MG/DL	0.9	0.8		0.9		
GLU:70-115 MG/DL	88	86		83		
T.BILI:0.3-1.2MG/DL	0.4	0.4		0.4		
D. BILI:0.1-0.4MG/DL	0.0	0,0		0.0		
CA:9.0-11.0 MG/DL	9.7	9-8		9.7		
PO4:3.0-4.5 MG/DL	3.9	3.8		4.3		
URIC A:4.2-8.8MG/DL	3.6	3.3		3.8		
T. PROT:6.0-8.5G/DL	7.1	7.2		6.8		
ALB.:3.2-5.3 G/DL	4.5	4.4		4.5		
AST:0-35 IU/L	22	16		15		
ALT:0-30 IU/L	15	20		17		
ALK PHOS:0-95 IU/L	80	84		79		
CHOL:151-268 MG/DL	CN	193		220		
LDH:0-200 IU/L	ND	93		83		
CPK:0-160 U/L(male)	ND	19]	145	125	119	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AT F	83	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopkins Horgital

Screen Predrug

Study

NA NA 02 NWEST Date ddmmmyy ddmmmyy ddmmmyy ddmmmyy TEST: NORMAL NA:135-148 MEQ/L 137 4.0 K:3.5-5.0 MEQ/L CL:96-109 MEQ/L 98 CO2:24-30 MEQ/L 26 SUN:12-25 MG/DL 12 CREAT: 0.4-1.5 MG/DL 0.9 GLU:70-115 MG/DL 84 T. BILI: 0.3-1.2MG/DL 0.6 D. BILI: 0. 1-0. 4MG/DL 0.1 CA:9.0-11.0 MG/DL ND PO4:3.0-4.5 MG/DL 4.7 URIC A:4.2-8.8MG/DL 3.7. T. PROT: 6. 0-8.5G/DL 7.6 4.9 ALB.:3.2-5.3 G/DL AST:0-35 IU/L 17 ALT:0-30 IU/L 22 ALK PHOS:0-95 IU/L 74 CHOL: 151-268 MG/DL 220 LDH:0-200 IU/L 84 CPK:0-160 U/L(male) 137

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	A T G F M L	03	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOBY VALUES Laboratory Johns Hopking Hospital

		Screen	Predrug				Study
TEST	NORMAL	24 0d 85 ddmmmyy	280185 ddmmmyy			ddmmmyy	Date
Ндь	13.9-16.3	13. 4	13.4	13.6	13.3		
PCV	41.0-53.0	41.0	40.9	45.0	41.1		
Plt	150-350	286	263	289	271		
RBC	4.50-5.90	4.96	4.86	5.57	4.89		
WBC	4500-11000	5300	6800	6400	6900		
Bands	2-6%	5	3	10	0		
Polys	31-76%	56	5 3	47	66		:
Eos	1-4%	4	8	8	9		
Bas)	1	J	0		
Lymphs	24-44%	29	27	22	26		
Atyp Lym		0	2	0	0		
Monos	2-11%	5	6	12	8	-	
Other		0	0	Ð	0,,		
Retics	0.5-1.5%	N.D.	1.6	1.7	2.0		
		-					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A76 FML	03	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Johns Hooking Hospital CN Clinical Pharmacology
een Predrug

Screen Predrug

Study

TEST	NORMAL	250d85 ddmmmyy		300t85			Date
Color/Ap		ND	ND	ND	619		
Sp. Gr.	,	1,021	1.014	1.018	20		
рН		6.5	6.0	6.5	6.0		
Protein		trace	100	ner	~~		
Ketones		necex	ney	7-99	ner		
Occ Bld		neg	nex	ney	nes		
Bili.		reg	res	res	ner		=
RBC		1	0	0	raine		
WBC		0	0	2	0-1		
Casts		0	S	0	0		
Epi. Cel	•	0	0	0	occapana	L	
Crystals		D	0	O	0		
Bacteria		0	0	D	0		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
250t85	w		
28 0 185	V		
30 ot85	V		
02 NW85	v		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AIG	03	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

	VERSE	ONSET (Date/Time)	CESSATION Date/Time	Attendance of the second	REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB.	 Treatment
				Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
				 Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ARK	04	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	18NOV 85	Screening laboratory
_	19 NOV 85	History, Physical Exam
0	211185	Admission
2	22 NOV 85	P.O.
5	25 NOV 85	I.V.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject $#_{04}$.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	ARK	04	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	19,	1 NW	85
			dd	mmm	УУ

Examiner

Date of birth

14/Sep/60 dd mmm yy

Age

25 yrs

Sex

<u>m</u>

Race

	No	Yes	Comments
Allergy			
Tobacco Use		V	1887
Alcohol Use		V	6 pack / weekend
Recreational Drug Use		1	mg g2 weeks
Medications past 2 weeks	1		
Experimental Drug Exposure		V	Tobramyein, Interfora
Blood or plasma donor			last 1984
Prior Surgery	~		
Eye, ear, nose, throat	1		
Endocrine(diabetes,thyroid)	V		
C-V (heart murmur, HBP)	1		
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal	/		
Genito-urinary	~		·
Musculoskeletal		/	fx hip as child
Neuropsychiatric	V		
Pesticide/herbicide use	V	Z.	
Other Pets		/	Birds, dogs & recent Py For Fleas

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	ARK	04	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 19/N0/85 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt.(kg)
<u>37.8</u> c	_ 68/min	2Q/min	_L _Q	<u>6/58</u>	174.0	63.5

	GENERAL EXAMINATION:								
(check)	Nor.	Abn.	Provide details of abnormalities						
Head/Neck	/								
EENT		/	some singivities mild arterialer narrowing						
Chest, lungs	~		8						
Heart	V								
Abdomen	~								
Genitalia			ND						
Rectal			ND						
Extremities		/	Scar (1) fream mild antecubith Scarring from studies + blood don't						
Skin			0						
Neurologic	1								

CHEST X-RAY
Date \G\DoV &S

NORMAL	X	ABNORMAL	Describe	abnorma	liti	es:		
			-	1				

Brent G. Petty

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	ARK FML	٥٧	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 No.85	0850	0920	IV	NA

Syringe + Byride = 41.38393 g DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B01, B47	0	25 Na 85	0845	0845	B01: *	B47:13.20
B02	0.08	٠.	0855	0855	BO2:15,4	
B03	0.16	٠,	0900	0900	B03: 24.5	
B04, B48	0.25	k.,	0905	0905	B04:27.1	B48:10.54-
B05	0.33	١,	0910	0910	BO5:30,5	
B06	0.42	1,	0915	0915	BOE:37.6	
B07, B49	0.50	4.5	0920	0920	B07:375	B49:8.50
B08	0.58	٠,	0925	0925	BO8: /8.6	
B09	0.66	l.	0930	0930	B09: /3,4	
B10, B50	0.75	4,5	0935	0935	B10:/2,0	B50: 9.70
B11	0.83	ŧ,	0940	0940	B11:10,3	
B12	0.92	4.1	0945	0945	B12: 850	
B13, B51	1.0	t,	0950	0950	B13:8.34	B51:10.12
B14, B52	1.33	1,	1010	1010	B14:5,87	B52://.0/
B15, B53	1.66	1,	1030	1030	B15:3,80	B53: 10.02
B16, B54	2.0	1,	1050	1050	B16:3,40	B54: 10. 7.7
B17	2.5	11	1120	1120	B17:3.08	

* below assay sensitivity 185

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A R K F M L	04	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 Naves	0850	0920	IV	NA

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	J5NN85	1150	1150	B18:/.5/	B55://.99
B19	3.5	1.	1220	1220	B19: *	
B20, B56	4.0	1,	12.50	1250	B20: *	B56: _{12.41}
B21	5.0	4.4	1350	1350	B21: N.S.	
B22, B57	6.0	1,	1450	1450	B22: 1.67	B57: 12.55
B23	7.0	1,	1550	15-50	B23: ★	
B24	8.0	i ,	1650	1650	B24: 💥	
B25	10.0	i,	1850	1850	B25: ★	
B 26	12.0	1.	2050	2050	B26:*	
B27, B58	24.0	26Nov87	0850	0850	B27: *	B58: _{12.59}
					e e	

* Inlaw assay pensitivity N.S. = no Sample

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A N K	04	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	25 Nw85	0850	0920	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	25 N ~ 81	0845	0845	#75	#33/39	#ok
	0.8	١,	0855	0855			
	0.16	14	0900	0900			
	0.25	٠,	0905	0905	#74		
	0.33	*4	0910	0910			
	0.42	41	0915	0915			
	0.50	*	0925	0925	# 68	#38/40	#ck
	0.58	٠,	0925	0925			
	0.66	6 q	0930	0935			
	0.75	4.	0935	0935	# 68		
	0.83	**	0940	0940			
	0.92	6.4	0945	0945			
	1.0	16	0950	0950	# 70	# 35/32	
	1.33	14	1010	1010	#66		
	1.66	1, 3	1030		#72		
	2.0	1.	1050	1050	# 74	#36/32	
	2.5	ı _i	1125	1120		·	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ARK	04	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 No 185	0850	0920	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	25 Noves	11.50	1150	#74	·	
	3.5	٠.	1220	1220			
	4.0	٠,	1250		# 60	#32/34	
	5.0	• 4	1350	1350			
	6.0	M	1450	1450	#68	#25/36	
	7.0	14	1550	1550			
	8.0	•	1650	1650			
	10.0	٠,	c281	1850			
	12.0	.,	2050	2050			
	24.0	26NW85		0230	#70	#41/40	# ok.
						/	
				,			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A R K	04	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	35NW82	0845	0845	PO	NA

sorginge + pyride = 6.72348 orgunge alore = 4.64543

DOSAGE (total)

20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B28, B59	0	22 Nov 85	0345	0845	B28: *	B59: 13.87
B29, B60	0.25	٠.	0900	0900	B29:*	B60: 13.39
B30, B61	0.50	٠,	0915	0915	B30: *	B61: /3.98
B31, B62	0.75	٠,	0935	0930	B31:2.23	B62:/3.23
B32, B63	1.0	٠,	0945	0945	B32: 7.93	B63://. 79
B33, B64	1.33	4,	1005	1005	взз: 853	B64:10.98
B34, B65	1.66	* ₁ =	1025	1025	B34:11.2	B65:9.77
B35, B66	2.0	٠,	1045	1045	B35:16.5	B66:9.02
B36	2.5	1,	7115	1115	B36:17.1	
B37, B67	3.0		1145	1145	B37: /8.8	B67: 8.30
B38	3.5	£ ,	1215	1215	B38:18.0	
B39, B68	4.0	4.	1245	1245	B39: 20.3	B68:7.57
B40	5.0	/ =	1345	1445*	B40: 14.7	
B41, B69	6.0	e,	1445	1445	B41: /0./0	B69: 9.63
B42	7.0	1.	1545	1545	B42: 7,90	
B43	8.0	fi	1645	1645	B43:7.35	B69A://.23
B44	10.0	I,	1845	1840	B44:3,10	

* Lebow assay sensitivity 189 * copied from original workshort

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	ARK	04	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	23 NN 85	0845	0845	PO	NA

DOSAGE (total) 20 mg

PLASMA CONCENTRATIONS

page 2

Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
12.0	27 Nove 5	2045	2045	B45: /.86	
24.0	230085	0845	0845	B46:2,77	B70: 13.5.7
			·		
	Time (hours)	Time (hours) 12.0 21NNSS 24.0 23NNSS	Time (hours) ddmmmyy Clock Time (0-2400) 12.0	Time (hours) ddmmmyy Clock Time (0-2400) 12.0 21 Naes 2045 2045 24.0 23 Naes 0845 0845	Time (hours) ddmmmyy Clock Time (0-2400) NG/ML 12.0 21 Nass 2045 2045 B45: /86 24.0 330085 0845 0845 B46: 2,77

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	ARK	04	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	22 N n 85	0845	084.5	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
-	0	23100085	0845	0845	#80	#3 5/38	# olc
	0.25	· 14	0900	0900	# 74		
	0.50	C ₂	0915	0915	#74	#26/36	#ok
	0.75	4.	0935	0930	* 76	,	
	1.0	٠,	0945	0945		25/29	
	1.33	•,	1005	1005	#72	/	
	1.66	t _e	1025	1025	#62		
	2.00	· · ·	1045	1045	# 60	#26/34	
	2.5	14	1115	A\$1115			
	3.0	· .	1145	1145	# 70		
	3.5	15	1215	1215			
	4.0	14	1245	1245	#74	#27/29	
	5.0	t.	1345	1345			
	6.0	١,	1445	1445	* 76	#25/26	
	7.0	١,	1545	15 45 1645 LA	*		
	8.0	٠.	1645	1645			
	10.0	١,	1845	1845			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A A K F M L	04	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	22 NW85	0845	0845	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	23Nm85	2045	2045	# 74		
	24.0	23101185	2045	0845	#75	#28/29	ck#
			-				
			,				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A A L F M L	04	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopking Hospital

Screen Predrug

Study

21Nov85 23Nov 85 26Nov85 Date 18 NW 85 TEST: NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy NA:135-148 MEQ/L ND 139 137 142 K:3.5-5.0 MEQ/L 4.0 4.3 ND 4.6 CL:96-109 MEQ/L ND 104 102 100 CO2:24-30 MEQ/L 24 22 (Li 26 SUN: 12-25 MG/DL 11 21 16 20 CREAT: 0.4-1.5 MG/DL 10 10 1.0 1.0 GLU:70-115 MG/DL 68 141 111 117 T. BILI: 0. 3-1. 2MG/DL 1.5 13 1.4 1. 1 D. BILI: 0. 1-0. 4MG/DL 0.1 0.1 0.1 5.0 CA:9.0-11.0 MG/DL 9.8 9.8 9.7 PO4:3.0-4.5 MG/DL 3.8 4.1 4. 3 45 URIC A:4.2-8.8MG/DL 55 5.1 6.2 6.0 T. PROT:6.0-8.5G/DL 7.6 7.17.5 7.2 ALB.:3.2-5.3 G/DL 4.9 4.9 4.7 5.0 AST:0-35 IU/L 23 22 13 17 ALT:0-30 IU/L 8 11 10 10 57 ALK PHOS:0-95 IU/L 5.Top 71 67 60 CHOL: 151-268 MG/DL ND ND GM ND. LDH:0-200 IU/L NB 117 119 117 CPK:0-160 U/L(male) ND 80 83 102

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	A R K	04	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

		Screen	Predrug				Study
TEST	NORMAL	18 Noves	21Noves ddmmmyy	9182	36 Nov87		Date
Ндь	13.9-16.3	14.7	14.6	15.3	14.4		
PCV	41.0-53.0	43.6		45.8	429		
Plt	150-350	267	263	273	271		
RBC	4.50-5.90	4.14	4.61	4.79	4.48		
WBC	4500-11000	5000	6900	4900	5000		
Bands	2-6%	0	9	10	اده	,	
Polys	31-76%	76	67	44	45		
Eos	1-4%	1	0	Ч	Q		
Bas		0	0	O	<u>ي</u>		
Lymphs	24-44%	17	16	31	3;		
Atyp Lym		0	O.	2	1		
Monos	2-11%	6	8	9	U		
Other		0	0	0	0		
		÷					
Retics	0.5-1.5%	int de	1.4	1.3	1.2		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A CK F M L	04	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

URINALYBIB VALUEB
Laboratory Johns High in Clinical Phirmocole gy

		Screen	Predrug			Study
TEST	NORMAL	9 18 17 (182	21 Nov85 ddmmmyy	25 Noves ddmmmyy	26 N (185	Date
Color/Ap		ND	ND	ИД	N.D	
Sp. Gr.		1-001	1.015	1.011	1.016	
pН		6.0	7.0	5.0	5.0	
Protein		Nes	Trace	Trace	troce	
Ketones		Nex	New	Noc	Na	
Occ Bld		Ney	Nei	Nei	Nax	
Bili.		Nex	Nex	New	Noe.	
RBC		0	0	0	ن	
MBC		0	0	O	0	
Casts		0	6	0	O	
Epi. Cel		Ο.	0	O	O	
Crystals		0	Ó	0	Ö	
Bacteria		0	ō.	0	0	
						1

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
19NN 85	V		
23 N N 8 5	V		
26NW85	√		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A A K F M L	04	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

1			1	1	
1				-1	
1	1	^		- 1	
3	/		-	J	
/1	_	_	_	_	`

	OVERSE	ONSET	CESSATION			ACTION
	EVENT	(Date/Time)	Date/Time	(Check)	TEST DRUG	(Check)
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	☐ Mod	PROB. POSS.	Treatment
	~			Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M K	05	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	19 NOV 85	Screening laboratory
_	20 NOV85	History, Physical Exam
0	21 No18	Admission
ಇ	22 NOV85	P.6.
5	22 NOV 85	I.V.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	FML	05	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	301	Nov,	<u>18</u> 5
			dd	mmm	УУ

Examiner

Brent G. Potty

Date of birth

all/Arc/63

print name

Age

22 yrs

Sex

 $\overline{\nu}$

Race

B

	No	Yes	Comments
Allergy	1	•	
Tobacco Use		V	T gk. / month
Alcohol Use		/	6 pk/month
Recreational Drug Use		V	mg, last exposure last your
Medications past 2 weeks	·		Tylend Tweek 150 for 41A
Experimental Drug Exposure	V		0
Blood or plasma donor	V		
Prior Surgery		~	I+Dax- @ 3rd mcPjoint 19:2
Eye, ear, nose, throat	1		
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	~		
Genito-urinary	/		
Musculoskeletal	/		
Neuropsychiatric	V		
Pesticide/herbicide use	~		
Other Pets			dogs 5 Fler Powder

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	E M L	<u>٥</u> 5	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 20/100/85 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt. (kg)
37.40	_ 9 _Ymin	⊋ ⊴⁄min	-1 -L	0176	167.6	60.3

	GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities					
Head/Neck	~							
EENT		V	51 1 @ tons, 1 5 erythoma or exidete tonis galatinus					
Chest, lungs	1							
Heart	~							
Abdomen	/		\					
Genitalia			N.D.					
Rectal			\mathcal{N} . \mathcal{D} .					
Extremities	~							
Skin		~	Scarl shoulds over @ 3rd mcp, on ablance					
Neurologic	/							

CHEST X-RAY
Date 20/10/85

NORMAL	×	ABNORMAL	Describe	abnormalities:	

Examiner

Brent F. Petty

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	05	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 NN 85	0855	0925	IV	NA

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG /AL	RBC ACHe uM/m1/min
B01,B47	0	25 Noves	0855	5850	B01:*	B47:/3.35
Bos	0.08	ι.	0900	0900	B02: 17.3	
B03	0.16	* *	0905	0905	B03:37.1	
B04, B48	0.25	*,	0913	0910	B04:38.7	B48:11.04
B05	0.33	٤.	0915	0915	BO5:38.8	
B06	0.42	1,	0920	0920	BOE: 31.9	
B07, B49	0.50	•	0925	0925	B07:31.3	B49: 8, 89
B08	0.58	t _t	0930	0930	BOB:30.2	
B09	0.66	4,	0935	0935	B09:22.0	
B10, B50	0.75	1,	0945	0940	B10:16.2	B50:9.78
B11	0.83	t,	0945	0945	B11:/6.8	
B12	0.92	4s	0950	0950	B12://.7	
B13, B51	1.0	4.1	0955	0955	B13: //.7	B51:/0-38
B14, B52	1.33	^	1015	1015	B14: 5.74	B52: 10.87
B15, B53	1.66	I,	1035	1035	B15:5.93	B53: 10,59
B16, B54	2.0	4.	1055	1055	B16:4.37	B54://./0
B17	2.5	(1125	1125	B17: 252	
* below	v assay	1 Rensit	wity 200			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M L	05	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 N. 85	0855	0925	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	REPYRIDOS NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	25 Na185	1155	1155	B18:4.56	B55:12.41
B19-	3.5	13	1225	1225	B19: 3.65	
B20, B56	4.0	1,	1255	1255	B50:*	B56: /2.72
B21	5.0	1,	1355	1355	B21: *	
B22, B57	6.0	i,	1455	1455	B55:*	B57: 13.24
B23	7.0	L _\	1555	1555	B23: ₩	
B24	8.0	r,	1655	1655	B24: *	
B25	10.0	1,	1855	1855	B25:2.14	·
B26	12.0	4	2055	2055	B26: *	
B27, B58	24.0	284495	0855	0855	B27:*	B58:12.91

* below assay sensitivity 201

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M L	05	PROTOCOL <u>DAMD_17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 Noves	0855	0955	IV	27

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	25 Nov85	0850	0850	#78	#23/29	# (,15
	0.8	N _n	0900	0950		,	
	0.16	V ,	0905	0905			
	0.25	١,	0910	0910	#64		
	0.33	•	6915	0915			
	0.42	*<	0920	0920			
	0.50	4.	0925	0915	#64	#47/49	# ok
	0.58	4,	0930	0930		,	
	0.66	1,	0935	0935			
	0.75	4.	0940	0940	# 60		
	0.83	C,	0945	0945			
	0.92	١.	0950	0950			
	1.0	• •	0955	0955	# 68	#36/37	
	1.33	٠,	1015	1015	#68		
	1.66	(-	1035	1035	#70		
	2.0	**	1055	1055	#64	# 38/42	
	2.5	i.	1125	1125		,	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	EM K FM L	05	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25NN85	0855	0925	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	2500085	1155	1155	# 68		
	3.5	l	1225	1225			
	4.0	13	1255	1255	# 60	#36/42	
	5.0	. (1355	1355			
	6.0	1	1455	1455	#88	#37/49	
	7.0	(1	1555	1555			
	8.0	րձ	1655	1655			
	10.0	£,	1855	1855			
	12.0	ţı	2055	2055			
	24.0	26NN85	0855	0822	#76	#46/55	# ok

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M L	05	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	22 NOV85	0850	0850	PO	NA

DUSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B28, B59	0	22 Na 85	0800	0800	B28: *~	B59: 13.86
B29, B60	0.25	1,	0905	09100	B29:*	B60: 13.53
B30, B61	0.50	٠,	0920	0925	B30:8,24	B61:12.73
B31,B62	0.75	16	0935	0935	B31:/4.0	B62: /1.83
B32, B63	1.0	1	0950	0950	вза: 19,9	B63: 10,13
B33, B64	1.33	٠,	1010	1010	взз: 22.0	B64:9,17
B34, B65	1.66	4.1	1030	1035	B34: 20.0	B65: 8,87
B35, B66	2.0	1	1050	1056	B35257	B66: 8.42
B36	2.5	.,	1120	1120	B36: 17,1	
B37, B67	3.0	1.	1150	1150	B37:/5,0	B67:9.34
B38	3.5	1.	1220	1220	B38:/3.4	
B39, B68	4.0	1,	1250	1250	вз9:9,42	B68: 9.79
B40	5.0	4	1350	1350	B40: 4.98	
B41,B69	6.0	4	1450	1450	B41:3,73	B69: 12.12
B42	7.0	4,	1550	1550	B42:3.62	
B43	8.0		1650	1650	B43:357	B69A:12.93
B44	10.0	١,	1850	1850	B44:1.41	

* below assay sensitivity 204

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M L	05	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	22 Noves	0850	0220	PO	NA

DOSAGE (total) 20mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO3	RBC ACHe uM/ml/min
B45	12.0	22 N N E S	2050	2050	B45:2.06	
B46, B70	24.0	23 mass	0850	0850	B46: 1.42	B70:14.04
			·			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	F M L	05	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	25 10/0 87	- 28G	0850	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	2211285	0800	0630	#78	#40/45	#
	0.25	t,	0905	0905	# 78		+
	0.50	4 ,	0920	0920	# 80	#35 144	#
	0.75	.,	0935	0935	#82		
	1.0	٠,	0950	0950	# 76	#38/48	
	1.33	1.	1010	1010	# 72		
	1.66	(,	1030	1030	# 72		
	2.00	٠.	1050	1050	#74	#41/48	
	2.5	×, **	1120	1120			
	3.0	ι.	1150	1150	#70		
	3.5	t ,	1220	1220			
	4.0	1,	1250	1250	#78	#49155 #55/49	
	5.0	٠,	1350	1350		de	
	6.0	ζ,	1450	1450	* 70	# 45 MR	
	7.0	٠,	1550	1550		Th	
	8.0	٠,	1650	1650			
	10.0	(1850	1850			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E M L	0.5	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	55100182	0850	087_0	PO	NA

DOSAGE (total) ___

20mg

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	22 NW85	2 050	2050	# 70		
	24.0	230085	0850	0850	#66	#49/54	ok#
		•					
	,						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	EME	05	PROTOCOL _ DAMD 17-85-C-5133-02

CHEMISTRY VALUES Laboratory Jahns Hopkins Hospital -

	Screen	Predrug			SN	Study
TEST:NORMAL	19 Nov85 ddmmmyy		23 NOVES	26 12 12 18 18 18 18 18 18 18 18 18 18 18 18 18	23.NW89	Date
NA:135-148 MEQ/L	143	144	141	139		
K:3.5-5.0 MEQ/L	4.2	4.2	3.6	4.0		
CL:96-109 MEQ/L	106	104	107	108		
CO2:24-30 MEQ/L	23	24	21	26		
SUN:12-25 MG/DL	- 8) t	_10	10		
CREAT:0.4-1.5 MG/DL	0.9	0.9	1, 0	0.9		
GLU:70-115 MG/DL	88	91	120	88		
T.BILI:0.3-1.2MG/DL	0.6	0.9	1. 0	0.6		
D.BILI:0.1-0.4MG/DL	0.1	0.1	0.1	0.1		
CA:9.0-11.0 MG/DL	10.4	10.0	9.8	9.6		
PD4:3.0-4.5 MG/DL	3.7	4.2	3.8	4.1	•	
URIC A:4.2-8.8MG/DL	6.1	5.9	6,0	5.6		
T. PROT:6.0-8.5G/DL	7.4	7.4	6.9	6.6		
ALB.:3.2-5.3 G/DL	5, 1	4.9	4.4	4.1		
AST:0-35 IU/L	17	17	12	14		
ALT:0-30 IU/L	16	1)	12	14		
ALK PHOS:0-95 IU/L	64	68	58	57		
CHOL:151-268 MG/DL	197	ND	20	ND		
LDH:0-200 IU/L	ND	116	99	105		
CPK:0-160 U/L (male)		- 0		, i		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	05	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hogkins Hospital

		Screen	Predrug				Study
TEST	NORMAL	19 NNST	gquummyy 311/9887	admmmyy 33 Nagr		ddmmmyy	Date
Ндь	13.9-16.3	15.0	15.1	14.6	14.3		
PCV	41.0-53.0	45.3	44.8	433	42.6		
Plt	150-350	288	270	274	282		
RBC	4.50-5.90	4.93	4.95	4.73	4.63		
WBC	4500-11000	4800	6200	3700			
Bands	2-6%	0	S	1	5	*	
Polys	31-76%	54	70	56	59		
Eos	1-4%		ı	3	3		11
Bas		C	0	3	0		
Lymphs	24-44%	38	24	36	27		
Atyp Lym		0	(C) 7	6	0		
Monos	2-11%	7	0	3	6		
Other		٥	0	0	0		
Retics	0.5-1.5%	N.D.	1.2	1.2	1.7		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	EMK	05	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Johns Hooking Housted and Clinical Pharmaculagy

Screen Predrug

Study

TEST	NORMAL	201051 ddmmmyy	<u> </u>			Date
Color/Ap		CU	N	NB	UD	
Sp. Gr.		1015	1.018	1.015	1.023	
рH		6.5	6.0	6.5	6,0	
Protein		Trace	Nog.	New	New	
Ketones		nce	Nev	New	2	
Occ Bld		nev	Nex	Nec	200	
Bili.		nec	Nev	22	New	
RBC		<i>S</i>	rare	0	nere	
WBC		0	rane	0	~2~4	
Casts		0	O	0	_ a	
Epi. Cel		0	D	0	0	
Crystals		0	<i>O</i>	0	0	
Bacteria		O	0	9	0	
					a lot of amorphys	

ELECTROCARDIDGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
20 N(185	V		
23 Nov85	V		
36 min 82	V		axis shifted from 0° to 15°

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	EMK	05	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

₹	7	ń	١
λ	(
	\		
	X	X	X

ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R M L	06	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
		Screening laboratory
	21 NOV 85	History, Physical Exam
0	21 NOV 85	Admission
2	22NOV85	P.O.
5	25No185	I.V.
	-	
		·

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

M. D. Investigator's signature

24,0ct,86 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R 1	06	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	311NOV182
------	----	------------	-----------

dd mmm yy

Examiner Suluty

Date of birth

24/ Mar 59 dd mmm yy

Age

2 byrs

Sex

*T*J

Race

Fr.	No	Yes	Comments
Allergy	V		
Tobacco Use			
Alcohol Use	~		Quit 1'12 yes ago
Recreational Drug Use			Smoked MJ
Medications past 2 weeks	/		3
Experimental Drug Exposure		V	"Pharma-kinetics" 3185 Interferon 9185 Coffaziding
Blood or plasma donor	/		
Prior Surgery	/		
Eye, ear, nose, throat	V		
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	/		
Genito-urinary	/		
Musculoskeletal	/		
Neuropsychiatric	/		
Pesticide/herbicide use		/	dog dipped " at veterinarians ~3 months ago + "dog soey" before that
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	R K	06	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 21/Nov/ &5

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt. (kg)
37.40	_78/min	J 8/min	_1 1_	8176	17.1.0	52.2

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	/		
EENT		V	
Chest, lungs	/		
Heart	/		
Abdomen	~		
Genitalia			ND
Rectal			N.D.
Extremities			
Skin		~	tattoos Byper orm, Dupper am, Deforem, Dhank
Neurologic			hyporeflexic diffisely

CHEST X-RAY
Date 05/A06/85

NORMAL	\propto	ABNORMAL	Descri be	abnormalities:	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R K F M L	06	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	25 Novi 85	0935	1005	ΙV	NA

Syrige Ane = 21.40249

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B01, B47	0	25 NN&5	0810	0810	B01:*	B47:14.26
B02	0.08	١,	0945	0940	B02:17.0	
B03	0.16	• •	0945	0945	BO3: 19.9	
B04, B48	0.25	1~	0950	0950	B04:29.5	B48: //. 0 /
B05	0.33	6.4	0955	0955	B05: 28,2	
B06	0.42	l _e	1000	1000	BO6:34.8	
B07, B49	0.50	4,	1.0.0.5	1005	B07: 353	B49:9.65
вов	0.58	• (1010	1010	BOB: 20,4	
B09	0.66	• .	1015	1015	B09: 11.9	
B10, B50	0.75	• • •	1020	1020	B10:/0,/	B50:9.45
B11	0.83	٠,	1025	1025	B11: 9.9/	
B12	0.92	*5	1030	1030	B12: 7,60	
B13, B51	1.0	٠,	1035	1035	B13: 7./8	B51:/0.47
B14, B52	1.33	14	,055	1055	B14: 2.98	B52: 10.93
B15, B53	1.66	11	1115	1115	B15:2.02	B53: //.46
B16, B54	2.0	1.	1135	1135	B16:2.09	
B17	2.5	١.	1205	1205	B17: /.90	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R _ K F M L	06	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 Nov 85	0935	1005	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	25 NNE5	1235	1235	B18:3,20	B55:13.45
B19	3.5	T _A	1305	1305	B19: 2.53	
B20, B56	4.0	٠,	1335	1405	B20: *	B56:/3,//
B21	5.0	١,	1435	1435	B21:*	
B22, B 57	6.0	١,	1535	15-35	B22:*	B57:14.42
B23	7.0		1635	1635	B23:→	
B24	8.0	i i	1735	1035	B24: *	
B25	10.0	4.	1935	1935	B25: *	
B26	12.0	f _s	2135	2135	B26: *	
B27, B58	24.0	26 Noves	0935	0935	B27: *	B58:/4.37
_						
* Irlo	w assay	f sensi	tivity 216			

INVESTIGATOR'S NAM	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R M L	06	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
S	25 Nn 85	0935	1005	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	25 NNES	0810	0810	# 76	#39 /44	# olc
	0.8	1,	0940	0940			
	0.16	٠,	0945	0945			
	0.25	4,	0950	0950	# 72		
	0.33	L,	0955	0955			
	0.42	1,	1000	1000			
	0.50	٠,	1005	1005	#79	#42/39	# ok
	0.58	• ,	10	1010			
	0.66	• •	1015	1015			
	0.75	4	1020	1020	# 76	40/y	
	0.83		1025	1025		£n	
	0.92	'1	1030	1030			
	1.0	Tr.	1035	1035	# 76	#40/42	
	1.33	1.1	1055		#80		
	1.66	·, ·	1115	1115	#76		
	2.0		1135	1135		#40/41	
	2.5	1,	1205	1205			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostipmi</u> ne
Lietman	R M L	06	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	25 NN86	0935	1005	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	25 NN 85	1235	1235	#74	·	
	3.5	1,	1305	1305			
	4.0	.,	1335	1405	# 25	#42/45	
	5.0	1	1435	1435			
	6.0	e,	1535	1535	#80	#45-/39	
	7.0	14	1635	1635			
	8.0	٤,	1735	1735			
	10.0	*4	1935	1935			
	12.0	a _k	2135	2135			
	24.0	26 NN 85	0935	0935	# 90	#44/46	#ok
			2				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	06	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study Day	Date ddmmmyy	Start drug admin- istration (0-2400)	End drug admin- istration (0-2400)	Route	Bottle I.D. #
2	22 NW86	0855	0855	PO	NA

DOSAGE (total)

Spring Love = 4.63942

PLASMA CONCENT

20 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B28, B59	0	22 N N 81	0819	0319	B28: *	B59:14.86
B29, B60	0.25	* 1	0910	0910	B29:4,75	B60: 14.04
B30, B61	0.50	٠.	0925	0925	B30: 8.94	B61:13,19
B31, B62	0.75		0940	0940	B31:/3,8	B62: //.38
B32, B63	1.0	t,	0955	0955	B32: /4,7	B63: //. 98
B33, B64	1.33	ŧ.	1015	1015	B33: 44	B64: //./3
B34, B65	1.66	"	1035	1035	B34:14,5	B65:/0.58
B35, B66	2.0	6 ₀	1055	1055	B35:/0.6	B66:10.93
B36	2.5	K	1125	1125	B36:/0,8	
B37, B67	3.0	ι,	1155	1155	B37:/0,5	B67:/1.26
B38	3.5	1,	1225	1225	взв:9,33	
B39, B68	4.0	١,	1255	1257	B39:8.24	B68: 10.98
B40	5.0	4 ,	1355	1355	B40:4,55	
B41, B69	6.0	Ĭŧ	1455	1455	B41: 3,79	B69: 13.34
B42	7.0	٠.	1555	1555	B42:3,62	
B43 B69A	8.0	13	1655	1655	B43: 3,36	B69A: 13,74
B44	10.0	4.2	1855	1855	B44: *	

* below assay sensitivity 219

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	R K F M L	06	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	22NN85	0.855	0855	PO	NA

DOSAGE (total) 20 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/HL	RBC ACHe uM/ml/min
B45	12.0	25 NN 85	2055	2055	B45:*	
B46, B70	24.0	23 NN85	0855	0822	B46: *	B70: 15, 11
_						
			19			
					·	

* below assay sensitivity 220

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	R _ K	06	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	22 Nov85	0855	0855	PO	NA

DOSAGE (total) 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	22 NNES	0819	0819	#84	#44/45	# ck
	0.25	٠.	0910	0910	# 77	,	• •
	0.50	٠,	0925	0925	#63	#41/42	# ok
	0.75	*1	0940	0940	#72		
	1.0	ř _š	0955	0955	# 65	#4 Jyg	
	1.33	4.5	1015	1015	# 69	/	
	1.66	e ș	1035	1035	#70		
	2.00	<i>i</i> ,	1055	1055	#75-	#33/44	_
	2.5	1,	1125	1125		,	
	3.0	A	1155	1155	#79		
	3.5	1.	1225	1225			
	4.0	٧.	1255	1257	#68	#45/42	
	5.0	٠.	1355	1355			
	6.0	7.	1455	1455	* 50	#38/40	
	7.0	4	1555	1555-	×	/	
	8.0	1.	1655	1655			
	10.0	Ĉ ₃	1855	1855			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	P _ JS F M L	06	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	22 N~ &	0855	0855	PO	NA

DOSAGE (total) _ 20 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	22NN85	9027	2055	#87		
	24.0		7280	2055	#80	#41/42	cic#

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R_K FML	06	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

Johns Hephins Hospital

Screen Predrug

Study

TEST:NORMAL	$\frac{qqwwm\lambda\lambda}{N\mathcal{D}}$		2314.085 ddmmmyy		Date
NA:135-148 MEQ/L		146	142	140	97
K:3.5-5.0 MEQ/L		4.1	3.7	4.1	
CL:96-109 MEQ/L		109	106	108	
CO2:24-30 MEQ/L		25	24	24	
SUN:12-25 MG/DL		15	10	11	
CREAT:0.4-1.5 MG/DL		1.0	1.0	0.8	
GLU:70-115 MG/DL		82	160	72	
T.BILI:0.3-1.2MG/DL	otion of the	0.0	0.5	0.3	
D. BILI:0.1-0.4MG/DL	La July	MO.1	0.1	0.0	
CA:9.0-11.0 MG/DL		9.6	9.5	5.9	
PO4:3.0-4.5 MG/DL		4.4	2.7	3.5	
URIC A:4.2-8.8MG/DL		5.7	6.1	5.5	
T. PROT:6.0-8.5G/DL		7.1	6.8	6.8	
ALB.:3.2-5.3 G/DL		4.9	4.5	4.9	
AST:0-35 IU/L		15	13	15	i
ALT:0-30 IU/L		7	10	8	
ALK PHOS:0-95 IU/L		74	62	65	
CHOL:151-268 MG/DL		ND.	ND.	N.D.	
LDH:0-200 IU/L		118	107	N.D	
CPK:0-160 U/L(male)		51	49	N.D.	

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	R K	06	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Higher Hoghel

Screen Predrug

Study

N.D. 21Nov85 23 Nov85 26NN85 Date TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy Hgb 13.9-16.3 14 6 15.6 16.3 PCV 41.0-53.0 49.1 45.4 44.0 Plt 150-350 289 263 307 RBC 4.50-5.90 5.63 5.29 5.08 WBC 4500-11000 7300 5700 6900 2-6% Bands 7 6 9 Polys 31-76% 66 51 76 1-4% Eos 2 1 Bas 1 1 0 24-44% Lymphs 14 28 14 Atyp Lym 0 0 0 Monos 2-11% 5 11 3 Other 0 0 0 Retics 0.5-1.5% N.D 0.8 0.9

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	06	PROTOCOL DAMD 17-85-C-5133-02

URINALYSIS VALUES
Laboratory Christophins Hopkins Hopk

Screen Predrug

Study

TEST	NORMAL	ddmmmyy		250085 ddmmmyy		Date
Color/Ap			ND	20	ND	
Sp. Gr.			1.022	1.020	1.017	
рН			6.0	6.0	6.0	
Protein			Trace	Trace	Nex	
Ketones			Nex	ney	Ney	
Occ Bld			Nex	ner	Nin	
Bili.			New	×0.5	N.3	
RBC			0	0	0	
MBC			0	0	્	
Casts			ð	O	0	
Epi. Cel			0	0	9	
Crystals		-	0-4	O	0	
Bacteria			0	, 0	D	*

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
21 NN 85	/		
23 NW 85	V		
26NN85	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	P M L	06	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

N		ŀ
		ı
/	1	l
	\geq	X

			·	,		
1	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mww yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB.	Treatment
	•			□ Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T T	07	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	02JMV86	Screening laboratory
_	OBJANSO	History, Physical Exam
0	05JH486	Admission
2	06JAN86	P.O.
5	09 JAN 86	I.V.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of $-\sqrt{5}$ pages, for subject $+\sqrt{5}$.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

24,00,1,86 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	T/T FML	07	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

			03 TAN	HISTORY	1 + BH US	1
Date	of	evaluat:	ion 04/03/86	Examiner	Bungfollylle	1
Date	of	birth	At dd mmm yy At 01,57 Add mmm yy		Brent G Petty print name	
Age			28 _{yrs}			
Sex			<u>_</u> <u> </u>			
Race			<u>B</u>			

	No	Yes	Comments
Allergy	V		
Tobacco Use			
Alcohol Use			
Recreational Drug Use	1		
Medications past 2 weeks	V		
Experimental Drug Exposure		/	Tobamyan
Blood or plasma donor			
Prior Surgery			
Eye, ear, nose, throat	V		
Endocrine(diabetes,thyroid)			
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)			
Hepatitis, gastro-intestinal			
Genito-urinary		\checkmark	6 C Pard pills, varled
Musculoskeletal	/		
Neuropsychiatric	V		
Pesticide/herbicide use			Persule of home weaths ago. Sible trait
Other	1	/	Sible trait

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T T I	07	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION
Date 0/03/8()

dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
37.4 c	_ 7 <u>2</u> (min	20min		171.0	80.5

	GENERAL EXAMINATION:								
(check)	Nor.	Abn.	Provide details of abnormalities						
Head/Neck	/								
EENT		1	Wild selewin @ The few byuglind exerciseurs pot.						
Chest, lungs									
Heart									
Abdomen	0								
Genitalia	N	D							
Rectal	N	\mathcal{P}							
Extremities	4								
Skin 🖁		1	Scar Objection aut. dust Oforeaux, Went those.						
Neurologic			10						

CHEST X-RAY
Date Q3/3AN/S6

NORMAL	×	ABNORMAL	Describe	abnormalities:
			'	
L			 	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	TIT	07	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	234ACPO	0820	0850	IV	NA

Sympe + pyrile = 42.74206 g DOSAGE (total) 1.32 mg.

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B01, B47	0	09JAN86	0815	0815	B01:	B47: _{/0.06}
B02	0.08	٠,	0825	0825	BO2:8.51	
воз	0.16	ν,	0830	0830	B03:13.8	
B04, B48	0.25	٠,	0835	0835	B04:18,4	B48:8.20
B05	0.33	٠,	0840	0840	BO5:15,6	
B06	0.42	*1	0845	2845	BO6:16.3	
B07, B49	0.50	٠.	0850	0850	B07:17,9	B49: 7.12
B08	0.58	١.	0855	0855	BOB: 13.0	
B09	0.66	/1	0900	0900	B09: /0./	
B10, B50	0.75	٠,	0905	0905	B10:9,95	B50: 7.47
B11	0.83	• (0910	0910	B11:7.65	
B12	0.92	1.,	0915 0920AL	0915	B12:6.16	
B13, B51	1.0	11	0920 0940ex	0920	B13:6.34	B51: 7,79
B14, B52	1.33	l.	10940 1000pc	0940	B14:4.14	B52: 8,27
B15, B53	1.66	1.	1000 102021	1000	B15:2,86	B53: 8.55
B16, B54	2.0	4.5	1020	1020	B16:2,58	B54:8,86
B17	2.5	1,	1050	1050	B17:2,7/	

+ below assay sinsitivity

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FIL	0.7	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	09JAN86	0820.	0280	IV	MA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	09 JAN86	1120	1120	B18: 1.95	B55:9.23
B19	3.5	ε,	1150	1150	B19:2.15	
B20, B56	4.0	ħ	1220	1220	B20: *	B56:9.56
B21	5.0	١,	1320	1320	B21: *	
B22, B57	6.0	: i.,	1420	1420	BS5:*	B57: 9. 74
B23	7.0	b _k	1525	1520	B23: ≭	
B24	8.0	r,	1620	1 620	B24:*	
B25	10.0	٠.(1820	1820	B25:_ <u>*</u>	·
B26	12.0	٠,	2025	2020	B26: *	
B27, B58	24.0	584PCO1	0820	0820	B27:*	B58:/0.30
			8			<i>2</i>
* belo	wassa	y Bens	etivity 23	1		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FFT	07	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	8UAC PO	6 0820	0850	IV	NA

DOSAGE (total) 1,32 mg.

PHYSIOLOGIC VARIABLES

	N.B	: Handa	r.o met	er not we	ckine relie		P 1000.	on 09
	Sample No.	Scheduled Time (hours)		Scheduled	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
		0	OGJANE6	0815	0815	#72	#53/55	# ck
-		0.8	~	0825	0725		7	
		0.16	1,	0830	0830			
		0.25	c,	0835	0835	#75		
		0.33	1,	0845	0840			
		0.42		0845	0845			
		0.50	4	0850	0850	#80	#58/54	# 31
		0.58	1 <	0855	0855			٠
		0.66	. (0900	0900	_		
		0.75	£ ;	0905	0905	# 75		
		0.83	14	0910	0910			
		0.92	ι,	0915	0915			
		1.0	(,	0920	0920	#74	#59/56	
		1.33	ι,	0940	0940	#64	,	
		1.66	ι,	1000	1000	# 72		
		2.0	1.	1020	1020	# 90	#45/48	
		2.5	ł.	1050	1050			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	T i X	07	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	809JAN&	0890	0230	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	23NACPO	1120 -	1120	#78	٠.	
	3.5	()	1150	1150			
	4.0	`\	1220	1220	#60	# N.D	
	5.0	t.	1320	1320			
	6.0	61	1420	1420	#80	Q.11 #	
	7.0	11	1520	1520			
	8.0	i,	1620	1620			
	10.0	- 1	1820	1820			
	12.0	F.	2020	2020			
	24.0	10 JAN86	0820	0820	#70	# ND	# ok
	•						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	I I I	07	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	06 JANE6	0855	0855	PO	NA

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B28, B59	0	23 MAC 30	0810	0810	B28: _*	B59:10.17
B29, B60	0.25		0910	0910	B29: 🛨	B60:9.94
B30, B61	0.50	1,	0925	0925	взо:6,43	B61: 9.05
B31, B62	0.75	• •	0940	0940	B31:8.24	B62: 8.79
B32, B63	1.0		0955	0955	B32: 8.33	B63:8.48
B33, B64	1.33	4,	1015	1015	B33:8.69	B64:8,27
B34, B65	1.66	۱.	1035	1035	B34:12.3	B65: 7.46
B35, B66	2.0	1.4	1055	P1553 1055	вз5:9,72	B66: 7.22
B36	2.5	11	1125	1125	взє: 9.63	
B37, B67	3.0	C.	1155	1155	B37: 13.8	B67: 7,03
B38	3.5	1,	1225	1255	B38: /0.9	
B39, B68	4.0	41	1255	1255	вз9:8.48	B68: 7.53
B40	5.0	κ,	1355	1355	B40:5.08	
B41, B69	6.0	٠,	1455	1455	B41:2,69	B69:8,59
B42	7.0	١	1555	1555	B42:4,25	
B43	8.0	4,	1655	1655	B43:	B69A:9,02
B44	10.0	4 ,	1855	1855	B44:*	9.55

* below assay sensitivity 234

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T - T	07	PROTOCOL _ DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
02	06JAN86	0822	0855	PO	NA

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/M L	RBC ACHe uM/ml/mir
B45	12.0	cs Jange	2055	5022	B45: *	
B46, B70	24.0	BUALLO		2822	B46: *	B70: 9.94
-						
						191
	,					
+ lion	IL O AAI	11/2 DO2	setwity	235	<u> </u>	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	T · T	07	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
02	06 JAN 88	0855	0855	PO	NA

DOSAGE (total) 16.0 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	06JAN86	0810	0810	#64	#5//59	# ok
	0.25	٠.	0910		#60		
	0.50	4 }	0925	0925	#60	#43/51	# ok
	0.75	٠.	0940	0940	#58		
	1.0	4.6	0955	0955	#64	#46/59	
	1.33	10	1015	1015	#80		
	1.66	/t	1035	1035	#64		
	2.00	n	1055	1055	#72	#45/51	
	2.5	C _i · ·	1125	1125			
	3.0	1.	1155	1155	# 78		
	3.5	1,	1225	1225			
	4.0	L	1255	1255	# 80	#43/56	
	5.0	*,	1355	1355			
	6.0	£ į	. 1455	1455	# 78	#60/45	
	7.0	4.	1822	1555			
	8.0	* ,	1655	1655			
	10.0	٠.	1855	1855			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	J-I	07	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
02	OS JANSE	0855	0855	PO	NA

DOSAGE (total) 16.0 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	OGJANE	2055	2055	#N)		
	24.0	OPJANEG	2055	2055	470	#100	12 D#
		U					
		٠					
	·						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T T	07	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

CHEMISTRY VALUES
Laboratory Johns Highers Higher

Screen	Predrug	

Study

TEST:NORMAL	OS JUN 82	0274787	ddmmmyy	ddmmmyy	DSJAP86 ddmmmyy
NA:135-148 MEQ/L	142	141		144	
K:3.5-5.0 MEQ/L	3.9	4.3		4.2	
CL:96-109 MEQ/L	103	104		110	
CO2:24-30 MEQ/L	27	28		25	
SUN:12-25 MG/DL	10	12		9	
CREAT: 0.4-1.5 MG/DL	1.0	1. /		0.7	
GLU:70-115 MG/DL	77	81		83	
T.BILI:0.3-1.2MG/DL	2.5	1.2		1.0	
D. BILI:O. 1-O. 4MG/DL	0.2	0.1		0.1	
CA:9.0-11.0 MG/DL	9.8	9.8		8.8	
PO4:3.0-4.5 MG/DL	3.9	4.5		4.1	
URIC A:4.2-8.8MG/DL	6.8	6.5		ND	
T. PROT:6.0-8.5G/DL	7.2	7.1		6.4	
ALB.:3.2-5.3 G/DL	5.0	4.9		4.3	
AST:0-35 IU/L	2	14		13	
ALT:0-30 IU/L	18	19		25	
ALK PHOS:0-95 IU/L	46	44		40	
CHOL:151-268 MG/DL	еч	ND		181	
LDH:0-200 IU/L	ND	151	141	138	150
CPK:0-160 U/L(male)	20	218	155	123	

page 184 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T - T	07	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopkins Hopetal

Screen Predrug

Study

Date

TEST:NORMAL	ddmmmyy	NA_	ddmmmyy	qqwwwyy	ddmmmyy] [
NA:135-148 MEQ/L			141			
K:3.5-5.0 MEQ/L			4.1			
CL:96-109 MEQ/L			105			
CO2:24-30 MEQ/L			27			
SUN:12-25 MG/DL			13			
CREAT: 0.4-1.5 MG/DL			1.7			
GLU:70-115 MG/DL			80			
T.BILI:0.3-1.2MG/DL			0.9			
D. BILI:0.1-0.4MG/DL			0.1			
CA:9.0-11.0 MG/DL			9.3			
PO4:3.0-4.5 MG/DL			4.4			
URIC A:4.2-8.8MG/DL			5.8			
T. PROT:6.0-8.5G/DL			6.6			
ALB.:3.2-5.3 G/DL			4.5			
AST:0-35 IU/L			25			
ALT:0-30 IU/L			36			
ALK PHOS:0-95 IU/L			38			
CHOL:151-268 MG/DL			178			
LDH:0-200 IU/L			145			
CPK:0-160 U/L(male)			136			

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	<u> </u>	07	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOBY VALUES
Laboratory Johns Hopkus Hospital

Screen Predrug

Study

TEST	NORMAL	02 JAN86		OT JANSE ddmmmyy		Date
Hgb	13.9-16.3	14.9	14.5	13.7	13.]	
PCV	41.0-53.0	43.2	42.2	39.8	38.7	
Plt	150-350	227	224	197	223	
RBC	4.50-5.90	4.95	4.79	4.79	4.43	
MBC	4500-11000	5600	6200	4400	5600	
Bands	2-6%	8.	10	14	П	
Polys	31-76%	61	52	54	54	
Eos	1-4%	0	0	1	2	
Bas		1	0	6	0	
Lymphs	24-44%	27	26	23	30	
Atyp Lym		0	2	0	0	
Monos	2-11%	3	10	8	7	
Other		0	0	0	0	
Retics	0.5-1.5%	NP	1.4	1./	1.4	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T - T F M L	07	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

Clinical Pharmacology

Screen Predrug

Study

TEST	NORMAL	03JM 86 ddmmmyy	OS JAN86			qqwwwyy	Date
Color/Ap		Clear	milky	clear	Clear		
Sp. Gr.		1.019	1.021	1.024	1.023		
рН		5,0	5.0	6.0	6.0		
Protein		~2~	trea	~~01	res.		
Ketones		0	~es)	~ 6	~es		
Occ Bld		~24	0 ~98/	nei	res		
Bili.		may	nex	~ 26-	~~×		
RBC		()	0	0	0		
WBC		٥	0	ی	Q		
Casts		0	6	Ċ	0		
Epi. Cel		C	0	<u>ا</u>	0		
Crystals		0	0	٥	0		
Bacteria		0	0	0	D		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
03)AN86			
57 JAN 88			·
10.140.56	/	±.	and done of

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	T T T	07	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

ADVERSE EXPERIENCES

check if none occurred

г	·	⊀
ı	V	1
ı	\wedge	VΙ
١.	/	Y

ADVERSE	ONSET	CESSATION			ACTION
EVENT	(Date/Time)	Date/lime	(Check)	TEST DRUG	(Check)
#			☐ Mild	DEF.	None
	dd mmm yy	dd mmm yy	W110	PROB.	None
		40.0400			
	(0-2400)	(0-2400)	Mod	POSS. DEF. NOT	Treatment
				DEP. NOT	
			Sev	UNKNOWN	Stop test drug
#					
	dd mmm yy	dd mmm yy	Mild	DEF.	None
				PROB.	
	(0-2400)	(0-2400)	Mod	POSS.	Treatment
				DEF. NOT	
			Sev		Stop test
			Sev	Chilliowit	drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	OHW FML	08	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	20DEC85	Screening laboratory
_	DB MATEGO	History, Physical Exam
0	05JAN84	Admission
2	38 MAT 30	P.O.
5	28 MMC PO	IV.
		·

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 25.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

brew yoully M.D.

Investigator's signature

24/00/100

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	OHW FML	08	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	021	MAC	186
			dd	mmra	УУ

Examiner

Brent G. Pettn

Date of birth

10/0CT/50 dd mmm yy

Age

35 yrs

Sex

m

Race

w

	No	Yes	Comments
Allergy	/		
Tobacco Use	V		
Alcohol Use		V	4 bears link
Recreational Drug Use	/		
Medications past 2 weeks	~		
Experimental Drug Exposure		V	740
Blood or plasma donor	V		
Prior Surgery		V	1977 (D. Server Johnson 1978 (D. wist parties, regar
Eye, ear, nose, throat	1		
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	/		
Genito-urinary	/		
Musculoskeletal	/		
Neuropsychiatric	1		
Pesticide/herbicide use		~	Externator of home 10 mois
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	Q M W	08	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 22/ JAN 86 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height	(cm)	Wt.(kg)
3 7.5c	_ 6 4/min	2 ⊵/min	1 3	4/28_	17	8.0	75.9

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	/		
EENT		/	Septum > ®
Chest, lungs	~		
Heart	/		
Abdomen	1		
Genitalia			ND
Rectal			N.D.
Extremities	/		
Skin		~	Diffuse Signalsed prob. hyperglastic
Neurologic	V		·

CHEST X-RAY
Date O2/ <u>ごれい</u>を

NORMAL	×	ABNORMAL	Describe	abnormalities:
			•	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	Q H L	08	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	C9JAN8	0813	0843	IV	22

Symplet P = 42.54756 DOSAGE (total) 1.32 mg plasma CONCENTRATIONS

NB: pung indusing too quickly tel solo 15 minutes [PYRIDO] Sample Schedule Date Scheduled Actual RBC Time Clock Time Clock Time **ACHe** No. ddmmmyy NG/ML (0-2400)(0-2400) uM/ml/min (hours) B47: 13.16 BO1, B47 B01: * 09 JAN 86 2180 0812 B02:9,69 B02 0.08 0818 3180 воз: 20,9 B₀3 0.16 0823 0823 B04, B48 0.25 BO4: 18,9 B4B:10.31 1. 0828 0828 BO5:/4.5 B05 0.33 $t \in$ 0833 0833 BOE: 13,5 B06 0.42 1. 8180 0838 B07, B49 0.50 BO7: 11.7 B49: 9.45 0843 0843 BOB: 10,8 BOB 0.58 0848 0848 BO9: 8,15 **B**09 0.66 0853 0853 11 B10: 5,82 B50:10-28 B10, B50 0.75 0858 0858 B11: 5,26 B11 0.83 0903 0903 B12: 4,34 B12 0.92 0905 0908 B13: 3.51 B51: 10.85 B13, B51 1.0 0913 0913 B14, B52 1.33 B14:3,37 B52: 11,56 093] 0933 B15: 2.75 B53: 11.99 B15, B53 1.66 0953 0953 B16: * B16, B54 2.0 B54: 12.10 1013 1013 B17: 1.59 2.5 B17 1043

* below away sensitivity

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	0 <u>H</u> 3 F M L	08	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	OGJANZ	0813	0843	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/MC	RBC ACHe uM/ml/mir
B18, B55	3.0	09 JAN86	1113	1113	B18: *	B55: _{12.39}
B19	3.5	1.	1143	1143	B19: *	
B20 , B56	4.0	*1	1213	1213	B20:*	B56: 13.17
B21	5.0	A ₃	1313	1317	B21:*	
B22, B57	6.0	21	1413	1413	B22:*	B57: _{13.52}
B23	7.0	١, ,	1513	1513	B23: *	
B24	8.0	ls.	1613	1613	B24: *	
B25	10.0	17	1813	1813	B25: *	
B26	12.0	. 17	2013	2013	B26: *	
B27, B58	24.0	10JAN8	08,3	0813	B27: *	B58:13.62
			•			-
	7					
	w assa					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	C H W F M L	08	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	BUALPO	2 0813	0843	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample Scheduled Date Scheduled Actual Pulse Hand Coord	NB:	Hand gu	p nete	- stopped	water	cens	tale -	10000
0.16	Sample	Scheduled Time	Date	Scheduled Clock Time	Actual Clock Time	Rate	Grip	Coord Test
0.16		0	09JAN86	0812h	10812	# 84	#47/44	#ok
0.25		0.8	c,	0818	0818			
0.33		0.16	٠.	082]				
0.42 0.838 0.838 0.838 0.838 0.838 0.838 0.843 88 89 <th></th> <th>0.25</th> <th>l,</th> <th>0828</th> <th>0828</th> <th>#78</th> <th></th> <th></th>		0.25	l,	0828	0828	#78		
0.50		0.33	ž _g	0833	0833			
0.58 0.66		0.42	4 (0838	0838			
0.58 0.66		0.50	*1	0843	0843	* 88	#49/5	# ok
0.75 $\frac{1}{1}$ 0858 0858 $\frac{1}{82}$ 0.83 $\frac{1}{1}$ 0903 0903 0.92 $\frac{1}{1}$ 0913 0913 $\frac{1}{1}$ 70 $\frac{1}{1}$ 33 $\frac{1}{1}$ 093 093 $\frac{1}{1}$ 70 $\frac{1}{1}$ 34 $\frac{1}{1}$ 35 $\frac{1}{1}$ 36 $\frac{1}{1}$ 37 $\frac{1}{1}$ 37 $\frac{1}{1}$ 38 $\frac{1}{1}$ 3953 0953 $\frac{1}{1}$ 70 $\frac{1}{1}$ 3953 0953 $\frac{1}{1}$ 3953 $\frac{1}{1}$ 3953 $\frac{1}{1}$ 3953 $\frac{1}{1}$ 3953 $\frac{1}{1}$ 3954 $\frac{1}{1}$ 36 $\frac{1}{1}$ 37 $\frac{1}{1}$ 36 $\frac{1}{1}$ 37 $\frac{1}{1}$ 38 $\frac{1}{1}$ 38 $\frac{1}{1}$ 39 $\frac{1}{$		0.58	4.3	0848	0848			
0.83		0.66	42	0853	0853			
0.83		0.75	ч	0828	0858	#82		
1.0 " 0913 0913 # 70 #53/43 1.33 . 0933 0933 # 70 1.66 . 0953 0953 #70 2.0 1013 1013 #68 #31/36		0.83	4.	0903				
1.33		0.92	11	0908	0905			
1.33		1.0	j 4	0913	0913	# 70	#53/43	
1.66 0953 0953 #.70 2.0 1013 1013 #68 #31/36		1.33	. 1	. 0933	0933	# 70		
20		1.66	()	0953				
2.5 1043 1043 En		2.0		1013	1013	#68	#31/36	
		2.5		1043	1043		fr	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	Q H W F M L	08	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	09JAN8	. 0813	0843	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	09 JHN86	1113	1113	#74		
	3.5	6.5	1143	1143			
	4.0	- 1	1213	1213	#86	# 20	
	5.0	1.	1313	1313			
	6.0	* 55	1413	1413	#84	#ND	
	7.0	1.	1517	1513			
	8.0	t 4	1613	1613			
	10.0	*1	1813	1813			
	12.0	١,	2013	2013			
	24.0	BURTOI	0813		#84	#20	#ok
			-				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	OHL FML	08	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	06JAN88	0858	0858	PO	NA

synge + pyrids: 7.628749 DOSAGE (total) 16 mg.
synge 5 pyrids: 5.983889
PLASHA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B28, B59	0	OG JANES	0805	0805	B28:	B59: 13.63
B29, B60	0.25	•	0913	0913	B29: ¥	B60: 13.50
B30, B61	0.50	"(0928	0928	B30:2.2/	B61: /3.03
B31, B62	0.75	15	0943	0943	B31:6.44	B62:12.30
B32, B63	1.0	• •	0958	0958	B32:5,14	B63:12.09
B33, B64	1.33	ι,	1018	1018	B33:4.72	B64:12.27
B34, B65	1.66	1.	1038	1038	B34: 6.56	B65://.07
B35, B66	2.0	j e	1058	1058	B35: 8,03	B66: 10.95
B36	2.5	· /4	1128	1128	B36:7,15	
B37, B67	3.0	1,	1158	1158	B37:122	B67: 10.04
B38	3.5	"	1228	1228	взв: 9,30	
B39, B 68	4.0	٠,	1258	1258	вз9: 9,24	B68:/0.19
B40	5.0	١,	1358	1358	B40: 4.96	
B41, B69	6.0	6.	1458	1458	B41:3.22	B69: 12.12
B42	7.0	• (1558	1558	B42:2.24	
B43	8.0	4 (1658	1658	B43:1.80	B69A:/2.67
B44	10.0	11	. 1 8 5 8 Siturity 25	1858	B44: *	13.00

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	0 <u>4 2</u> F M L	08	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	OGJAN8	0858	0828	PO	NA

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/mir
B45	12.0	06 JAN86	2058	2058	B45: _★	
B46, B70			0858	0858	B46: *	B70: 13.63
				(9)		
	ii.					
				·		
					€,	
		, .				
* . lu	MUT ask	OAL ALD	witivity	251		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	OHW FML	○ 8	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	06JAN89	0858	0858	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	06.JAN8	- 0805		#88	#29/32	# _O L
	0.25	((0913		#90		1
	0.50		0928		*78	#41/45	# 02
	0.75	<i>(</i> ,	0943		# 78		
	1.0	/,	0958		#84	#37/45	
	1.33	4.	1018		#76	,	
	1.66	4	1038		#72		
	2.00	ć	1058		#64	#37/93	
	2.5	, ,	1388				
	3.0	۸	1058 en		# 76		
	3.5	//	1228				
	4.0	1.	1258		#76	#38/41	
	5.0	1.	1358				
	6.0	• (1458		# 80	#33/42	
	7.0	1.	1658 pm				
	8.0	1,	1758 gm				
	10.0	1,,	1858				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	는 M 기 기 기	08	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	06JAN8	08.78	0858	PO	NA

DOSAGE (total) _____ | 6 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	BUAC DO	2058	2058	#ND		
	24.0	07 JANSE	100	0858	#ND	# NO	ND#
	,		·				
	-						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u> </u>	08	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES, Laboratory Johns Hospital

	Screen	Predrug				Study
TEST:NORMAL	26Dec85		OBJAU86 ddmnmyy	SUAC FO		Date
NA:135-148 MEQ/L	145	145		144		
K:3.5-5.0 MEQ/L	5.0	4,9		4.7		
CL:96-109 MEQ/L	108	104		104		
CO2:24-30 MEQ/L	28	30		27		
SUN:12-25 MG/DL	19	10		14	•	
CREAT:0.4-1.5 MG/DL	0.9	1.0		0.9		
GLU:70-115 MG/DL	93	108		79		
T.BILI:0.3-1.2MG/DL	0.4	0,5		0,5		
D. BILI:0.1-0.4MG/DL	0.0	0.1		0.0		
CA:9.0-11.0 MG/DL	9.9	10.2		9.8		
PO4:3.0-4.5 MG/DL	ND	4.1		4.3		
URIC A:4.2-8.8MG/DL	6.7	5.9		5.7		
T. PROT:6.0-8.5G/DL	7.2	7.1		7.3		
ALB.:3.2-5.3 G/DL	4.8	5.0	,	5.0		
AST:0-35 IU/L	23	15		١٦		
ALT:0-30 IU/L	17	15		13		ji
ALK PHOS:0-95 IU/L	54	49		49		
CHOL:151-268 MG/DL	ND	ND		ND		
LDH:0-200 IU/L	ND	122	200	131	119	
	1					

66

49

ND

41

CPK:0-160 U/L(male)

page 12Aof 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>0</u> <u>H</u> <u>w</u> F M L	08	PROTOCOL _ DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

	6	D. a. d.			
		Predrug			
TEST:NORMAL	,	N A ddmmmyy	10JBN86	ddmmmyy	ddmmmyy
NA:135-148 MEQ/L			144		
K:3.5-5.0 MEQ/L			4.9		
CL:96-109 MEQ/L			105		
CO2:24-30 MEQ/L			29		
SUN:12-25 MG/DL			19		
CREAT:0.4-1.5 MG/DL			1.0		
GLU:70-115 MG/DL			85		
T.BILI:0.3-1.2MG/DL			0.3		
D. BILI:0.1-0.4MG/DL			0.0		
CA:9.0-11.0 MG/DL			9.8		
PD4:3.0-4.5 MG/DL			4.0		
URIC A:4.2-8.8MG/DL			6.6		
T. PROT:6.0-8.5G/DL			6.9		
ALB.:3.2-5.3 G/DL			21.8		
AST:0-35 IU/L			13 .		
ALT:0-30 IU/L			17		
ALK PHOS:0-95 IU/L			51		
CHOL:151-268 MG/DL			215		
LDH:0-200 IU/L			116		

56

CPK:0-160 U/L(male)

Study

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	0 <u>H</u> <u>w</u> F M L	08	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

Screen	Predrug		

		2100 00	OS JAN85	07 TO UST	10 JAN86		Date
TEST	NORMAL	qqwwwyy				ddmmmyy	Date
Ндь	13.9-16.3	15.0	15.4	15.0	13.8		ļ
PCV	41.0-53.0	43.1	44.1	42.1	40.1		
Plt	150-350	273	298	276	301		
RBC	4.50-5.90	4.64	4.74	4.66	4.32		
WBC	4500-11000	7600	9200	7000	7800		
Bands	2-6%	10	9	8	4		
Polys	31-76%	54	69	59	75-		
Eos	1-4%	4	1	2	2		
Bas		j	0	0	0		
Lymphs	24-44%	22	14	25	17		
Atyp Lym		6	1	0	0		
Monos	2-11%	3	6	6	2		
Other		0	0	0	0		
Retics	0.5-1.5%	20	1.0	1.0	1.0		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	P M L	08	PROTOCOL DAMD 17-85-C-5133-02

Laboratory John Hamacolo yy

Screen Predrug

Study

e

TEST	NORMAL	O2 JAN86	OSJAN86 ddmmmyy	077ANR ddmmmyy	DJAN25 ddmmmyy	ddmmmyy	Dat
Color/Ap		ND	yelln	-xellow	clear		
Sp. Gr.		1.019	1.021	1.023	1.020		
рH		6.5	6.0	6.5	7.0		
Protein		neg	Arece	nel)	trace		
Ketones		neer	760.	neu	ver.		
Occ Bld		nez	S meet	nag	7 es		
Bili.		new	ner	Len	3		
RBC		0	0	0	0		
MBC		0	0	0	0		
Casts		0	0	0	0		
Epi. Cel		0	0	0	0		
Crystals		0	Q	.0	0		
Bacteria		0	D	O	O		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
027AN86	1		
TRUATICO	V		
Janac OI	1		non-specific ST-T changes, questionally
	·		, , , , , , , , , , , , , , , , , , , ,
			257

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	2 <u>7</u> <u>3</u> F M L	08	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERBE EXPERIENCES

check if none occurred



	VERSE VENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	☐ Mod	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JRS FML	09	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
	26 Dec 85	Screening laboratory
_	26 Dec 85	History, Physical Exam
		Admission
2	DYJANOG	P.O. IV.
5	10 JAN 860	エV.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

__m.D. 24, Octi 86
ure dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JR S FM L	09	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	26, Dec 185	,
			dd mrara vv	

Examiner Bunt Petty M.D.

print name

Date of birth

05, JAN, 56 dd minm yy

Age

Sex

Race

	No	Yes	Comments
Allergy	/	,	
Tobacco Use		V	I pack per day
Alcohol Use		/	1 pack per day 2 cans beer / day on weekend Marijuana 1 cigaretti/minth
Recreational Drug Use		V	Marijuana I cigaretti/month
Medications past 2 weeks	V		
Experimental Drug Exposure		V	Pharmakinetico, FSK (apt 10/13/85
Blood or plasma donor	V		last 1983
Prior Surgery	V		
Eye, ear, nose, throat	V		
Endocrine(diabetes, thyroid)	V		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal	V		
Genito-urinary	V		
Musculoskeletal	V		
Neuropsychiatric	1		
Pesticide/herbicide use	/		No pets. No insected we.
Other	1		J

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	IRS FML	09	PROTOCOL DAMD 17-85-C-5133-02

PHYBICAL EXAMINATION Date 26 1 Dec 185

dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
37.2c	_ 68_/min	16/min	112178	182.00	76.5

			GENERAL EXAMINATION:
			GENERAL EXHITINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	/		
EENT		/	rew lymphad excrecences (R) portion pharyna, fair dentition, muld gingivitis, pigmentation of gums.
Chest, lungs	V		
Heart	/		
Abdomen	V		
Genitalia	~	D	
Rectal	Λ	D	
Extremities	V		•
Skin		V	Scalo Qupper arm, UQ, (R) lower back, forehead, Kneed from MVA 1979
Neurologic	/		ieft handed

CHEST X-RAY
Date 20/Nov/85

NORMAL Describe abnormalities:	
Small calcified granulomata in each apex	
Examiner Lent Methy Med Brent G. Petty, M.D.	
print name	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JR5 FML	09	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	10 JAN 86	0940	1005	ΙV	NA

Syringe + pyrido = 42.74 748 Syringe - pyrid= 22.655 Ho

- Infusion faster than scheduled
2° pump going too fast
Initially, then was slowed down.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
BO1, B47	O	10 JAN 86	0940	0940	B01: N.R.	B47: 13.35
802	0.08	11	0945	0945	B02:475	
B03	0.16	//	0950	0950	B03:8,73	
B04, B48	0.25	11	0955	0955	B04: 13.6	B48: 11, 48
B05	0.33	//	1000	1000	B05: 25.9	
B06	0.42	11	1005	1005	BOE:/8.3	
B07, B49	0.50	u	1010	1010	B07:203	B49:9,70
BOB	0.58	11	1015	1016	BOB: /4.0	
B09	0.66	. <i>II</i>	1020	1020	B09:9.24	
B10, B50	0.75	11	1025	1025	B10:9,19	B50: 9.53
B11	0.83	(1)	1030	1030	B11:629	
B12	0.92	l (1035	1035	B12: 2.63	
B13, B51	1.0	lt.	1040	1040	B13:627	B51: 10.34
B14, B52	1.33	H	1100	1100	B14:4.04	B52: 10.51
B15, B53	1.66	li .	1120	1/20	B15: 1.8/	B53: 11.05
B16, B54	2.0	11	1140	1140	B16: 2.38	B54: 11.55
B17	2.5	(1	1210	1210	B17:2.21	

N.R. - not run 262

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>JR5</u> FML	09	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	10 JAN 86	0940	1005	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	10 JRN 86	1240	1240	B18: 🙀	B55: 12.04
B19	3.5	11	1310	1310	B19: *	
B20, B56	4.0	lj	1340	1340	B20: *	B56:12.54
B21	5.0	li	1440	1440	B21: *	
B22, B57	6.0	0	1540	1540	B55: *	B57:13,18
B23	7.0	4	1640	1640	B23: *	
B24	8.0	11	1740	1740	B24: *	
B25	10.0	11	1940	1940	B25: *	
B26	12.0	Н	2140	2140	B26: *	
B27 , B58	24.0	11 JAN 86	0940	0940	B27: *	B58: 12.78
				7		
* bel	ow assa	y Deni	itivity 26	3		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JR5 FML	09	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

		Start drug ad istration (0				Bottle I.D. #
5	10 JAN 86	0940	100	5	IV	NIA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

NB Ele	retrove In	endering	En ston	accurate	on M	us da	λ ·
Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	10 JAN 86	0940	0940	# 40	#22/22	# OK
	0.8	· (1	0945	0945		LA	
	0.16	t _i	0950	0950			
	0.25	ti	0955	0955	#40		
	0.33	. ((1000	1000			
	0.42	(t	1005	1005			
	0.50	tı	1010	1010	#46	#24/21	# OK
	0.58	Į(1015	1016		en	
	0.66	lı	1020	1020			
	0.75	()	1025	1025	#60		
	0.83	Ц	1030	1030			
	0.92	h	1035	1035			
	1.0	11	1040	1040	#60	#21/22	
	1.33	tı	1100	1100	#68	xn	
	1.66	U	1120	1120	# 80		
	2.0	h	1/40	1140	# 78	# 23/24	-
	2.5	h	1210	1210		LR	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	J R 5 F M L	09	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	10 JAN 86	0940	1005	IV	NIA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	10 JAN 86	1240	1240	#72		
	3.5	· #	1310	1310			
	4.0	Ü	1340	1340	#80	#21/21	
	5.0	l/	1440	1440		3	
	6.0	11	1540	1540	# 84	23/25	
	7.0	11	1640	1640		建へ	
	8.0	11	1740	1740			
	10.0	li .	1940	1940			
	12.0	u	2140	2140			
	24.0	11JAN 86	0940	0940	#84	\$1/20	# OK
						La	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	J R 5 F M L	09	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	07 JAN 86	0840	0840	PO	NA

Syringe + pyrido= 7.63517 Syringe - pyrido= 5.95228

16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/m1/min
B28, B59	0	07 JAN 86	0811	0811	B28: *	B59: 12.76
B29, B60	0.25	t/	0855	0655	B29: *	B60: 12.38
B30, B61	0.50	t	0910	0910	взо: 227	B61: 12.03
B31,B62	0.75	t/	0925	0935	B31:3,95	B62: /1.5/
B32, B63	1.0	it	0940	0940	B32:4,0/	B63: _{11.35}
B33, B64	1.33	t _i	1000	1000	B33:5,46	B64: 11.02
B34, B65	1.66	li	1020	1020	B34: 6.01	B65: 10.91
B35, B66	2.0	li .	1040	1040	B35: 4.91	B66:10.59
B36	2.5	u.	1110	1110	B36: 4.76	
B37, B67	3.0	u	1140	1140	B37: 421	B67:10.35
B38	3.5	И	1210	1210	B38:5,00	
B39, B68	4.0	il.	1240	1240	B39: 6.19	B6B: 9.98
B40	5.0	11	1340	1340	B40:6.65	
B41, B69	6.0	μ	1440	1440	B41:2,//	B69:10.94
B42	7.0	(1	1540	1548	B42: 1,75	
B43	8.0	1,	1640	1640	B43: *	B69A: //.7/
B44	10.0	(ı	1840	1840	B44: *	11.78

* below assay sensitivity 266

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	J R 5 F M L	09	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	07JHN86	0840	0840	PO	WA

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B45	12.0	07 JAN 86	2040	2040	B45: ★	
B46, B70	24.0	08 JAN 86		0840	B46: *	B70: 12.63
			4			
					•	
		•				
			•			
* held	w asso	14 000	situity			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	JRJ FML	09	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	07 JAN 86	0840	0840	PO	NA

DOSAGE (total) 10mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	07 JAN 86	0802	0802	# 64	#55/54	# OK
	0.25	, H	0855	0855	#NotI		-
	0.50	. "	0910	0910	#60	#56/10	# OK
	0.75	11	0925	0925	# 64	,	
	1.0	/ (0940	0940	# 64	* 59/53	
	1.33	lı .	1000	1000	# 58		
	1.66	ħ	1020	1020	# 48		
	2.00	t ₁	1040	1040	#62	#49/49	
	2.5	11	1110	1110			
	3.0	i li	1140	1140	# 60		
	3.5	l)	1210	1210			
	4.0	11	1240	1240	#64	#50/59	
	5.0	H	1340	1340	2	,	
	6.0	U	1440	1440	#70	* 57/54	
	7.0	H	1540	1548			
	8.0	l:	1640	1640			
	10.0	ti .	1840	1840			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>I</u> <u>R</u> <u>5</u> F M L	09	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	07JAN 86	0840	0840	PO	NA

DOSAGE (total) 16 mg

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	09 JAN 86	2040	2040	#80		
	24.0	08JAN 86	2040 0840	2040 0840	# 46	#62/49	00
						,	
	-						
						+	
			•				
						4	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u> </u>	09	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES Laboratory Johns Hopkins Hospital

	Screen	Predrug	gam		448m	Study
TEST:NORMAL	<u>26 Dec 85</u> ddmmmyy	06 JAN 86 ddmmmyy	075AN 86	08 JAN 86 ddmmmyy	07 Jan Blo ddmmmyy	Date
NA:135-148 MEQ/L	143	141	138	139		
K:3.5-5.0 MEQ/L	4.1	4.7	4.3	4.6		
CL:96-109 MEQ/L	111	104	107	105		
CO2:24-30 MEQ/L	28	28	21	25		
SUN:12-25 MG/DL	18	13	14	18		
CREAT:0.4-1.5 MG/DL	1.2	1.2	1.4	1.3		
GLU:70-115 MG/DL	74	68	70	82		
T. BILI:0.3-1.2MG/DL	0.4	0.3	0.9	0.7		
D. BILI:0. 1-0. 4MG/DL	0.1	0.1	0.1	0.1		
CA:9.0-11.0 MG/DL	9.2	9.9	9.5	10.2		
PO4:3.0-4.5 MG/DL	ND	2.8	3.8	4.1		
URIC A:4.2-8.8MG/DL	5.1	4.5	5,1	5.4		
T. PROT:6.0-8.5G/DL	6.9	7.0	ND	ND		
ALB.:3.2-5.3 G/DL	4.1	4.3	4.5	4.5		
AST:0-35 IU/L	26	18	24	20		
ALT:0-30 IU/L	16	13	16	13		
ALK PHOS:0-95 IU/L	64	59	65	64		
CHOL:151-268 MG/DL	ND	ND	197	210		
LDH:0-200 IU/L	175	195	224	170		
CPK:0-160 U/L(male)	ND	¥385	341	270	325	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietmæn	JRS FML	09	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Chins Hopkins Duspital

Screen Predrug

Study

TEST:NORMAL	NA	10. TAN 86 ddmmmyy	11JHN80 ddmmmyy	ddmmmyy	Date
NA:135-148 MEQ/L		-	138		
K:3.5-5.0 MEQ/L			4.0		
CL:96-109 MEQ/L		22	105		
CO2:24-30 MEQ/L			24		
SUN:12-25 MG/DL			16		
CREAT: 0.4-1.5 MG/DL			1.3		
GLU:70-115 MG/DL			104		
T. BILI:0.3-1.2MG/DL			0.4		
D. BILI:0.1-0.4MG/DL			0.1		
CA:9.0-11.0 MG/DL			9,5		
PD4:3.0-4.5 MG/DL			3.5		
URIC A:4.2-8.8MG/DL			5.1		
T. PROT:6.0-8.5G/DL			7.4		
ALB.:3.2-5.3 G/DL			4.3		
AST:0-35 IU/L			27		
ALT:0-30 IU/L			26		
ALK PHOS:0-95 IU/L			65		
CHOL:151-268 MG/DL			205		
LDH:0-200 IU/L			168		
CPK:0-160 U/L(male)		257	2104		

INVESTIGATOR'S NAM	E PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	J R 5 F M L	09	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES Laboratory Johns Hopkins Hospital

		Screen	Predrug				Study
TEST	NORMAL	26Dec 85 ddmmmyy	OLEJANSO ddmmmyy	04 JAN 80 ddmmmyy	08 JAN 80 ddmmmyy		Date
Hgb	13.9-16.3	15,1	15.7	16.4	16.0	15.2	
PCV	41.0-53.0	44.2	44,9	48.7	47.7	45,1	
Plt	150-350	229	226	232	243	a <i>38</i>	
RBC	4.50-5.90	4.82	4.89	5.37	5.22	4.95	
WBC	4500-11000	7400	9400	6500	6900	2200	
Bands	2-6%	4	6	8	4	8 .	
Polys	31-76%	64	71	43	53	58	
Eos	1-4%	5	2	5	3	/	
Bas		0	0	/	/	/	
Lymphs	24-44%	17	15	32	26	24	
Atyp Lym		0	0	0	/	0	
Monos	2-11%	10	6	//	12	8	
Other		0	0	0	0	O	
Retics	0.5-1.5%	1.2	1.5	0,7	0,5	0.7	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JRS FML	09	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Johns Hopkins Hospital of Clinical Pharmacole yy

Screen Predrug

Study

TEST	NORMAL	clo Dec 85 ddmmmyy	DOJAN 86 ddmmmyy		10 JAN 86 ddmmmyy	ddmmmyy	Date
Color/Ap		ND	yellow	Cloudy	clear		
Sp. Gr.		1.018	1.033	1.021	1.021		
pН		0.5	6.5	6.0	6.0		
Protein		neg	- negr	neg	teace		
Ketones		neg	hea	reg	ula		
Occ Bld		neg	neg	near	neg		
Bili.		neg	neg	neg	near		
RBC		e	.Ø	\mathcal{E}^{I}	\mathcal{O}^{0}		
WBC		e	Ð	Ø	E		
Casts		Ø	Ø	D	Ø		
Epi. Cel		Ø	D	æ	Q		
Crystals		B	Ø	Ð	Ø		-
Bacteria		Ø	Ø	D	B		٠

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
02 JAN 86	/		
08 JAN 86	1		axis shifted rightward
11 JAN 86	/		axis Shifted leftward toward original

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JRS FML	09	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

4	
IV	
1 /	ΧI
V	

		1			1
ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mrnrn yy	dd mmm yy	Mild	DEF.	None
	(0-2400)	(0-2400)	□ Mod	PROB.	Treatment
			Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT # CO	DMPOUND Pyridostigmine
Lietman	W H M	610 PR	ROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
		Screening laboratory
-	MARTEO	History, Physical Exam
	06JAN 80	Admission
2	07.JAN 86	P.O.
5	28 CATEOI	IV

I certify that:

- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

24,001,000 dd/mmm/yy

INVESTIGATOR'S NAME	PT	INI	TIAL	PT #	COMPOUND	Pyridostigmine
Lietman	W F	H	n	010	PROTOCOL	<u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	02/JAN/86

dd mmm yy

Examiner

BRENT G. Petty M

Date of birth

27 /AUG 154 dd mmm yy

print name

Age

31 yrs

Sex

 \underline{m}

Race

B.

	No	Yes	Comments
Allergy	V		
Tobacco Use	V		
Alcohol Use	V		
Recreational Drug Use	V		
Medications past 2 weeks	V		
Experimental Drug Exposure		V	Pharmakinetics - last n 2mos ago
Blood or plasma donor	V		
Prior Surgery	V		
Eye, ear, nose, throat	V		
Endocrine(diabetes,thyroid)	V		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	1/		
Hepatitis, gastro-intestinal			
Genito-urinary	V		
Musculoskeletal	V		
Neuropsychiatric	V		
Pesticide/herbicide use		V	Dog washed for fleas in - month ago at his place residence but he did not do the washing. Extermination last done in 6 mos. ago.
Other	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	WH M	010	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 02/17/1/80 dd mmm yy

Temperature		7			_		
37.5 c	- 乾/mir	16/min	/	10,70	174	1.0	77.0

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	V		
EENT		V	Uvilla asymmetric and deviated to left
Chest, lungs	V		
Heart	/		50ft Short systolic flow murmur
Abdomen	/		
Genitalia	N	D	
Rectal		V.D	
Extremities	/		
Skin		/	Tatuo ® Forearm, scars ® anterolateral chest
Neurologic			

CHEST X-RAY Date 02/JAN / 86

NORMAL X ABNORMAL	Describe abnormalities:	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHAL	010 ·	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	10 JAN 86	0950	1:020	IV	NA

Syringe + pyrido = 42.44876 Syringe - pyrido = 22,43366

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B01, B47	0	10 JAN 86	0945	0945	B01:**	B47: 12.04
BOS	0.08	u	0955	0955	B02:2.27	
B03	0.16	11	1000	1000	BO3:5,23	
B04, B48	0.25	li.	1005	1005	B04: 3,94	B48: 10.39
B05	0.33	U	1010	1010	BO5:6.06	
B06	0.42	U	1015	10 15	BOE: 9.61	
B07, B49	0.50	l)	1020	1020	B07:12.4	B49: 8.93
B08	0.58	ll .	1025	1025	BOB:7,46	
B09	0.66	l(1030	1030	B09:5.32	
B10, B50	0.75	ų ·	1035	1035	B10:4,78	B50:9.37
B11	0.83	11	1040	1040	B11:3,68	
B12	0.92	Ц	1045	1045	B12:299	
B13, B51	1.0	U	1050	1050	B13: 2.8/	B51: 9.98
B14, B52	1.33	l)	1110	1110	B14: 1.62	B52: 10.09
B15, B53	1.66	Ц	1130	1130	B15: *	B53: 10.38
B16, B54	2.0	lı .	1150	1150	B16: *	B54: 10.84
B17	2.5	u	1220	1220	B17: *	

* below assay sensitivity

INVESTIGATOR'S NAME	PT	INI.	TIAL	PT #	COMPOUND	Pyridostigmine
Lietman	W F	H	th.	010	PROTOCOL	DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	10 JAN 86	0950	1020	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/MC	RBC ACHe uM/ml/min
B18, B55	3.0	10JAN 86	1250	1250	B18: *	B55:10.90
B19	3.5	(1	1320	1320	B19: *	
B20, B 5 6	4.0	11	1350	1350	B50: *	B56: //,57
B21	5.0	П	1450	1450	B21: 💥	
B22, B5 7	6.0	U.	1550	1550	B55: ¥	B57:/2,30
B23	7.0	Ц	1650	1650	B23: *	
B24	8.0	t _l	1750	1750	B24: ★	
B25	10.0	li	1950	1950	B25: 💥	
B26	12.0	ų .	2150	2150	B26: *	
B27, B58	24.0	11 JAN 86	0950	0950	B27: *	B 58: /2.52

* below assay sensitivity 279

INVESTIGATOR'S NAME	PT INITIE	L PT #	COMPOUND Pyridostigmine
Lietman	W H M	010	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	10.JAN 86	0950	1020	·IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

NB: EC	Rectanic de	and Sul	meter not a	wohing ace	gletary	on The	itab c
Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	10 JAN 86	0945	0945	#80	*22/20	# OK
	0.8					LA	
	0.16						
	0.25	11	1005	1005	# 84		
	0.33						
	0.42						
	0.50	11	1020	1020	# 78	# 22/21	# OK
	0.58		700			#22/21 In	
	0.66		,	70			
	0.75	11	1035	1035	# 80		
	0.83						
	0.92		9				
	1.0	11	1050	1050	# 72	* 21/23	
	1.33	"	11 10	1110	* 88	# 24/23 La	
	1.66	1,	1130	1130	#80		
	2.0	l I	1150	11 50	# 70	# 25/23	
	2.5	50 1				La	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHM	010	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	10 JAN 86	0 950	1020	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	10 JAN 86	1250	1250	*70		
	3.5	-					
	4.0	10 JAN86	1350	1350	*74	*33/25	
	5.0					St	
	6.0	10.JAN 86	1550	1550	#80	\$6/23	
	7.0					Lo	
	8.0		,				
	10.0						
	12.0						
	24.0 (4)	11 JAN 86	0950	0950	*60	*21/19	# OK
						da	
					·		
	·						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHM	010	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	07 JAN 86	0843	0843	PO	WA

Syringe + pipido = 7.62954 Syringe - pipido = 5.95975 DOSAGE (total) 16 mg.

PLASHA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B28, B59	0	07 JAN 86	0815	0815	B28: *	B59: //. 3/
B29, B60	0.25	li .	0858	0858	B29: *	B60: //.02
B30, B61	0.50	17	0913	0913	B30: ★	B61: 10. 70
B31,B62	0.75	l i	0928	0928	B31: 2.33	B62: 10.35
B32, B63	1.0	14	0943	0943	B32:2.66	B63: 10.31
B33, B64	1.33	l _l	1003	1003	взз:3.28	B64: 9, 75
B34, B65	1.66	П	1023	1023	B34: 3,56	B65: 9.35
B35, B66	2.0	1/	1043	1043	B35:4.57	B66: 8.88
B36	2.5	11	1113	1113	B36:454	
B37, B67	3.0	11	1143	1143	B37: 4.21	B67:8.49
B38	3.5	и	1213	1213	B38:662	
B39, B68	4.0	u	1243	1243	B39: 5,67	B68:8.83
B40	5.0	11	1343	1343	B40:576	
B41,B69	6.0	(e	1443	1443	B41:2,01	B69: _{10.08}
B42	7.0	1,	1543	1543	B42: 2.21	
B43	8.0	H	1643	1643	B43: ★	B69A: 10.68
B44	10.0	()	1843	1843	B44: *	10.86

* below assay sensitivity 282

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	WHML	DIO	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	07 JAN 86	0843	0843	PO	NIA

DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/MC	RBC ACHe uM/ml/min	
B45	12.0	07 JAN 86	2043	2043	B45:*		
B46, B70	24.0	08JHN86	0843	0905	B46: *	B70://./3	
F							
		·					
							
			·		-		
* 10	Dour ax	0004	nemostició	41			
* below appay pensitivity							

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHM FML	010	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	07 JAN 86	0843	0843	PO	NIA
			1	/	1

DOSAGE (total) 16 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	075AN86	0808	0808	#76	#51/55	#OK
	0.25	11	0858	0858	*74		
	0.50	1/	0915	0915	#68	\$5/48	# OL
	0.75 ⁻	17	0928	0928	#78		
	1.0	4	0943	0943	*64	#51/48	
	1.33	(/	1003	1003	# 80		
	1.66	<i>(</i> /	1023	1023	#60		
	2.00	11	1043	1043	#64	#57/58	
	2.5						
	3.0	//	1143	1143	#62		
	3.5	4					
	4.0	11	1243	1243	# 72	#54 <i> </i> 54	
	5.0						
	6.0	(1	1443	1443	#72	* 53/54	
	7.0						
	8.0						
	10.0						

INVESTIGATOR'S NAME	PT INITIAL PT	COMPOUND Pyridostigmine
Lietman	W H M DIO	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	045AN86	0843	0843	PO	NA

DOSAGE (total) 16 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	OFJAN80	2043		#ND		
	24.0	08JHN 86	2043 0843	0905	#60	#53/54	OK#
						/	
	_						
				,			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WH ML	010	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VACUES Hopkins Haspital

Screen	Predrug

	Screen	Preurug				Study
TEST:NORMAL	D2JAN86 ddmmmyy	DO JANSO domininyy		07JAN86 ddmmmyy		Date
NA:135-148 MEQ/L	142	138	136		138	
K:3.5-5.0 MEQ/L	4.7	4.5	5.0		5.1	
CL:96-109 MEQ/L	104	104	99		104	
CD2:24-30 MEQ/L	30	29	23		25	
SUN:12-25 MG/DL	10	14	9		14	
CREAT: 0.4-1.5 MG/DL	1.0	1.0	1.0		1.1	
GLU:70-115 MG/DL	78	71	75		81	
T.BILI:0.3-1.2MG/DL	0.8	0.3	0.7		0.6	
D. BILI: 0. 1-0. 4MG/DL	0.1	0.0	0.1		0.1	
CA:9.0-11.0 MG/DL	10.0	10.0	9.8		10.4	
P04:3.0-4.5 MG/DL	3.6	3.0	3.8		4.1	
URIC A:4.2-8.8MG/DL	5.6	5.5	5.5	,	5,5	
T. PROT:6.0-8.5G/DL	7.6	7.3	7.4		ND	
ALB.:3.2-5.3 G/DL	5.1	4.9	4.8		5.1	
AST:0-35 IU/L	29	20	24		22	
ALT:0-30 IU/L	35	28	27		27	
ALK PHOS:0-95 IU/L	40	36	33		39	y.
CHOL:151-268 MG/DL	ND	ND	211		236	
LDH:0-200 IU/L	. ND	137	173	139	150	
CPK:0-160 U/L(male)	ND	147	122	110	94	

page 12Aof 15

Study

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H M	810	PROTOCOLDAMD_17-85-C-5133-02

Screen Predrug

CHEMISTRY VALUES Hipking Haspital

35

220

TEST:NORMAL	0	A	N	iuuaaa T	10JANSU_		ddmmmyy	Date
NA:135-148 MEQ/L		1				139		
K:3.5-5.0 MEQ/L						4,5		
CL:96-109 MEQ/L						105		
CO2:24-30 MEQ/L						25		
SUN:12-25 MG/DL						20		
CREAT: 0.4-1.5 MG/DL						0.9		
GLU:70-115 MG/DL						77		
T.BILI:0.3-1.2MG/DL						0.5		
D. BILI:0.1-0.4MG/DL						0.0		
CA:9.0-11.0 MG/DL						9.6		
P04:3.0-4.5 MG/DL	T					4.1		
URIC A:4.2-8.8MG/DL	T					4.9		
T. PROT:6.0-8.5G/DL						7.5		
ALR.:3.2-5.3 G/DL						4.8		
AST:0-35 IU/L		*				44		
ALT:0-30 IU/L						31		

ALK PHOS:0-95 IU/L

CHOL:151-268 MG/DL

CPK:0-160 U/L(male)

LDH:0-200 IU/L

INVESTIGATOR'S NAME	PT INITIAL PT #	COMPOUND Pyridostigmine
Lietman	W It M DIO	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hopkins Hospital

Screen	Predrug				Study
02 JAN 86	06JAN86	07 JAN 86	08 JAN86	11 JAN86	Date

TEST	NORMAL	DA TAN 86 ddmmmyy	06JHN86 ddmmmyy	07 JAN 86 ddmmmyy	08 JAN 80	
НдЬ	13.9-16.3	15.2	14.2	14.6	15.1	13.4
PCV	41.0-53.0	44.5	40.5	43,5	45.5	39.9
Plt	150-350	224	215	208	241	261
RBC	4.50-5.90	5,49	4.88	5.35	5.54	4.89
MBC	4500-11000	4900	6700	6300	6100	7700
Bands	2-6%	/	5	9	4	16
Polys	31-76%	60	66	47	60	45
Eos	1-4%	0	0	4	0	1
Bas		0	/	0	0	2
Lymphs	24-44%	30	21	27	25	25
Atyp Lym		C	/	0	0	0
Monos	2-11%	9	6	13	//	10
Other		0	0	0	0	0
Retics	0.5-1.5%	ND	ND	1.0	0.8	1.2

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHA	Ď10	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

Laboratory Christophens Vniverily Clinical Pharmacology

Screen Predrug

Study

TEST	NORMAL:	COJAN86 ddmmmyy		OB JAN86 ddmmmyy		ddmmmyy	Date
Color/Ap		yellow	uellow	Clear	clear		
Sp. Gr.		1.023	1.017	1.018	1.019		
рН		6.5	6.0	6.0	5.0		
Protein		neg	nea	neg	trace		
Ketones		neg	nea	nea	neg		
Occ Bld		neg	nea	neg	neg		
Bili.		neg	noa	nea	rea		
RBC		0	Ø	e	Ø		
WBC		rare	Ø	Ø	Ø		
Casts		0	Ð	Ø	Q		
Epi. Cel		0	Ø	Ø	a		
Crystals		D	Ø	Ø	D		
Bacteria		D	Ø	0	Ø		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
02 JAN 86	1		
02 JAN86 08 JAN86 11 JAN86	J		
11 JAN86	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WKM	010	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

1	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	Ust a dodd	10 Jan 86 dd mmm yy		Mild	DEF.	None
1	Hot and cold Hadie, initable	1430 (0-2400)	<u>1500</u> (0-2400)	Mod	PROB.	 Treatment
		•		Sev	DEF. NOT	Stop test
#			dd mmm vv	☐ Mild	DEF.	None
		(0-2400)	dd mmm yy (0-2400)	Mod Mod	PROB. POSS.	Treatment
				□ Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R A H F M L	011	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	DOWNER	Screening laboratory
_	28 CHATPO	History, Physical Exam
0	98/HEE61	Admission
2	13JAN81	I.V.
5	162AN 86	P.O.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 16.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigatoff's signature

24,01,06 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H	011	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	09,	MAK	86
			dd	mtata	УУ

Examiner

Brent & Potty

Date of birth

13/Aug 57
dd mmm yy

Age

28 yrs

Sex

 $-\infty$

Race

B

any different con			
	No	Yes	Comments
Allergy			
Tobacco Use		/	1/2 - 1 ppd
Alcohol Use		~	2 cans/day on weekens
Recreational Drug Use		/	Marijaure 93 wks
Medications past 2 weeks	~		0
Experimental Drug Exposure		/	Pharmakinetics 2 yrs go
Blood or plasma donor			Dest 00 1985
Prior Surgery		V	Appy 1978, tonoclastony
Eye, ear, nose, throat	~		
Endocrine(diabetes, thyroid)	1		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	V		
Genito-urinary		~	5-364.115 1977 6.C. 1979- Rx PCN
Musculoskeletal	/		¥
Neuropsychiatric	/		
Pesticide/herbicide use			
Other	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date ON JAN ST dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height	(CM)	Wt. (kg)
37.40	_ & D/min	16/min		0120	_1 ?	9.0	77.1

	GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities					
Head/Neck	/							
EENT	/		Residual tousillar tissue on @					
Chest, lungs	/							
Heart	/							
Abdomen	/							
Genitalia			N.D.					
Rectal			N.D.					
Extremities	/							
Skin								
Neurologic	/							

CHEST X-RAY Date 09/3/N/86

NORMAL	×	ABNORMAL	Describe	abnormalities:
	,			

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	R L H F M L	(j) (PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	13 JAN86	1015	1045	IV	NA

Syringe + gyrido = 42.356948 DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
BO1, B47	0	13 JAN 86	NA.	08:14 0826gn	B01: *	B47:15.08
B02	0.08	٠,	1050	1020	B02:5.11	
B03	0.16	c _k	1025	1025	B03: N.R.	
B04, B48	0.25	i,	1035	1030	B04:3.37	B4B: 13,19
B05	0.33	C ₄	1035	1035	B05: 19,2	
B06	0.42	4,	1040	1040	воб: 23.3	
B07, B49	0.50	Ĺq.	1045	1045	B07: 27.8	B49: 10.76
BOB	0.58	1,	1050	1020	BOB:25,7	
B09	0.66		1055	1055	B09:195	
B10, B50	0.75	h	1100	1100	B10:13.9	B50: 10.79
B11	0.83	м	1105	1105	B11:12,7	
B12	0.92	4	1110	1110	B12:9.38	
B13, B51	1.0	h	1115	1115	B13:8,65	B51: //. 36
B14, B52	1.33	þ.	1135	1135	B14:6,10	B52: //.85
B15, B53	1.66	7,	1155	1155	B15:6,07	B53: 12.99
B16, B54	2.0	fs.	1215	1215	B16:4,07	B54: /2.99
B17	2.5	1.	1245	1245	B17: 3.3j	

* thelow assay sinsettirty 294

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	Du	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	13 JAN86	1015	1045	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	13JAN82	1315	1315	B18:3,86	B55: 14.12
B19	3.5	ν,	1345	1345	B19:3,23	
B20, B56	4.0	٠,	1415	1415	B20:2,70	B56:14.75
B21	5.0	I,	1515	1515	B21: 2,0/	
B22, B57	6.0	٠,	1615	1615	B55: *	B57:14,76
B23	7.0	t,	1715	1715	B23: 👍	
B24	8.0		1815	1815	B24: ¥	
B25	10.0	l,	2015	2015	B25: *	
B26	12.0	- N	2215	2215	B26: *	
B27, B58	24.0	٠,	1015	1015	B27: 💥	B58:/5,48
			·			
			10:4: 1-4:			

- below assay sensitivity 295

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	13 JAN &	1015	1045	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

1 Nec	hanical. h	and sixp	moter used	- Bolieved	to be	accu-	. ta
Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	13 JAN82	N.A.	0826	#64	#43/ss	# o/c
	0.8	* · · · ·	1020	1020			
	0.16	r,	1025	1025			
	0.25	4	1030	1030	#68		
	0.33	11	1035	1035			
	0.42	4,	1040	1040			
	0.50	٠,	1045	1045	#70	#43/56	# o Ł
	0.58	٠.	1050	1050			
	0.66	4.	1055	1055			
	0.75	٠,	1100	1100	# 68		
	0.83	- 4	1105	1103			
	0.92	• .	1110	1110			
	1.0	1.	1115	1115	*68	#43/50	
	1.33	ı,	. 1135	1135	#75		
	1.66	1	1155	1155	#66		
	2.0	•,	1215	1215	#70	# 39/52	
	2.5	4	1245	1245			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietmæn	R H F M L	011	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	13 JAN86	1015	1045	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	13 JAN86	1315	1315	#70		
	3.5	٠, ,	1345	1345			
	4.0	1,	1415	1415	#64	#44/48	
	5.0	١,	1515	1515			
	6.0	• • 1	1615	1615	#68	# 48/48	
	7.0		1715	1715			
	8.0	15	1815	1815			
	10.0	٧,	2015	2015			
	12.0	13	2215	2215			
	24.0	JB CHECHI	1015	1015	#60	#54/44	#ok
				Э			
			1				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	DII	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	163A N&	0820	0890	PO	NA

Dyrige & pyride = 7.61223 of DOSAGE (total) 16 mg.

Syrige & pyride = 5.94095

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min			
B28, B59	0	16 JAN86	N.A.	0804	B58:*	B59:14.81			
B29, B60	0.25	• •	0835	0835	B29: 5.11	B60: 14.99			
B30, B61	0.50	13	0850	0850	B30: 12.7	B61: 12.99			
B31,B62	0.75	V.	0905	0905	B31:17,2	B62: //.5/			
B32, B63	1.0	٠,	0920	0920	B32: 19.3	B63: 10.29			
B33, B64	1.33	× .	0940	0940	B33: 23,7	B64: 9.76			
B34, B65	1.66	۲,	1000	1000	B34:19,9	B65: 974			
B35, B66	2.0	• <	1030	1020	B35:16.3	BE6:10.55			
B36	2.5	*\	1050	1050	B36: 14.6				
B37, B67	3.0	41	1120	1120	B37:14,8	B67: 10.58			
B38	3.5	l,	1150	1150	B3B:01,84				
B39, B68	4.0	4	1220	1223	B39:11,9	B68:11.17			
B40	5.0	bi,	1320	1320	B40: 9,/8				
B41, B69	6.0	٠,	1420	1425	B41:7.99	B69:12.24			
B42	7.0	٧ -	1520	1520	B42: 6.88				
B43	8.0	l.	1620	1620	B43: 5.08	B69A:/3,39			
B44	10.0	44	1820	1822	B44: 2.04	14.07			
* Jule	* Iulaur assay sensitivity 298								

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: DRAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	16JAN88	೧೯೨೦	0890	PO	NA

DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	RE/ML	RBC ACHe uM/ml/mir
B45	12.0	16Jnn86	2020	2020	B45:1	
B46, B70	24.0	17 JAN86	_	0810	B46: 🔆	B70: 14.99
					•.	
-						
	*					
						_
0 0					7	
* belo	wason	ay se	noitivity			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	R L H F M L	O) T	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
S	16 JAN86	0820	0820	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	16 JA 1186	NA	0810	#84	# ND	# ok
	0.25	,	0835	0 839	# 78		_
	0.50		0850	0850	# 80	# ND	# ot
	0.75	• (0905	0905	# 78		
	1.0	44	0920	0920	# 72	4 ~ 4	
	1.33	٠,	0940	0940	# 76		
	1.66	٠,	1000	1000	#80		
	2.00	٠,	1020			# ND	
	2.5	£ q	1050	1050			
	3.0	c _t	1120	1120	#64		
	3.5	1,	.1150	1150			
	4. 0	٠	1220	1220	# 84	# ~ D	
	5.0	I.	1320	1320			
	6.0	٠.	1420	1420	#88	#~'9	
	7.0	1,	15 22	15 Qui			
	8.0		1620	1620			
	10.0	۲,	1820	1820			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	16JAN86	0832	2Q 2D	PO	Na

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	38 UAC 31	20 20	مدمد	# ND		
	24.0	17 JAN86	i	0820	# ND	# N D	ND#
	6						
			Y				
	© .						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopkin Hopkin Hopkin

Screen Predrug

	screen	Preurug				Study
TEST:NORMAL	OZJAN86		13JANS6 ddmmmyy		16 Jang	Date
NA:135-148 MEQ/L	143	141	143	141		
K:3.5-5.0 MEQ/L	4.3	4.4	4.2	4.1		
CL:96-109 MEQ/L	109	107	107	104		
CO2:24-30 MEQ/L	23	24	19	26		
SUN:12-25 MG/DL	12	13	12	15	•	
CREAT: 0. 4-1.5 MG/DL	1.1	1.3	1. 2	1.4		
GLU:70-115 MG/DL	74	153	63	65		
T.BILI:0.3-1.2MG/DL	0.8	0.4	0.7	ND		
D. BILI:0.1-0.4MG/DL	0.1	0.0	0.1	0.1		
CA:9.0-11.0 MG/DL	9.3	8.9	9.6	9.9		
PO4:3.0-4.5 MG/DL	4.0	4.3	3.9	3.8		
URIC A:4.2-8.8MG/DL	4.7	4.9	51/	5.3		
T. PROT:6.0-8.5G/DL	6.4	6.1	6.6	6.8		
ALB.:3.2-5.3 G/DL	4.2	3.8	4.1	4.3		
AST:0-35 IU/L	31	34	24	32		
ALT:0-30 IU/L	23	25	25	27		
ALK PHOS:0-95 IU/L	45	41	52	46		
CHOL:151-268 MG/DL	ND	160	189	188		
LDH:0-200 IU/L	ND	126	ND	129	136	
CPK:0-160 U/L(male)	46	644	401	3 59	339	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	Dil	PROTOCOL DAMD 17-85-C-5133-02

CHEMIBTRY VALUEB
Laboratory Johns Hopkin Hospital

N.A. N.A Screen Pregrug

	12	JAP			17 JAN86			Date
TEST: NORMAL	dg	DAMBAA	ddm	nunyy	ddmmmyy	ddmmmyy	ddmmmyy	
NA:135-148 MEQ/L					144			
K:3.5-5.0 MEQ/L					4.4			
CL:96-109 MEQ/L					106			
CO2:24-30 MEQ/L					26			
SUN:12-25 MG/DL					ND			
CREAT: 0.4-1.5 MG/DL					1.2			
GLU:70-115 MG/DL					78			
T.BILI:0.3-1.2MG/DL					0.9			
D. BILI:0.1-0.4MG/DL					0.1			
CA:9.0-11.0 MG/DL					9.8			
P04:3.0-4.5 MG/DL					4.1			
URIC A:4.2-8.8MG/DL					5.3			
T. PROT:6.0-8.5G/DL					7.1			51
ALB.:3.2-5.3 G/DL					4.4			
AST:0-35 IU/L					3 2			
ALT:0-30 IU/L					22			
ALK PHOS:0-95 IU/L					46			
CHOL:151-268 MG/DL					180			
LDH:0-200 IU/L					132			
CPK:0-160 U/L(male)			1		3.20			

INVESTIGATOR'S	NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman		R L H F M L	011	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hopkins Hospital

Screen Predrug

TEST	NORMAL	02]AN&	12JAN86 ddmmmyy	14 JAN& ddmmmyy	qquumyy 1JJVAU86	qqwwmyy	Date
НдЬ	13.9-16.3	14.7	14.5	15.8	ND		
PCV	41.0-53.0	41.8	42.7	46.7	ND		
Plt	150-350	243	254	265	20		
RBC	4.50-5.90	4.34	4.47	4.88	ND		
WBC	4500-11000	7900	7900	7300	ND		
Bands	2-6%	5	8	4	ND		
Polys	31-76%	42	42	40	7		
Eos	1-4%	0	0	0	2		
Bas		1	0	1	29		
Lymphs	24-44%	44	46	37	20		
Atyp Lym		2	0	17	ND		
Monos	2-11%	6	4	7	\sim 0	(*)	
Other		0	0	0	ND		
Retics	0.5-1.5%	ND	0.8	N.D	ND		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L M F M L	011	PROTOCOL DAMD 17-85-C-5133-02

URINALYBIB VALUEB
Laboratory Johns Hopkins Hopkins Hopkins

Screen Predrug

Study

TEST	NORMAL	CG JAN86		284ACP1		Date
Color/Ap		yellow	zellow	rellas	milka	
Sp. Gr.		1.019	1.017	1.023	1.019	}
рH		7.0	6.0	7.5	7.0	
Protein		trace	drace_	2+	there.	
Ketones		near	nal	neg	208	
Occ Bld		rea	معا	ney	~0	
Bili.		neev	~ 37	~~X		
RBC		0	0	2	0	
MBC		0	0	0	0	
Casts		0	0	0	2	
Epi. Cel		0	0	0	0	
Crystals		O	O	O	0	
Bacteria		ט	O	U	0	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
09JAN8	V		
14JAN86	V		
17 JAN86	V		
	,		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R L H F M L	011	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred



	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	Mild	DEF.	□ None
		(0-2400)	(0-2400)	□ Mod	PROB.	
				 Sev	DEF. NOT	Stop test
#			dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB.	 Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AWB	012	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
		Screening laboratory
	09 JAN 86	History, Physical Exam
0	JBUAT 61	Admission
2	13JAN86	IV
5	16JAN86	Po
		.*

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject # 12.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

______M.D.

Investigator's signature

24/04/186 dd/mmm/yy

INVESTIGATOR'S NAME	PT	INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	₽	WB M L	012	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	09/	JAH	86
			dd	mmm	УУ

Date of birth

27/JUN 56

Age

29 yrs

Sex

 \underline{m}

Race

B

	No	Yes	Comments
Allergy			
Tobacco Use		/	1/2882
Alcohol Use		/	; et ein lute. 2 6 pades/whend Occ. mangaum - ent - 3 du ago, auroge - 3/wh
Recreational Drug Use		1	Occ. mangama - lest - 3 de
Medications past 2 weeks	/		
Experimental Drug Exposure		/	Tobramyen - here Pharme - Kincties
Blood or plasma donor		/	Last ~ 4 months ago
Prior Surgery			Broke foot, but some bone
Eye, ear, nose, throat			alesses x 8yrs.
Endocrine(diabetes, thyroid)			
C-V (heart murmur, HBP)			
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal			
Genito-urinary	/		
Musculoskeletal	/		
Neuropsychiatric			
Pesticide/herbicide use		V	Last just before Chustmas and then 2 days ago.
Other	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	A W IS F M L	012	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION
Date 09/ 10/86
dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height (cm)	Wt.(kg)
37.40	_68/min	2 @/min	_12	0164	182.0	79.2

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	/		
EENT	_		
Chest, lungs	/		
Heart	~		
Abdomen	/		
Genitalia			N.D.
Rectal			N.D.
Extremities	/		
Skin		/	ocens both musts, (B) upper arm, (D) freem
Neurologic	/		

CHEST X-RAY
Date ON JAN 86

NORMAL	ABNORMAL	Describe	abnormalities:

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	A W B F M L	012	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	BUAL EI	1018	1048	IV	NA

ognife + pyrids = 42, 55140 g DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B01, B47	0	13 JAN8	2 %	0810	B01: *	B47:15.28
B02	0.08	٠.	1023	1023	B02:3.37	
B03	0.16		1028	1028	B03:10.4	
B04, B48	0.25	٠, (1033	1033	B04:20.3	B48: 13.37
B05	0.33	ι,	1038	1078	B05:30.9	
B06	0.42	1,	1043	1043	BOE: 28,1	
B07, B49	0.50	• 4	1048	1048	B07: 22,1	B49:10.37
BOB	0.58	•	1053	1053	BOB:16-3	
B09	0.66	· •	1058	1058	809:12,8	
B10, B50	0.75	·.	1103	1103	B10:173	B50:10.84
B11	0.83	ās.	1108	1108	B11:8.30	
B12	0.92	٤,	11/3	1113	B12:8,36	
B13, B51	1.0	٠,	1118	1118	B13: 9.06	B51://.55
B14, B52	1.33	۲,	1138	// 38	B14:617	B52: /1.68
B15, B53	1.66	٠.,	1158	1158	B15: 4,66	B53: 13.61
B16, B54	2.0	11	1218	1218	B16:3,3/	B54: 13,37
B17	2.5	6.	1248	1248	B17: 2.53	
* lu	elow c	issay	Sensitivi	ty 310		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A WB F M L	012	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	13 JAN86	1018	1048	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B18, B55	3.0	13 JAN86	1318	1318	B18:1,74	B55:13.67
B19	3.5	١,	1348	1348	B19: 1.57	
B20, B56	4.0		1418	1418	B20: *	B56: 14,51
B21	5.0	٠.	1518	1518	B21: *	
B22, B57	6.0	4	1618	1618	B55: *	B57: 14.84
B23	7.0	ω,	1718	1718	B23: *	
B24	8.0	٠,	1818	1818	B24: *	
B25	10.0	٠,	2018	2018	B25: *	
B26	12.0		2218	2218	B26: *	
B27 , B 5 8	24.0	14 JAN86		1018	B27: *	B58: 15.80
* belo	w assi	ay Sens	itivity	311		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AWB FML	012	PROTOCOL <u>DAMD_17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	13 JAN86	1018	1048	IV	NA

DOSAGE (total) 1.32mg.

PHYSIOLOGIC VARIABLES

Sample No.		I	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	13 JAN88	NA	0827	* 52	#46/45	#0/c
	0.8		1023	1027			
	0.16	٠,	1028	1028			
	0.25	kog	1033	1033	# 60		
	0.33	4.1	1038	1038			
	0.42	٤,	1043	1043			
	0.50	• (1048	1048	#64	#53/56	#ok
	0.58	t ₃	1053	1053			
	0.66	٤,	1058	1058			
	0.75	4	1103	1103	# 64		
	0.83	~	1108	1108			
	0.92	۱.	1113	11/3			
	1.0	4-	1/18	1/18	# 64	#49/58	
	1.33	٠,	1138	1138	#60		
	1.66	۲,	1158	1158	#52		
	2.0	£ <	1218	1218	#60	#51/53	
	2.5	**	1248	1248			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	A W L	012	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	13 JAN 86	1018	1048	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	13 JAN86	1318	1318	#76		
	3.5	.,	1348	1348			
	4.0	٠,	1418	1418	# 60	#48/56	
	5.0	• ,	1518	1518			
	6.0	4.1	1618	1618	* 72	#45/48	
	7.0	• 1	1718	1718			
	8.0	٠.	1818	1818			
	10.0	17	2018	2018			
	12.0	11	2218	2218			
	24.0	147AN86	1018	1018	(N#	#ND	# N D
					+		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A W B	012	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	16JAN&	0823	0823	PO	NA.

Syringe & pyride = 7.65615 g DOSAGE (total) ______ [My. Syringe & puride = 5.97785 g PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B28, B59	0	16 Jan 86	NA	0812	B28:_ *	B59: 14.98
B29, B60	0.25	, O	0838	0838	B29: *	B60: 15.25
B30, B61	0.50	1,	0853	0853	взо: 5.67	B61: 13.96
B31, B62	0.75	4	0908	0908	B31:4.49	B62: 13.38
B32, B63	1.0	• .	0923	092]	B32:6.28	B63:12.86
P33, P64	1.33	V ₁	0943	0943	B33:7.04	B64: 1223
B34, B65	1.66	1,	1003	1003	B34: 11,7	B65: //.53
P35, P66	2.0	٤ (1023	1023	B35:13.9	B66:10.94
B36	2.5	ι,	1053	1053	B36:13.2	
B37, B67	3.0	i ş	1123	1123	B37:8.75	B67:10.56
B38	3.5	i,	1153	1153	B38:122	
B39, B68	4.0	έχ	1223	1223	B39:10.2	B68: 11.28
B40	5.0	٠,	1323	1323	B40:6.62	
B41, B69	6.0	1,	. 1423	1423	B41:4,10	B69: 13.37
B42	7.0	i v	1523	1523	B42: 2.70	
B43	8.0		1623	1623	B43:250	B69A: 13.96
B44	10.0	į, i	1823	1823	B44: ₩	14.74
* Itelai	w assi	ry Ser	witivity	314		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	A W L	012	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	16 JAN86	0823	0823	PO	N.A.

DOSAGE (total) 16mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B45	12.0	16 30-86	2023	2023	B45: *	
B46, B70	24.0	17 Jan 86		0813	B46:*	B70: 15.37
		U				
			1			
					•	
	2					
* hel	awass	DIL Son	octivita			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A W B F M L	012	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	16JAN86	0823	0893	PO	NIA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	16JAN86	N.A.	0815	#64	0 m	#~0.
	0.25		0838	0839	# 64		_
	0.50		0853	0853	# 70	#20	# ok.
	0.75	k,	0908	0908	# 72		
	1.0	11	0923	0923	#64	#ND	
	1.33	C)	0943	0943	#68		
	1.66	٠,	1003	1003	# 70		
	2.00	V,	1023	1023	#60	# ND	
	2.5	در	1053	1053			
	3.0	ι,	1123	1123	#64		
	3.5	4,	1153	1153			
	4.0	L ₁	1223	1223	#84	@~#	
	5.0	ار	1323	1323			
	6.0	l.	1423	1423	#84	# ~ D	
	7.0		1523	1523			
	8.0	4	1623	1623			
	10.0	lų .	1823	1823			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	AWR FML	012	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	16 JAN86	0833	0873	PO	NIA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	16JAN86	2023	2023	# ~5		
	24.0	17 JAN &		0873	#~9	# NO	N8#
				<i>h</i>			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A W B	012	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VOLUES
Laboratory Johns Highins Highling

Screen Predrug

Study

TEST:NORMAL	QQWWWAA		qquumyy 130086		TP JUNEP	D
NA:135-148 MEQ/L	138	140		140		
K:3.5-5.0 MEQ/L	4.0	4.3		4.1		
CL:96-109 MEQ/L	106	106		108		
CO2:24-30 MEQ/L	27	26		24		
SUN:12-25 MG/DL	8	12		9		
CREAT: 0.4-1.5 MG/DL	1.0	1.]		1.2		
GLU:70-115 MG/DL	90	100		84		
T.BILI:0.3-1.2MG/DL	0.3	N.D.		N.D.		
D. BILI:0.1-0.4MG/DL	0.1	0.0		0.0	1	
CA:9.0-11.0 MG/DL	9.1	8.9		9.4		
PO4:3.0-4.5 MG/DL	2.8	3.4		2.4		
URIC A:4.2-8.8MG/DL	5.3	6.7		6.5		
T. PROT:6.0-8.5G/DL	6.1	6.2		6.0		
ALB.:3.2-5.3 G/DL	3.9	3.9		4.0		
AST:0-35 IU/L	13	19		18		
ALT:0-30 IU/L	14	9		13		
ALK PHOS:0-95 IU/L	44	38		43		
CHOL:151-268 MG/DL	ND	127 123 es		153		
LDH:0-200 IU/L	20	123	132	110	122	
CPK:0-160 U/L(male)	ND	181	94	79	78	

NB - no fellow my labo on 17 Jan 86

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	AWS FML	012	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hoghi

		Screen	Predrug	Bre gunt			Study
TEST	NORMAL	GO JAH 86	12JANSE ddmmmyy			1) JAN 86	Date
Ндь	13.9-16.3	14.1	14.7	14.7	15.1	Q.N	
PCV	41.0-53.0	41.5	42.9	46.0	443	6 14	
Plt	150-350	225	184	154	215	ND	
RBC	4.50-5.90	4.17	4.34	4.54	4.42	N.D.	
MBC	4500-11000	5700	5300	4400	6600	N.D.	
Bands	2-6%	١	10	0	Ч	ND	
Polys	31-76%	72	59	59	5. 8	20.	
Eos	1-4%	2	3	6	7	N.D.	
Bas		0	0	0	1	N D.	
Lymphs	24-44%	18	24	35	26	N.D.	
Atyp Lym		0	0	0	1	23	
Monos	2-11%	۲٦	4	5	8	N.D	
Other		٥	0	0	0	N.P.	
Retics	0.5-1.5%	20	1.0	N.D.	ND.	N.D.	
		3					
			•				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A WR F M L	012	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Clinical Pharmacol gy

		Scr	een	Predrug			Study
TEST	NORMAL		mmyy	13 Janes			Date
Color/Ap				yellan	Clear	Clear	
Sp. Gr.				1.023	1.019	1.024	
рН				6.0	6.0	70	
Protein				nee	trace	nee	
Ketones				~ev	~ 68/	ne	
Occ Bld				ney	~~	- or	
Bili.				nex	~~~~	ner	
RBC				₽	0	0	
WBC				0	0	0	
Casts				6	O.	0	
Epi. Cel				0	0	۵	
Crystals				a ()	0	٥	
Bacteria			V	0	0	Ŋ	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
BURGED	/		
147AN 86	1		
17 Jan 86	1		

INVESTIGATOR'S NAME	PT IN	IITIAL	PT	#	COMPOUND Pyridostigmine
Lietman	A L	1 L	01	7	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

ADVERSE	ONSET	CESSATION	CEUEDITY	REL'N TO	ACTION
EVENT	(Date/Time)	Date/Time		TEST DRUG	
#	dd mmm yy	dd www yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	 Treatment
			Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB.	 Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W F R	013	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	07 JAN84	Screening laboratory
_	09JAN86	History, Physical Exam
0	13JAN86	Admission
a	14JAN86	Po
5	17JAN84	IV

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WFR	013	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	M. JAN	86
			dd mmm	УУ

Examiner

Brent G. Petty

Date of birth

27/500/55 dd mmm yy

print name

Age

30 yrs

Sex

11

Race

<u>C</u>

	No	Yes	Comments
Allergy			
Tobacco Use	1		
Alcohol Use		/	Six pods prale; ou orditaile
Recreational Drug Use			
Medications past 2 weeks			
Experimental Drug Exposure		/	Interferon, cettipione.
Blood or plasma donor			
Prior Surgery			
Eye, ear, nose, throat	/		
Endocrine(diabetes,thyroid)	/		
C-V (heart murmur, HBP)	1		
Pulmonary (cough, asthma)			
Hepatitis, gastro-intestinal	/		
Genito-urinary	1		
Musculoskeletal	1		
Neuropsychiatric			
Pesticide/herbicide use			Justifizede last year (annie 1985)
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WER	013	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 1/11/1/2/ dd mmm yy

1			Blood Pressure	
_375 c	_ 68/min	20/min	1/2,78_	 _62.7

	GENERAL EXAMINATION:						
(check)	Nor.	Abn.	Provide details of abnormality	ies			
Head/Neck	/						
EENT			Sclertz QTM, (B) EHC chumière				
Chest, lungs	V						
Heart	V						
Abdomen							
Genitalia	μ	10					
Rectal	N	0					
Extremities	V						
Skin			Ears Dinder Luiger + (Planteutital aux				
Neurologic							

CHEST X-RAY
Date Lo/JAN/84

NORMAL X ABNORMAL	Describe	abnormalities:
	· 	

Examiner

Brent & Patty

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	WFR FML	013	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
S	17 JAN86	0827	0857	IV	NA

Syring + pyride = 41.77491

syring - pyride = 21.83540 DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min		
B01,B47	0	17 JANSG	0815	0815	B01: *	B47:13.41		
B02	0.08		0832	0832	B02: 8.78			
B03	0.16	• 4	0837	0837	B03:11.7			
B04, B48	0.25	٠,	0842	0842	B04:25.0	B48: 10.90		
B05	0.33	٠,	0847	0847	BO5:32.5			
B06	0.42	t,	0852	2827	BOE: 19,7			
B07, B49	0.50		0857	0857	B07:46.8	B49: 8.16		
BOB	0.58		0902	0902	BOB:15.2			
B09	0.66	1.	0907	0907	B09:/4./			
B10, B50	0.75	a _L	0912	0912	B10: //.2	B50:10.17		
B11	0.83	4	0917	0917	B11: 8.34			
B12	0.92	i,	0922	0922	B12: 6.12			
B13, B51	1.0	4	0927	0927	B13: 7,06	B51:10,75		
B14, B52	1.33	ų.	0947	0947	B14:2,50	B52: 11.60		
B15, B53	1.66	در	1007	1007	B15: 2.3A	B53:12.18		
B16, B54	2.0	4	1027	1027	B16: *	B54: 12.67		
B17	2.5	*	1057	1057	B17:-			
* lul	* below assay sensitivity 325							

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	CUFR FML	013	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	17 JAN86	0827.	0857	IV	WA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	17 JAH8	1127	1127	B18:	B55: 12.62
B19	3.5	٠,	1157	1157	B19: *	
B20, B56	4.0	1,	1227	12 27	B20:*	B56: 13.18
B21	5.0	14	1327	1327	B21:*	
B22, B57	6.0	· ·	1427	1425	B22: *	B57: 13,39
B23	7.0	4	1527	1527	B23:*	
B24	8.0	Εţ	1627	1627	B24: *	
B25	10.0	r,	1827	1827	B25: *	
B26	12.0	Łζ	2027	2027	B26:*	
B27, B58	24.0	784AC81	0827	0827	B27:*	B58:13.50
				N_		

* below assay Rensitivity 326

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WFR FML	013	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	17 JAN86	0827	0857	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	BUACTI	NA.	Not Reended	#68	# N.D.	#ok
	0.8	• •	0832	0832			
	0.16	· ·	0837	0831			
	0.25	E _X	0842	0842	# 70		
	0.33		0847	0847		,	
	0.42	٠,	0852	0852			
	0.50	۲,	0857	0827	# 78	614#	# 01
	0.58	۸.	0902	0902			,
	0.66	⁴ u	0907	0907			
	0.75	هر	0912	0912	#70		
	0.83	*	0917	0917			
	0.92	45	0922	0922			
	1.0	ι,	0927	0927	#64	m)	
	1.33	1,	0947	0947	#74	NO	
	1.66	1,	1007	1007	612#		
	2.0	(,	1027	1027	#60	# ND	
	2.5	11	1057	1057			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	WF C F M L	013	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug	admin-	End drug	admin-	Route	Bottle
Day	ddmmmyy	istration	(0-2400)	istration	(0-2400)		I.D. #
5	17 JAN86	0897		082	ጎ	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	J8UAG FI	1127	1127	#84		
	3.5	е,	(157	1157			
	4.0	4,	1227	1227	#84	# NO	
	5.0	1,	1327	1327			
	6.0	٠,	1427	1427	# 45	QN #	
	7.0	1)	1527	1257			
	8.0	• .	1227	1627			
	10.0	فر	1827	1827			
	12.0		2027	2021			
	24.0	18JDN82	0827	0827	#88	#10	# 0 K
							i.
		*					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	013	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	384AC 21	0823	6280	PO	とも

DOSAGE (total) 16 mg. organge - grado = 7.653288

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO3	RBC ACHe uM/ml/min
B28, B59	0	147AN8	. 0822	0892	B28: *	B59: 13.15
B29, B60	0.25	41	0908	8090	B59:*	B60: 13.19
B30, B61	0.50	••	0923	0927	B30:7.61	B61: 11.55
B31, B62	0.75	14	0938	0978	B31: /3,5	B62: 10.47
B32, B63	1.0	4	0953	0953	B32:12.8	B63: 9.52
B33, B64	1.33	i,	1013	1013	B33:16.8	B64: 8.86
B34, B65	1.66	``	1037	1033	B34:14,4	B65: 8,80
B35, B66	2.0	67	1053	1053	B35: A.3	B66:9.00
B36	2.5	ł s	1123	1123	B36:16.7	
B37, B67	3.0		1153	1123	B37:13.5	B67: 8.49
B38	3.5	ι,	1223	1753	B3B: 13.4	
B39, B68	4.0	((1253	1523	B39: //./	B68: 8,64
B40	5.0	ιι	1353	1323	B40: 16,2	
B41, B69	6.0	ч	425	1453	B41: 5,55	B69:10.72
B42	7.0	در	47653	1553	B42:3,/7	
B43	8.0	ار	1653	1653	B43:4,21	B69A: 11,47
B44	10.0	Ŋ	1853	1853	B44: ★	12.15

* below assay sensitivity

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	WF AL	013	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	14 JAN86	0823	0823	PO	NA

DOSAGE (total) / 16mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO)	RBC ACHe uM/ml/mir
B45	12.0	14 JAN86	2053	2053	B45: ★	
B46, B70	24.0	15 DANGE		0853	B46:Ж	B70: 13,47
			,			
					* .	
						
	,					
			·			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	F M L	013	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
2	14 JAN86	0853	0853	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	14 JAN 86	0851	2851	# 60	# NO	# ok
	0.25	. (1	2908	0908	# 72		-
	0.50	ι,	0923	0923	# 70	#ND	# 0/
	0.75	4	0738	0933	# 80		
	1.0	٤,	0953	0953	# 78	QN #	
	1.33	١,	1013	1013	#60		
	1.66	1,	1033	1033	#60		
	2.00	*1	1053	1053		eu #	
	2.5	<u>.</u>	1123	1123			
	3.0	1,	1153	1153	#60		
	3.5	١,	1223	1223			
	4.0	4,	1253	1253	#64	# ND	
	5.0	L .	1353	1353			
	6.0	(,	1453	1453 41653	#68	# ND	
	7.0	٠,	1553	41653			
	8.0		1653	1653			
	10.0	•	1853	1853			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WEL FML	013	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	14 JAN86	0853	0853	PO	NA

DOSAGE (total) 16 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	147)AN86	2053	De 53	000#		
	24.0	PANAC EI		0853	# 72	4 N D	٥£#
					-		
				9			,
				,			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WFT.	013	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

	screen	Preurug				Study
TEST:NORMAL		7378486			18 Janse ddmmmyy	Date
NA:135-148 MEQ/L	143	140		144	141	
K:3.5-5.0 MEQ/L	5.0	4.1		4.9	4,0	
CL:96-109 MEQ/L	105	104		105	1.07	
CD2:24-30 MEQ/L	25	28		24	24	
SUN:12-25 MG/DL	16	14		16	AN 19 12	
CREAT:0.4-1.5 MG/DL	1.0	1.0		1. 0	4120.9	
GLU:70-115 MG/DL	87	89		91	1 76	
T.BILI:0.3-1.2MG/DL	0.3	0.4		0.3	0.3	
D. BILI:0.1-0.4MG/DL	0.0	0.0		0.0.	0.0	
CA:9.0-11.0 MG/DL	10.1	27		10.4	9.6	
PD4:3.0-4.5 MG/DL	4.2	5:3		4.1	3.5	
URIC A:4.2-8.8MG/DL	5.1	4.9		5.9	5.3	
T. PROT:6.0-8.5G/DL	7.4	7.1		7.8	7.4	
ALB.:3.2-5.3 G/DL	4.7	4.5		4.8	4.5	
AST:0-35 IU/L	22	35		36	33	
ALT:0-30 IU/L	26	40		44	45	
ALK PHOS:0-95 IU/L	69	72		7/	70	
CHOL:151-268 MG/DL	199	212		251	153 214	
LDH:0-200 IU/L	27	123		139	137	
CPK:0-160 U/L(male)	ND	84	62	67	68	

N.B. CPK net done on 17 Jan 86

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	WFR FML	013	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Stars Hopkins Hospital

Screen Predrug Study Pre dru B84ACCO 13.JAN86 14 JAN86 15 Jan 86 18 JANS6 Date TEST NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy Hgb 13.9-16.3 14.8 14.9 15.7 14.2 15.0 PCV 41.0-53.0 42.8 44.4 43.6 46.7 41.7 Plt 150-350 340 333 323 328 336 4.50-5.90 RBC 4.89 4.84 4.75 5.26 505 WBC 4500-11000 6400 6300 8300 6100 6200 Bands 2-6% 18 17 7 Polys 31-76% 54 42 44 60 1-4% Eos 5 1 7 1 4 My o Bas 0 0 0 Lymphs 24-44% 28 34 26 38 32 Atyp Lym 2 0 0 0 Monos 2-11% 9 10 8 12 Other 0 0 0 0 0 0.5-1.5% Retics 1.5 J.D. 1.6 N.D 1.0

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	013	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Canical Planmacule gy

		Sci	reen	Predrug			Study
TEST	NORMAL		D_		15 JAN86		Date
Color/Ap				Clear	yellow	N.D.	
Sp. Gr.				1.021	1.019	1.022	
рН	-			5.0	6.0	6.0	
Protein				+ ~~	trece	ner	
Ketones				nee	mes)	~ = /	
Occ Bld	4			nec	neel	nes	
Bili.				~e1)	mej/	nex	
RBC				0	6	0	
MBC				0	O	0	
Casts				6	0	0	
Epi. Cel				O	0	0	
Crystals				.0	0	Ö	
Bacteria		,	/	0	ي	0	

ELECTROCARDIOGRAM

٦	Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
P	E8 90A 10	1		
	15 JAN86	V		a .
	BUNGC 81			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	J F R F M L	013	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

Г	10	
L	X	
Ł		
	$\overline{}$	

ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	☐ Mod	PROB.	Treatment
	*		 Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatmerit
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JDS FML	014	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
	070184	Screening laboratory
₩P	09 JM 86	History, Physical Exam
0	\$3 JAN 86	Admission
2	14:IMN 80	
_5	17 JAN 86	IV

I certify that:

1)	I	have	carefully	examined	all	entrie	s on	the	data	collection
	fo	orms,	consisting	a of LIL	5_	pages,	for	subje	ect #	OH

2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/y)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	ID S	014	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

Date of evaluation 170780 Examiner brent G. Setty

Date of birth dd mmm yy

Age

Sex

Race

tied band
in Ax1
+ puts

INVESTIGATOR'S NAME	РТ	INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	I	D 5	014	PROTOCOL DAMD 17-85-C-5133-02

PHYBICAL EXAMINATION
Date 01/09/86
dd mmm yygu

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt.(kg)
c	_ND/min	ND/min	ND ND	/76	865

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck			
EENT		V	Druse at 10 dash () eye ~ 2 dies d'auteus from disc (D) Tru el sufficientem space septement to right (D) bung & some abouchi
Chest, lungs		/	(2) lung & some shorelie -
Heart	/		
Abdomen	/ن		
Genitalia	V	p	
Rectal	U	2	
Extremities			
Skin			Tattos Ouzon arun
Neurologic	1		

CHEST X-RAY
Date 09/3AN/86

NORMAL	X	ABNORMAL	Describe abnormalities:	
V			7	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JDS FML	014	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	17 JAN8	c [86	0900	IV	กก

Superg - Pyrido = 41.94086 & DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B01, B47	0	17 DANSE	0820	0820	B01:	B47:12.46
B02	0.08	٠.	0835	0835	BO2: 11.6	
B03	0.16	1.	0840	0840	B03:17.7	
B04, B48	0.25	Z c	0845	0845	B04: 23./	B48:10.03
B05	0.33	4,	0850	0850	BO5: 19.0	
B06	0.42	1,	0855	0827	BOE:333	
B07, B49	0.50	4	0900	2900	B07:24.5	B49: 8,51
B08	0.58	T1	0905	0905	BOB: 20.2	
B09	0.66	4	0910	0910	B09://.3	
B10, B50	0.75	٠,	0915	0915	B10:6.59	B50: 9.37
B11	0.83	1,	0920	0920	B11:6,70	
B12	0.92	4	0925	0925	B12:407	
B13, B51	1.0	4,	0930	0930	B13:442	B51:10.33
B14, B52	1.33	٠,	0955	0950	B14:304	B52:10.77
B15, B53	1.66	٠,	1010	1010	B15: 2,92	B53: 11.23
B16, B54	2.0	4	1030	1030		B54: 11.68
B17	2.5	fo.	1100	1/33	B17: 1.62	

+ helminashay sensitivity 340

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JDS FML	014	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study Day	Date ddmmmyy	Start drug admin- istration (0-2400)	End drug admin- istration (0-2400)	Route	Bottle I.D. #
5	17 JAN86	0830	0690	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/MC	RBC ACHe uM/ml/mir
B18, B55	3.0	13/AUSS	1130	1130	B18: 💃	B55: 11.76
B19	3.5	0,	1200	1200	B19: <u>≭</u>	
B20, B56	4.0	4 4	1230	1230	B20: *	B56: 12.28
B21	5.0	- , '(1330	1335	B21:*	
B22, B57	6.0	'(1430	1430	B55: *	B57: 12.54
B23	7.0	11	1530	1530	B23:*	
B24	8.0	l _i	1630	1630	B24:	
B25	10.0	4	1830	1830	B25: *	
B26	12.0	4	2020	28 3°	B26:*	
B27, B58	24.0	38 MAC 81	0835	0830	B27: *	B58: 12.27
* bel	low as	Day S	ensitivita	1_		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	7 0 S F M L	019	PROTOCOL <u>DAMD 17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	17 JAN86	0830	0900	IV	NV

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	38MAC (1	0820	0820	#78	# ~D	# ck
	0.8	ϵ_{I}	0837				
	0.16	4.4	0840				
	0.25	. 1	0845		#80		
	0.33	•1	0850				
	0.42	()	0855				
1	0.50	•.	090	0900	#84	# 20	#ck
	0.58	41	0005				
	0.66	· 1,	0910	7			
	0.75	4	0915	0915	#92		
	o. 83	4	0920				
	0.92	1	0925				
	1.0	4	0135	0930	#90	2N #	
	1.33	٠,	0950	0950	#90		
	1.66	٠,	1010	1010	#88		
	2.0	٠.	1030	1030	# 72	a.v. #	
	2.5	1	1100	1/00			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	J D S F M L	014	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	17 DANS 6	08 3.0	0900	IV	NA

DOSAGE (total) 1.32mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	17 JAN86	1130	1130	#80		
	3.5	•	1200	1200			
	4.0	١,	1230	1230	#84	# ~ D	
	5.0	"\	1330	1330			
	6.0	٠,	1435	1430	*~ D	# ND	
* /	7.0	4	1530	1530			
	8.0	4,	1630	1630			
	10.0	•(1830	1830			
	12.0	(2035	2030			
	24.0	18 JAN88	O830	0830	#60	# NO	#ok

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	I D S F M L	014	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	14 JAN86	0855	0855	PO	NA

Syringe + pyrido = 7. 601788 DOSAGE (total) _____ 16 mg.

Syringe - pyrido = 5. 94663 g

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B28, B59	0	14 JAN86	0825	0825	B28:*	B59: 12.12
B29, B60	0.25	٠,	0910	0910	B29:3,5/	B60: 12.29
B30, B61	0.50	•	0925	0925	взо:4.98	B61: 11.59
B31,B62	0.75	t,	0945	0940	B31:10.7	B62: 10.05
B32, B63	1.0	6.	0955	0955	B32:14.5	B63: 9,15
B33, B64	1.33	V	1015	1015	B33:10.0	B64: 8,79
B34, B65	1.66	Ν	1035	1035	B34: 9.07	B65: 8.70
B35, B66	2.0		1055	1055	B35:16,2	B66: 8,33
B36	2.5	4,	1125	1125	B36:20,9	
B37, B67	3.0	4)	1155	1155	B37:/4,0	B67: 8.33
B38	3.5	١,	1225	1225	B38:/3.9	
B39 , B68	4.0	b t	1255	1255	B39://,6	B68: 8.57
B40	5.0	٢.	1355	1355	B40:12.7	
B41, B69	6.0	١,	1455	1455	B41: 7.59	в69:9,57
B42	7.0	14	1555	1555	B42:6.45	
B43	8.0	• `	1655	1655	B43:NS	B69A:10,63
B44	10.0	t _i	1855	1855	B44=1.9.*	11.01
* below	v assai	pensiti	vity NS 344	nosam	ple	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D S F M L	014	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	14 Jan 86	0855	0855	PO	マカ

DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B45	12.0	284BC 41	2055	2055	B45: 1.92	
B46, B70	24.0	15 JAN88		0811-	B46: *	B70:12.63
					♥.	
-						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	70 S F M L	014	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	14 JAN86	0855	0855	PO	20

DOSAGE (total) 16 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	14JAN88	N.A.	NOT Recorded	#88	# 120.	#0 k
	0.25	ε,	0910	0910	# 76		-
	0.50	ė,	0925	0925			#ok
	0.75	43	0940	0940	#76		
	1.0	1.	0955	0955	# 80	#ND	
	1.33	1,	1015	1015	#76		
	1.66	1.	1035	1035	# 78		
	2.00	h	1055	1055	# 80	# ND	
	2.5	1.	1125	1125			
	3.0	i,	1155	1155	#80		
	3.5	44	1225				
	4.0	le	1255	1255	#84	#22	
	5.0	l.	1355	1355			
	6.0	l,	1455	1455	# 80	# ND	
	7.0	e _l	1555				
	8.0	*4	1655	*/			
	10.0	6),	1855				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JDS FML	014	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	14 JAN86	0855	0855	PO	NA

DOSAGE (total) 16 mg.

PHYSIOLOGIC VARIABLES

page 2

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	14 JAN86	2055	2055	10m		
	24.0	15 JAN86		0855		# 20	8×#
	#0						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D P L	014	PROTOCOL <u>DAMD</u> 17-85-C-5133-02
	F 171 L	0. /	•

CHEMISTRY VOLUES
Laboratory Johns Hazpelal

Screen Predrug

Study

TEST:NORMAL	ddmmmyy				18JANSE JSVMMDD	Date			
NA:135-148 MEQ/L	139	139		142	143				
K:3.5-5.0 MEQ/L	4.4	4.7		4.3	4.2				
CL:96-109 MEQ/L	104	103		109	107				
CD2:24-30 MEQ/L	29	26		18	25				
SUN:12-25 MG/DL	21	12		15	15-				
CREAT:0.4-1.5 MG/DL	1.3	0.9		1. 0	1.0				
GLU:70-115 MG/DL	88	95		92	79				
T.BILI:0.3-1.2MG/DL	0.5	ND		1,0	0.5				
D. BILI:0.1-0.4MG/DL	0,1	0.1		6.0	0.1				
CA:9.0-11.0 MG/DL	9.8	96		9.6	92				
PD4:3.0-4.5 MG/DL	4.5	3.4.		4.1	3.9				
URIC A:4.2-8.8MG/DL	7,2	6.0		5.3	ND				
T. PROT:6.0-8.5G/DL	6.7	6.7		6.7	6.2				
ALB.:3.2-5.3 G/DL	4.7	4.6	·	4.6	4.5				
AST:0-35 IU/L	14	17		12	14				
ALT:0-30 IU/L	13	10		10	12				
ALK PHOS:0-95 IU/L	44	45		46	39				
CHOL:151-268 MG/DL	253	250		249	228				
LDH:0-200 IU/L	んり	160	*	148	142				
CPK:0-160 U/L(male)	N.D	218	174	152	153				

NB: CPK not done on 17 Jan 86

INVESTIGATOR'S NAME	PT INITIAL F	PT #	COMPOUND Pyridostigmine
Lietman	JDS FML	014	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

Date

TEST	NORMAL	OJJHU86	13 JAN86			
НдЬ	13.9-16.3	16.4	16.7	16.8	15.7	
PCV	41.0-53.0	47.3	50.3	48.8	45.2	
Plt	150-350	283	314	332	308	Ì
RBC	4.50-5.90	4.81	5.06	5.02	4-66	
MBC	4500-11000	7000	6000	7500	6600	
Bands	2-6%	5	2	13	4	
Polys	31-76%	54	54	40	46	
Eos	1-4%	3)	1	7	
Bas		0	Û	0	6	
Lymphs	24-44%	33	38	37	34	
Atyp Lym		0	0	3	0	
Monos	2-11%	5	5	6	9	
Other		0	0	0	0	
Retics	0.5-1.5%	h'Ú)./	N.A.	1.3	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u> </u>	014	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Clinical Pharmacology

		Screen	Predrug			Study
TEST	NORMAL		147AN86	15JAN86 ddmmmyy		Date
Color/Ap		yellas	velles	nellen	N.D.	
Sp. Gr.		1.017	1.019	1.024	1.025	
рН		5.0	6.0	5,0	6.0	
Protein		nev	mes!	000	ney	
Ketones		ner	nen	new	~~	
Occ Bld		25	res	ves	22 (184	
Bili.		~~~	new	nev	coop	
RBC			, 5	0	0	
WBC		0	0	<u>ي</u>	0	
Casts		0	Ö	0	0	
Epi. Cel		0	0	٥	0	
Crystals		0	0	0	۵	l l
Bacteria		0	0	O	۵	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
09 JAN 86	V		
15JAN86	V		axis Shifted hightward
18 Jan 86	V		axis shifted leftward toward original

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JOS FML	014	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

1	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	Mild	DEF.	None
		(0-2400)	(0-2400)	Mod /	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug
#	-	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. Poss.	Treatment
	*			□ Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAH FML	015	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
-	08 JAN 86	Screening laboratory
-	15 JAN 86	History, Physical Exam
0	19 JAN 86	Admission
2	21 JAN 86	deare ii
5	24 JAN 86	P.O.
		•
		·

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of _______ pages, for subject #__O_____.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	TAH FM L	015	PROTOCOL DAMD 17-85-C-5133-02

Date of evaluation of the print name

Date of birth of the print name

Age

Sex

Race

	No	Yes	Comments
Allergy	/		
Tobacco Use		V	Ippd x3yrs
Alcohol Use		1	1 ppod x 3 yrs 3 beers/d, oec hajur
Recreational Drug Use	/		MT years ogs
Medications past 2 weeks		V	ryleud ard ago for Hot
Experimental Drug Exposure	B	V	University of Mid (diamber stuly)
Blood or plasma donor	/		
Prior Surgery	1		
Eye, ear, nose, throat	1		
Endocrine(diabetes, thyroid)			
C-V (heart murmur, HBP)	1		
Pulmonary (cough, asthma)			
Hepatitis, gastro-intestinal	/		
Genito-urinary			BC Bell PEN
Musculoskeletal	/		
Neuropsychiatric	/		
Pesticide/herbicide use		/	Over Six wath ago
Other	/		

INVESTIGATOR'S NAME	PT	INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	714	AH	015	PROTOCOL DAMD 17-85-C-5133-02

Date dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
_37. 4c	_ 72vmin	20/min	128174	184.0	<u> 79.2</u>

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck		V	bump just left of midline or occipat from recent haven
EENT		/	Middletted new filers bitativally. (P) con common, Seften & > (2)
Chest, lungs			
Heart	V		
Abdomen			
Genitalia	M	P	
Rectal _.	^	\mathcal{D}	
Extremities	/		
Skin		/	St. legez guarted bithungh O Hank, for some fun
Neurologic			left handed

CHEST X-RAY
Date 15/1AH 86

NORMAL	\propto	ABNORMAL	Describe	abnormalities:	

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	JAH FML	015	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study Day	Date ddmmmyy	Start drug admin- istration (0-2400)	End drug admin- istration (0-2400)	Route	Bottle I.D. #	
2	21 Jan 85	0912	0942	IV	NA	
Junio - Pyrida - 42.46026 (total) 1.32 mg.						

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B01,B47	0	21 Jan 85	29	0825	B01: ¥	B47: 14.13
B02	0.08	Ű.,	0917	0917	B05:8'/	
B03	0.16	١,	0922	0922	B03: 13.9	
B04, B48	0.25	r_{i}	0927	0927	B04:18.2	B48:/1.52
B05	0.33	15	0932	0932	BO5:16.4	
B06	0.42	1,	0937	0937	BOE: 14.0	
B07, B49	0.50	4	0942	0942	B07:18.0	B49: 10.20
BOB	0.58	4.	0947	0947	BOB: 125	
B09	0.66	٠,	0952	0952	B09: 867	
B10, B50	0.75	4,	0957	0957	B10: 8.96	B50: 10.69
B11	0.83	<i>e</i> 1	1002	1002	B11:5.37	
B12	0.92	4.	1007	1007	B12:3.86	
B1 3, B 51	1.0	1.	1012	1012	B13:530	B51: //. 09
B14, B52	1.33	4.	1032	1072	B14: 25	B52: _{//.88}
B15, B53	1.66	1,	1052	1052	B15:1,97	B53: /1.97
B16, B54	2.0	r _e	1112	1112	B16:2,63	B54: 12.76
B17	2.5	4	1142	1142	B17: 1.83	

INVESTIGATOR'S NAME	РТ	INI	TIAL	PT	#	COMPOUND	Pyridostigmine
Lietman	J F	A	Ŧ	0	5	PROTOCOL	DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study Day	Date ddmmmyy	Start drug admin- istration (0-2400)	End drug admin- istration (0-2400)	Route	Bottle I.D. #
2	21 Jan 86	0912	0942	IV	24
	0				

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	21 Jan 86	1212	1212	B18:2.07	B55: 13.34
B19	3.5	0.,	1242	1242	B19: 🗡	
B20, B56	4.0		1312	1312	B20: *	B56: 13.95
B21	5.0	iį	1412	1412	B21: ¥	
B22, B57	6.0	e v	1512	1512	BSS: *	B57: /4.00
B23	7.0	4	1612	1612	B23: 🗶	
B24	8.0	f t	1712	1712	B24:*	
B25	10.0	C _C	1912	1912	B25: ★	
B26	12.0		2/12	2112	B26:*	
B27, B58	24.0	22 Jan 88	0912	0912	B27: *	B58: /3.8/
		0				

* below assay sensitivity 356

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JA H F M L	~2 K	PROTOCOL <u>DAMD 17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	21 Jan 86	0912	0942	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	21/ANE8	N N.	0825	# 50	4ND	#ok
	0.8	0	0917	0917			
	0.16	٠,	0922	0922			
	0.25	C,	0927	0927	#60		
	0.33	4.4	0932	0932	390		
	0.42	/1	0937	0937			
	0.50	٠.	0942	0942	# 48	# ND	# 01
	0.58	• (0947	0947			
	0.66	٠.	0952	0952			
	0.75	٠,	0957	0957	# 48		
	0.83	c ę	1002	1002			
	0.92		1007	1007			
	1.0	4,	1012	1012	# 56	EU #	
	1.33		1032		#50		
	1.66	٠,	1052		#52		
	2.0	•-	1112		#60	#20	
	2.5	4	1142				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	PAH	015	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	21/2018	0912	0942	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	2170086	1212	1212	#60		
	3.5	i,	1242	1242			
	4.0	1,	1312	1312	#60	#20	
	5.0	٠,	1412	1412			
	6.0	4	1512	1512	#56	a~#	
	7.0	٨	1612	1612			
	8.0	4	1712	1712			
	10.0	41	1912	1912			
	12.0	4	2 1124	2112			
	24.0	22 JAN86			#64	@N#	# _o }

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAH FML	015	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
S	24JAN86	0810	0810	PO	NA

syringe + pyrids: 7.60 gy syringe - pyrids: 5.94 g

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B28, B59	0	38uac Pc	0807	0807	B28: #	B59: 14. 29
B29, B60	0.25	t ,	0825	0825	B29: *	B60: 14.07
B30, B61	0.50	٠,'	0840	0845	B30: -≭	B61: 14.16
B31,B62	0.75	£ 1	0855	0855	B31:5./4	B62:13.48
B32, B63	1.0	·, ·	0910	09/3	B32: 7.83	B63: 12.29
B33, B64	1.33	• (0935	0930	B33:10.9	B64: 11.16
B34, B65	1.66	*1	0950	0950	B34:4.6	B65: 10.50
B35, B66	2.0	٠,	1010	1010	B35:12.3	B66: 9.53
B36	2.5	٠,	1040	1040	B36: 109	
B37, B67	3.0	C _C	1110	1110	B37:/3.5	B67: 8,77
B38	3. 5	۲,	1140	1140	B38:/2,0	
B39, B68	4.0	1	1210	1210	B39:/3.2	B68: 9, //
B40	5.0	4.	1310	1310	B40: 115	
B41, B69	6.0	٠,	1410	1410	B41:6-22	B69: 11.21
B42	7.0	1.	1510	1510	B42: 2,56	
B43	8.0	. 4,	1610	1610	B43:★	B69A:/2,74
B44	10.0	1	1810	1810	B44: *	13.32

* below away senschirty

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	DA H FM L	015-	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	247228	0810	0810	PO	

DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B45	12.0	RURCHS	2010	2010	B45: *	
B46, B70	24.0	25 JANS6		0810	B46:★	B70: 14.84
7						
			,			
						
_						
				,		
			·			
* lie	low as	bay s	ensitivit	1		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	DAH FM L	015	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	24 JAN82	0810	08/0	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	24JANS1	0807	0807	#55	# N D	# ot
	0.25	6 q	0825	0807	* 52		1
	0.50	٧,	0840	0845		£ 64	#ok
+	0.75	c,	0855	0855	#60		
	1.0	٠ ١	09/0	0910	# 60	# ~	
·	1.33	٠,	0930	0930	#64		
	1.66		0950	0950	#60		
	2.00	٠.	1010	1010		En #	
	2.5	11	1040	1040			
	3.0	٠.	1110	11/0	# 60		
:	3.5	٠,	1140	1140			
	4.0	1)	1210	1210	#64	# NB	
	5.0	r _k	1310	1310			
	6.0	/.	1410	1410	#68	#~0	
	7.0	1,	1510	1510			
	8.0	′,	1610	1610			
	10.0	1.	1810	1810			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAH FM L	015	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	BUALPE	£ 0810	08/0	PO	NA

DOSAGE (total) /6mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	24 JA1186	2010	2010	#ND		
	24.0	25 JAN86		0810		# 20	ale#
			-				
	¥						

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAH	015	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

	screen	Predrug				Study
TEST:NORMAL	QQWWWAA			22 DANS		Date
NA:135-148 MEQ/L	140	152		140		
K:3.5-5.0 MEQ/L	4.3	4.3		4.1		
CL:96-109 MEQ/L	104	116		106		
CO2:24-30 MEQ/L	26	27		28		
SUN:12-25 MG/DL	17	1)		13		
CREAT: 0.4-1.5 MG/DL	1.0	1.1		1.1		
GLU:70-115 MG/DL	83	97		82		•
T.BILI:0.3-1.2MG/DL	77	0.9		1. 5		
D. BILI:0.1-0.4MG/DL	0.1	0.1		0.0		
CA:9.0-11.0 MG/DL	10.2	10.3		9.8		
PO4:3.0-4.5 MG/DL	4.2	3.6		4. c		1
URIC A:4.2-8.8MG/DL	5.2	7.2		6.4		
T. PROT:6.0-8.5G/DL	7.3	7.1		6.8		
ALB.:3.2-5.3 G/DL	4.7	4.6		4.5		
AST:0-35 IU/L	17	27		19		
ALT:0-30 IU/L	13	11	,	6		
ALK PHOS:0-95 IU/L	65	S8		51		
CHOL:151-268 MG/DL	173	183		185		
LDH:0-200 IU/L	ND	140		123		
CPK:0-160 U/L(male)	ND	739	262	17/	125	h D

page 12Aof 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAH FM L	015	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMIBTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

AM NA 38MAC 25 Date ddmmmyy TEST: NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy NA:135-148 MEQ/L 137 K:3.5-5.0 MEQ/L 4.0 CL:96-109 MEQ/L 103 CD2:24-30 MEQ/L 22 SUN: 12-25 MG/DL 12 CREAT: 0.4-1.5 MG/DL 1.1 GLU:70-115 MG/DL T. BILI: 0.3-1.2MG/DL 1.2 D. BILI: 0. 1-0. 4MG/DL 0.0 CA:9.0-11.0 MG/DL 9.7 PD4:3.0-4.5 MG/DL 4.3 URIC A:4.2-8.8MG/DL 5.9 T. PROT:6.0-8.5G/DL 6.9 ALB.:3.2-5.3 G/DL 4.5 AST:0-35 IU/L 14 14 ALT:0-30 IU/L ALK PHOS:0-95 IU/L 48 CHOL:151-268 MG/DL 203 LDH:0-200 IU/L ND CPK:0-160 U/L(male) 105

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	JAH	015	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hogkins Vaspedal

		Screen	Predrug	On		25 JAU8	Study
TEST	NORMAL	OS JANSO Deminimby	202AUS6 ddmmmyy	ddmmmyy	gqwwwyy 3574788	34 JANSE	Date
Hgb	13.9-16.3	15.3	14.3		14.6	13.8	
PCV	41.0-53.0	43.8	41.1		42.1	40.3	
Plt	150-350	287	250		280	262	
RBC	4.50-5.90	4.75	446		4.52	4.29	
MBC	4500-11000	5300	5000		4900	4000	
Bands	2-6%	6	4		1	7	
Polys	31-76%	51	46		5/	44	
Eos	1-4%	3	0		1	1	
Bas		0	0		0	Ö	
Lymphs	24-44%	35	43		46	41	
Atyp Lym		1	0		9	0	
Monos	2-11%	4	5		1	7	
Other		٥	0		0	0	
Retics	0.5-1.5%	ND	an	N.D.		2.2	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JAHL	015	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Christophers University
Christophers Charmacology

		Screen	Predrug			Study
TEST	NORMAL	15JQN86	21JAN81			Date
Color/Ap		clear	Coar	Clear	25	
Sp. Gr.		1.017	1.024	1.017	1.020	
pН		5.0	6.5	5,0	7.0	
Protein		ree	me!	men	~ os.	
Ketones		0-64	8	2011	~ ~ ~ ~ ~	
Occ Bld		051	0200	nes	new	
Bili.		nen	~~~	رز وس	nex	
RBC			0	0		
WBC .		0	reore	0	j	
Casts		0	0	0	0	
Epi. Cel		0	0	0	0	
Crystals		٥	0	0	0	
Bacteria		0	0	0	0	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
JANAC ZI		✓	mon-specific ST-T changes & min change between
22 JAN 82		√	mon specific ST-T changes & min Changes between
25 JAN 8:		~	mon Specific ST-T changes & min Changes between trajings
			, 0

INVESTIGATOR'S NAME	PT IN	NITIAL	PT #	COMPOUND Pyridostigmine
Lietman	7	A H	0,5	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

ADVERSE EXPERIENCES

check if none occurred

abla	
	(
1/	1

ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	 None
	(0-2400)	(0-2400)	☐ Mod	PROB.	 Treatment
			Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB.	 Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A	016	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	21 144 85	Screening laboratory
-		History, Physical Exam
0	DE JAN SE	Admission
2	BUALIS	1
5	24 JANS6	P. 3

I certify that:

- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

14,01,80 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	DLA FML	06	PROTOCOL DAMD 17-85-C-5133-02

MEDICAL HISTORY

Date	of	evaluation	01,15,86
------	----	------------	----------

dd mmm yy

Examiner

Brent G. Petty , M.D

Date of birth

09/39 158

print

27yrs

Sex

Age

m

Race

B

	No	Yes	Comments
Allergy	V		
Tobacco Use		V	1/2 ppd
Alcohol Use		V	1 pint brandy week
Recreational Drug Use		1	1 pint brandy week Cocains, last 2 mage MJ 1/4 ech
Medications past 2 weeks	~		<i>y y</i>
Experimental Drug Exposure		V	Pharmakinetics - July 85, Umed 9/85, Cestriarme Supp 12/55
Blood or plasma donor		V	last Oct 1985
Prior Surgery		/	Um bilical hernia repair age 4
Eye, ear, nose, throat		V	Umbilical hernia repair age 4 Classes x 152ps
Endocrine(diabetes,thyroid)	V		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	V		suppository & 2 days GC 94h grade
Genito-urinary		V	GC 9xh grade
Musculoskeletal	V		
Neuropsychiatric	V		
Pesticide/herbicide use	V		
Other			

INVESTIGATOR'S NAME	PΤ	INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	D	L A	016	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

PHYSICAL EXAMINATION
Date 0//5/86
dd mmm yy

Temperature	Pulse	Respir	Blood P	Pressure	Height	(Cm)	Wt.(kg)
_372c	_ <u>84</u> /min	_16/min	126	188	196	0	104.8

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck		V	Large SCAR @ neck from trauma 1983
EENT		V	Gold Earning (pierced) Deartobe
Chest, lungs	V		1 cm cept just @ ion thoracic spine -
Heart	V		
Abdomen	V		Umbilical hernierahaghy scar
Genitalia	ni	þ	V
Rectal	n	D	
Extremities		V	Gear medial aspect (1) foot, superiorly
Skin	V		
Neurologic	/		

CHEST X-RAY
Date 15/3AM8L

NORMAL &	ABNORMAL	Describe	abnormalities:	
	,			

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	DLA FML	016	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	21 JAN 86	0916	0946	IV	NA

Strunge + pipido 43.42423 Strunge - pipido 23.16032

DOSAGE (total)

1.32 mg

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
BO1, B47	o	21 JAN 86	0815	0815	B01: *	B47:12.64
B02	0.08	l)	0921	0921	BO2: 7.83	
B03	0.16	11	0926	0926	B03:/0.9	
B04, B48	0.25	.′′	0931	0931	B04:14.2	B48:10,54
B05	0.33	/-	0936	0936	BO5: 14.6	
B06	0.42	Ü	0941	0941	BOE:/9,4	
B07, B49	0.50	11	0946	0945	B07:2/.0	B49: 9.35
BOB	0.58	11	0951	0951	BOB: //.0	
B09	0.66	11	0956	0956	B09: 8,64	
B10, B50	0.75	U	1001	1001	B10: 10,7	B50: 9.55
B11	0.83	11	1006	1006	B11: 7.20	
B12	0.92	()	1011	1011	B12: 5.83	
B13, B51	1.0	17	1016	1016	B13:5,85	B51: 10.21
B14, B52	1.33	11	1036	1036	B14: 2.63	B52: 10. 78
B15, B53	1.66		1056	1056	B15: 1.67	B53: //,/4
B16, B54	2.0	f ,	1116	1116	B16: ¥	B54: 11.20
B17	2.5	(1	1146	1146	B17:205	
* chil	ow assi	y Rin	citarity 3	371		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	31 JAN 86	0916	0.946	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	SI JAN86	12.16	1216	B18: *	B55: 11.26
B19	3.5	(1	1246	1246	B19:★	
B20, B56	4.0	()	1316	1316	B20:*	B56: 12.50
B21	5.0	t _i	1416	1416	B21: *	
B22, B57	6.0	Ü	1516	1516	B55: ★	B57:12.51
B23	7.0	t ·	1616	1616	B23: *	
B24	8.0	17	1716	1716	B24:*	
B25	10.0	t _l	1916	1916	B25: *	
B26	12.0	u	2116	2116	B26: *	
B27 , B58	24.0	22JAN 86	0916	0916	B27: <u>★</u>	B58:12,38
* bel	owaso	ay De	nsituity	372		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	21 JAN86	0916	0946	IV	NA

DOSAGE (total) 1.32mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	21 JAIS Elo	0815	0815	#70	# ND	# 0 K
	0.8	1,	0921	0921			
	0.16	1'	0926	0926			
	0.25	17	1931	0931	# 60		
	0.33	Ti .	0936	0936			
	0.42	17	0941	0941			_
	0.50	17	0946	0946	#70	# NO	# OK
	0.58	1/	0951	0951			
	0.66	l į	0956	0956			
	0.75	11	1001	1001	# 74		
	0.83	l)	1006	100/0			
	0.92	l i	1011	1011			
	1.0	l,	1016	1016	# 80	*20	
	1.33	ţ.	1036	1036	# 78		
	1.66	le .	1056	1056	# 77		
	2.0	f _t	1116	1110	#72	* NO	
	2.5	t _i	1146	1146			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	DLA	016	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	21JAN86	0916	0946	IV	NA

DOSAGE (total) 1,32mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	21JAN86	1216	1216	#77		
	3.5	- 11	1246	1246			
	4.0	1)	1316	1316	#76	#20	
	5.0	11	1416	1416			
	6.0	1/	1516	1516	#76°	# 200	
	7.0	11	1616	Hollo			
	8.0	li .	1216	1716			
	10.0	lı	1916	1916			
	12.0	H	2116	2116			
	24.0	22 JAN 86	0916	0916	#77	# ND	# OK

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	DLA	016	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	24 JAN 86	0813	D813	PO	NA

orginge + projudo = 7.548 DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B28, B59	0	24 JAN 86	0810	0810	B28: *	B59: 12.70
B29, B60	0.25	11	0838	0828	B29: *	B60: 12.63
B30, B61	0.50	11	0843	0843	B30: *	B61: /2.72
B31, B62	0.75	74	0858	0858	B31: *	B62:12.83
B32, B63	1.0	11	0913	0913	832: -★	B63: 12.82
B33, B64	1.33		0933	0933	B33:167	B64:12.45
B34, B65	1.66		0953	0953	вз4: 3,75	B65:11.84
B35, B66	2.0	F	1013	1013	B35:10.2	B66:10,96
B36	2.5	7-	1043	1043	B36:5.90	
B37, B67	3.0	1	1113	1113	B37:496	B67: 10.14
B38	3.5	(1	1143	1143	B38:5,30	
B39, B68	4.0	Į)	1213	1213	B39:753	B68:10.24
B40	5.0	11 -	1313	1313	B40: 7,02	
B41, B69	6.0	17	1413 .	1413	B41:6.28	B69: 10.17
B42	7.0	1/	1513	1513	B42: 4.61	
B43	8.0	(1	1613	1613	B43: 3.57	B69A:10.90
B44	10.0	11	1813	1813	B44: 220	11.74

* below assay sensitivity 375

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	24 JAN 86	0813	0813	PO	NA

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/MC	RBC ACHe uM/ml/mir
B45	12.0	24.JAN 86	2013	3013 0873 engi	B45:	
B46, B70	24.0	25 JAN86	0813	0813	B46: *	B70: 12.78
				,		
					•	
	-					
			·			
* bell	was	by se	nsituitte			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admi	in- End drug	admin-	Route	Bottle
Day	ddmmmyy	istration (0-24	400) istration	(0-2400)		I.D. #
5	24 JAN 56	0813	C	813	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	24 JAN 860	0810	0810	#80	# 110	#OR
	0.25	, ti	0828	0838	# 70		-
	0.50	11	0843	0843	#77	# 40	# OK-
	0.75	4	0858	0858	#80		
	1.0	- 41	0913	0913	#80	# 100	
	1.33	f _I	0933	0933	#80		
	1.66	17	0953	0953	#84		
	2.00	q	1013	1013	#76	# ND	
	2.5	t_l	1043	1043			
	3.0	1,	1113	1113	# 72		
	3.5	11	1143	1143			
	4.0	//	1213	1213	#76	#ND	
	5.0	1/	1313	1313			
	6.0	11	1413	1413	* 90	# ND	
	7.0	α	1513	1513			
	8.0	l,	1613	1613			
	10.0	7.5	1813	1813			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	24 JAN 86	0813	0817	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	24 JAN 86	2013	2013	# NO		
	24.0	24 JAN 86 25 JAN 86	2013 0813	2013 0813	#84	#ND	DA.
					·		

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES Laboratory Johns Hopkins Hospital

	Screen	Predrug				Study
TEST:NORMAL	20. TAN 86 ddmmmyy		QLJHN86_ ddmmmyy	22JAN86 ddmmmyy	24JAN86 ddmmmyy	Date
NA:135-148 MEQ/L	142	141		138		
K:3.5-5.0 MEQ/L	5.1	4,5		3,9		
CL:96-109 MEQ/L	107	105		105		
CO2:24-30 MEQ/L	29	24		24		
SUN:12-25 MG/DL	14	14		12		
CREAT:0.4-1.5 MG/DL	1.2	0.9		1.0		
GLU:70-115 MG/DL	93	83		153		
T.BILI:0.3-1.2MG/DL	0.6	0.7		0.7		
D. BILI:0.1-0.4MG/DL	0.1	0.0		0.1		
CA:9.0-11.0 MG/DL	10.6	9,9		10.1		
PO4:3.0-4.5 MG/DL	3.3	3,6		3.6		
URIC A:4.2-8.8MG/DL	6.1	5,4		5.6		
T. PROT:6.0-8.5G/DL	7,7	7.4		7.3		
ALB.:3.2-5.3 G/DL	4.6	4.3	٠	4.5		
AST:0-35 IU/L	21	22		21		
ALT:0-30 IU/L	21	23		14		
ALK PHOS:0-95 IU/L	44	43		43		
CHOL:151-268 MG/DL	200	191		188		
LDH:0-200 IU/L	145	140	145	147	155	
CPK:0-160 U/L(male)	10	20	227	182	200	

page 12Aof 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	Ollo	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VOLUES Hospital
Laboratory Johns Hospital

Screen Predrug

Study

Date

TEST:NORMAL	NA_	 25JAN86 ddmmmyy	qqwwwyy	ddmmmyy
NA:135-148 MEQ/L		140		
K:3.5-5.0 MEQ/L		4,1		
CL:96-109 MEQ/L		101		
CD2:24-30 MEQ/L		29		
SUN:12-25 MG/DL		12		
CREAT: 0.4-1.5 MG/DL		1.1		
GLU:70-115 MG/DL		86		
T.BILI:0.3-1.2MG/DL		0.7		
D. BILI:0. 1-0. 4MG/DL		0.1		
CA:9.0-11.0 MG/DL		10.0		
PD4:3.0-4.5 MG/DL		3.6		
URIC A:4.2-8.8MG/DL		5.5		
T. PROT:6.0-8.5G/DL		7.8		
ALB.:3.2-5.3 G/DL		4.7		
AST:0-35 IU/L		19		
ALT:0-30 IU/L		13		
ALK PHOS:0-95 IU/L		42		
CHOL:151-268 MG/DL		195		
LDH:0-200 IU/L		143		
CPK:0-160 U/L(male)		248		

INVESTIGATOR'S NAME	PT INITIAL PT #	COMPOUND Pyridostigmine
Lietman	D LA 016	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hopkins Haspitul
Whitesty Clinical Pharmaculary

Screen Predrug

Study

OJHN SC ALTHNES 12 JANSC 24 JANSC Date

TEST	NORMAL	2014NS6 ddmmmyy	al Thyse ddmmmyy	12JANS6 ddmmmyy	34 JHN SU ddmmmyy	25TRNS6 ddmmmyy
НдЬ	13.9-16.3	14.7		14.6		14.7
PCV	41.0-53.0	43,4		43.3		42.6
Plt	150-350	303		385		297
RBC	4.50-5.90	4,9		4,9		4.87
WBC	4500-11000	6900		6000		1300
Bands	2-6%	3		/	22	5
Polys	31-76%	21		31		
Eos	1-4%	4		6		27 5
Bas		1		0		\mathcal{C}
Lymphs	24-44%	65		57		57
Atyp Lym		0		3		/
Monos	2-11%	6		2		10
Other		0		0		0
Retics	0.5-1.5%	N.D.		h' p		1.2
	1					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Johns Hopkins University Clinical Pharmacology

Screen Predrug

Study

TEST	NORMAL	15JAN 86 ddmmmyy	21 JANSLe ddmmmyy			 Date
Color/Ap		clear,	yellow	yellow	not recorded	
Sp. Gr.		1.021	1.020	1.019	1.028	
рН		7.0	6.0	6.0	5.0	
Protein		TRACE	TRACE	necj	hía	
Ketones		neg	129	neg	hia	
Occ Bld	-	neg	nec;	nec	neg	
Bili.		neg	11.89	nec	NG	
RBC		D	K	0	2	
MBC		0	É	0	Q-1	
Casts		e	Ř	O	e	
Epi. Cel		e .	e	E	0	
Crystals		U	D	e	e	
Bacteria		D	Ø	U	B	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
15 Jan 86	1		
22 Jan 86	/		
25 Jan 86	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D L A F M L	016	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

- 7	1	
-1	1	71
-1		
- 1	\sim	ιI
- 1	/	N

					1	
	DVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
		e e		Sev	DEF. NOT	Stop test drug
#		dd mṁm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	☐ Mod	PROB. Poss.	Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT	INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F	- W M L	017	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	17 JAN 80	Screening laboratory
_	22 JM 86	History, Physical Exam
0		Admission
a	38 JAN &	IV ORAL
5	BURTOE	ORAL

I certify that:

1

- 1) I have carefully examined all entries on the data collection forms, consisting of 5 pages, for subject 10.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

___m.D. 24/04/86 ture dd/mmm/yy Investigator's signature

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E-W FML	017	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

Examiner

Age

2/yrs

Sex

Race

,	No	Yes	Comments
Allergy	V		
Tobacco Use		V	1 pod
Alcohol Use		V	6 pack/week
Recreational Drug Use		~	1 ppd 6 pack/week Marijiana 9 month
Medications past 2 weeks	V		
Experimental Drug Exposure		V	JHU-Ithra, Cepteraxini
Blood or plasma donor	V		
Prior Surgery		V	MVA E Lacerations 1971
Eye, ear, nose, throat	V		URI last week
Endocrine(diabetes, thyroid)	V		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)		V	Smoker's cough
Hepatitis, gastro-intestinal	V		
Genito-urinary		V	GC 1987
Musculoskeletal	V		
Neuropsychiatric	V		
Pesticide/herbicide use	V		
Other	V		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	FML	017	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date # JAN 186 # dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
_N_D: c	-64/min	16/min	124 182		65.0

-			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	V		
EENT	V		
Chest, lungs		V	Deformity of chest wall at @ Ind intercostal arheulation
Heart	1		
Abdomen	V		
Genitalia	ni)	
Rectal	n	\mathcal{D}	
Extremities	~		
Skin		V	Tatuos D Chest, (1) arm, (1) hand; scars (R) medial &
Neurologic	V		

CHEST X-RAY
Date 20120185

NORMAL ABNORMAL	Describe abnormalities:
,	

Examiner

Brent G. Petty M.P.
print name

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	017	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	27JAN86	0835	0905	IV	NA

Syringe-pyride= 42.69 Eyringe-pyride= 22,68

DOSAGE (total:

1.32 mg.

PLASMA CONCENTRATIONS

tes it	. 716	ke				
Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B01, B47	0	2754N56	0805	0805	B01: ¥	B47: 15, 73
B02	0.08	11	0840	0840	B02:8.87	
воз	0.16	žį.	0845	0845	B03:6/17	
B04, B48	0.25	/1	0850	0850	B04:129	B48:13,17
B05	0.33	(i	0855	0855	BO5: 14.2	
B06	0.42	t,	0900	0900	BO6: 16-2	
B07, B49	0.50	۲,	0965	0905	B07:20,7	B49: 1.12
BOB	0.58	f 1	0910	0910	BO8: /0,/	
B09	0.66	٠,	0915	0915	B09: 8.92	
B10, B50	0.75	l _t	0920	0920	B10:6,16	B50: //.96
B11	0.83	41	0935	0925	B11:5.22	
B12	0.92	1/	0930	0930	B12:3,82	
B13, B51	1.0	t_{ℓ}	0935	0935	B13:3,10	B51: 12.75
B14, B52	1.33	1/	0955	0955	B14: 2.29	B52: 13,78
B15, B53	1.66	1,	1015	1015	B15:/,7/	B53: 14.31
B16, B54	2.0	٠,	1035	1035	B16: ¥	B54: 14.64
B17	2.5	11	1105	1105	B17: *	

* below assay sensitivity 38

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	47	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
à	27 JAN 86	0835 .	0905	IV	nA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/HL	RBC ACHe uM/ml/mir
B18, B55	3.0	27 JAN 86	1135 Ferrass	1135	B18: *	B55:15.06
B19	3.5	U	1205	1205	B19: *	
B20, B56	4.0	q	1235	1335	B20: NR	B56:15.28
B21	5.0	4	1335	1335	B21: *	
B22, B57	6.0	4)	1435	1435	B55: *	B57:15.69
B23	7.0	r/	1535	1635	B23: 🗡	
B24	8.0	- U	1635	1635	B24: *	
B25	10.0	f.,	1835	1835	B25:	
B26	12.0	q	2035	2035	B26:*	
B27, B58	24.0	38 JAN 86	0835	0835	B27:	B58: 15.33
			0			
* Ju	clow a	ssay	Sensitive 388	ity N.R.=	not rus	n

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F m L d	17	PROTOCOL <u>DAMD_17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	27 JAN 86	0835	0905	IV	na

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	27 JAN86	0805	0805	# 50	#N0	# OK
	0.8	- 11					
	0.16	ti					
	0.25	Ü	0850	0850	# 50		
	0.33	1					
	0.42	t(
	0.50	ti	0905	0905	#52	# 100	# ar
	0.58	ii .					:
	0.66	H					
	0.75	ll	0920	0920	#60		
	0.83	t _f					
	0.92	P					
	1.0	Q.	0935	0935	#64	* 20	
	1.33	ì	0955	0955	#70		
	1.66	ť	1015	1015	#64		
	2.0	a!	1035	1035	#64	au #	
	2.5						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	FML	017	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	22 JAN 86	0835	0905	IV	カト

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	27 JAN86	1135		#64		
	3.5	1/					
	4.0	4	1235		#64	# ND	
	5.0	1,					
	6.0	11	1435		#60	# 100	
	7.0	Tr.					
	8.0	/-					
	10.0	11					
	12.0	* 11					
	24.0	28JHN86	0835		#56	# ND	# DIC-
				<i>8</i> 1			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E - W	0/7	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	30JAN 86	0855	0855	PO	NA

Syrunge + pyride = 7.6394? DOSAGE (total)

Syrunge - pyride = 5.95118 PLASMA CONCENT

16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO3 NG/ML	RBC ACHe uM/ml/min	
B28, B59	0	BOJAN SU	0840	. 0840	B28: *	B59: 15,39	
B29, B60	0.25	1/	0910	0910	B29: *	B60: 15.62	
B30, B61	0.50	U	0925	0925	B30:7.75	B61: 4.06	
B31, B62	0.75	li	0940	0940	B31: /0.6	B62: 12.64	
B32, B63	1.0	μ	0955	0955	B32://./	B63: 11.90	
B33, B64	1.33	†i	1015	1015	B33:/3,8	B64: 10.96	
B34, B65	1.66	fi	1035	1035	B34: /0.2	B65: 10.64	
P35, P66	2.0	, II	1055	1055	B35:/0.5	B66: 10.73	
B36	2.5	ħ.	1125	1125	B364.42		
B37, B67	3.0	f i	1155	1155	B37:8.37	B67: , 0	
B38	3.5	łi	1225	1230	взв:6.28		
B39, B68	4.0	Ĥ-	1255	1255	B39: 6.53	B68:11,43	
B40	5.0	11	1355	1405	B40: 4,69		
B41,B69	6.0	10	1455	1455	B41:352	B69:13,44	
B42	7.0	li.	1555	1555	B42: *		
B43	8.0	(1	1655	11055	B43: X	B69A: 14,06	
B44	10.0	11	1855	1855	B44:*	14.70	

* below assay sensitivity 391

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E - W F M L	07	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	30 an 86	0855	0855	PO	TA

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/mir
B45	12.0	30 Jan 86	2055	2055	B45: ★	
B46, B70	24.0	30 Jan 86 31 Jan 86	0855	0855	B46:*	B70: 15,41
		U		a)		
						_
					۴.	
	,					
	7					
·			•			
+ lol	MIT ON	au Du	nsitivity			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	017	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	30 Jan 86	0855	0855	PO	nA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	30 an 86	0855	0855	#45	# ND	# OK
	0.25	.00	0910	0910	# 48		-
	0.50	U	0935	0925	#40	# NO	# OK_
	0.75	i	0940	0940	#68		
	1.0	(°	0955	0955	*64	# NO	
	1.33	Į.	1015	1015	#60		
	1.66	ri .	1035	1035	#56		
	2.00	Ч	1055	1055	#60	# ND	
	2.5	H	1135	1125			
	3.0	(ı	1155	1155	#64		
	3.5	19	1225	1225			
	4.0	łř	1255	1255	#60	# ND	
	5.0	ų	1355	1355			
	6.0	ti	1455.	1455	*60	# \/D	
	7.0	ti	1555	1555			
	8.0	TI.	1655	1655			
	10.0	[1	1855	1855			

INVESTIGATOR'S NAME	PT INITIAL P	рт #	COMPOUND Pyridostigmine
Lietman	FMLO	17	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	30 Jan 86	0855	0855	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	30 Czin 87c	2055	2055	# ND		
	24.0	30 Gan 86	2055 0855	2055 0855	# NO	C W#	#
		J					
	·						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML.	017	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VOLUES
Laboratory Johns Hopkins Hospital

			/			
	Screen	Predrug	1			Study
TEST:NORMAL	17 Jan 86	269u 86 ddmmmyy	28 Jan 86 ddmmmyy	27 Jan 86 ddmmmyy	31 Jan 86 ddmmmyy	Date
NA:135-148 MEQ/L	140	no	140		141	
K:3.5-5.0 MEQ/L	4.4	no	4.7		4.7	
CL:96-109 MEQ/L	104	nd	108		102	
CO2:24-30 MEQ/L	24	no	19		30	
SUN:12-25 MG/DL	16	17	15		16	
CREAT:0.4-1.5 MG/DL	0.9	1.2	0.9		1.0	
GLU:70-115 MG/DL	82	97	88		116	
T.BILI:0.3-1.2MG/DL	1.2	0.7	0.6		0.9	
D. BILI:0.1-0.4MG/DL	_	0.1	0.0		0.1	
CA:9.0-11.0 MG/DL	9.7	9.6	8.9		9,7	
PO4:3.0-4.5 MG/DL	4.1	4.2	4.2		3,6	
URIC A:4.2-8.8MG/DL	4.8	5.2	4. U		4.9	
T. PROT:6.0-8.5G/DL	6.6	7.1			7.0	
ALB.:3.2-5.3 G/DL	4.9	4.7	4.3		4.9	
AST:0-35 IU/L	//	34	22		17	
ALT:0-30 IU/L	4	16	8		13	
ALK PHOS:0-95 IU/L	49	48	45		45	
CHOL:151-268 MG/DL	219	212	186		227	
LDH:0-200 IU/L		167	179	172		
CPK:0-160 U/L(male)		492,	270	31.71		

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	F M L	117	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Johns Applins Hospital

Screen Predrug

Study

TEST	NORMAL	17 Jan 86 ddmmmyy	26 Jan 86 ddmmmyy	28 Jan 80	31 Jan 82 ddmmmyy	qqwwwyy	Date
Hgb	13.9-16.3	14.1	15.5	14,5	15,5		
PCV	41.0-53.0	41.7	45.3	42.5	45,3		
Plt	150-350	343	218	186	364		
RBC	4.50-5.90	4.57	5.07	4.75	5.13		
WBC	4500-11000	6000	4200	4200	4700		
Bands	2-6%	3	0	9	3		
Polys	31-76%	42	41	40	39		
Eos	1-4%	2	0	4	6		
Bas		/	0	/	/		
Lymphs	24-44%	47	40	36	48		
Atyp Lym		0	2	/	0		
Monos	2-11%	5	17	9	3		
Other		0	0	0	0		
Retics	0.5-1.5%	N.D.	-4D-	N.D.	N.D.		
				,			
		2					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	017	PROTOCOL DAMD 17-85-C-5133-02

Laboratory John Hopkins University Clinical Pharmacolcy

screen	Preorug				
22 Jun 56	28 Jan 86	31			

Study

TEST	NORMAL	22 Vin SU ddryfnmyy	38 Jan 86	31 Jun 86 ddmmmyy	qqwwwyy	ddmmmyy	Date
Color/Ap		Clear	yellow	Cloudy			
Sp. Gr.		1.020	1.027	1.019			
рН		7.0	6.5	6.0			
Protein		NG	- hea	neg	à.		
Ketones		nog	200	néa			
Occ Bld		neg	nea,	nea			
Bili.		nsc	nac	nlu			
RBC		e	E	Q			
WBC		P	Q	Ð			
Casts		8	Û	B			
Epi. Cel		D	D	Ø			
Crystals		Q	S	B			
Bacteria		₽′	Ø	D			

ELECTROCARDIDGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
22 Jan 86	V		
28 Jan 86 31 Jan 86	V		
31 Jan 86	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	E - W F M L	017	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

ADVERSE EXPERIENCES

check if none occurred



ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT	INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W. F	H_K	018	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

PROTOCOL

Study Day	Date ddmmmyy	Procedures
1		Screening laboratory
_	22JMV 8U	History, Physical Exam
0	26 JM 86	Admission
2	38MAC CE	IV 6 ORAL
5	30JAN8	CORAL

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of \sqrt{S} pages, for subject #OLS.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature M.D. 24/00/06 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	WHU FML	018	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

Date of evaluation #13 186 Examiner BRENT G. PETTY, M.D.

Date of birth 11,940,63 dd mmm yy

Age 23 JHN MEDICAL HISTORY

Examiner BRENT G. PETTY, M.D.

print name

Sex

Race

	No	Yes	Comments
Allergy	V		
Tobacco Use		V	1/2 ppd .
Alcohol Use		/	1 6 pack/week
Recreational Drug Use		V	mJa4weeks, last months age
Medications past 2 weeks		V	Fylenol ~ 2 d ago for HA
Experimental Drug Exposure		V	Jobra mycin
Blood or plasma donor		/	
Prior Surgery		V	Notunteer blood done at hospital Repair of hypospadia age 4 had bustered preumothorax &'d & chest out
Eye, ear, nose, throat		V	Wears glasses
Endocrine(diabetes,thyroid)	V		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal	V		8
Genito-urinary	V		
Musculoskeletal	V		
Neuropsychiatric	V		
Pesticide/herbicide use		V	2 dogs ¿ bathe for flew today
Other	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	W / W	018	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 22 / AN / 86 dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
_368c	_ <u>88</u> _/min	20/min	120 182	192.0	78.8

	GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities					
Head/Neck	V		Shoty node R & area					
EENT		V	TeleRotic TM's belaterally, Masal septem deviated to (1). Discoloration both top front feeth					
Chest, lungs	1							
Heart	1							
Abdomen	V							
Genitalia	no	5						
Rectal	ns							
Extremities	1							
Skin		1	Tattos R) shoulder + Q forearm, scars anterior elect belaterally from chest hibs docements, scars (Q) forearm, small lateration & produmal thumb					
Neurologic	/		Small laderation & productal thumb					

CHEST X-RAY Date 22/Jun/84

NORMAL X ABNORMAL	Describe abnormalities:
	0 1- 0
E	Examiner Beent Wetty My
	ORENT G. PETTY, M.D.

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H L	018	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	27 Jan 86	0838	1350908	IV	nA

Syng + Oyida = 42.438 Syng - Oyida = 22.4798

DOSAGE (total) 1.32mg

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
BO1, B47	0	27 Jan 86	0805	0805	B01: *	B47:13,41
B02	0.08	11	0842	0847	BO2: 6.90	B48
воз	0.16	14	084 8 5	0848	B03:/0.5	
B04, B48	0.25	ϵ_t	085 3 5	08535	B04:/2.3	B48 B 11.77
B05	0.33	C;	08585	0858 3	B05:/2.9	
B06	0.42	(/	0903 (3)	09033	BOE: 20.6	
B07, B49	0.50	(i	090859	0908 5	B07: /8.9	B49: 10,41
BOB	0.58	tt	0913	0913	BOB:/4,5	
B09	0.66	Ú	0918	918	B09: 10.5	
B10, B50	0.75	r _e	0923	0923	B10:5,67	B50:10.82
B11	0.83	łe	0928	0928	B11:5.50	
B12	0.92	(0933	0933	B12:4,6/	
B13, B51	1.0	t.	0938	0938	B13:4.37	B51: //.47
B14, B52	1.33	le	1958	0958	B14:3,19	B52:12.04
B15, B53	1.66	1,	1018	1018	B15:2.34	B53: 12.65
B16, B54	2.0	I,	1038	1038	B16: 🗶	B54: 12.80
B17	2.5	014 00	1108	1108	B17: *	

* below assay sensitivity 402

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W K L	018	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	37 Jan 86	0838 .	0908	IV	'nΑ

DOSAGE (total) 1,32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	22 Jansle	1138	1138	B18: *	B55:12,73
B19	3.5	li li	1208	1208	B19: *	
B20, B56	4.0	ř <u>.</u>	1238	1238	B20: *	B56: 13.41
B21	5.0	t _i	1338	1338	B21:*	
B22, B57	6.0	()	1438	1438	B55:*	B57: 13.40
B23	7.0	l _i	1538	1538	B53:¥	
B24	8.0	(t	1638	1638	B24: *	
B25	10.0	k	1838	1838	B25:*	
B26	12.0	a	2038	2038	B26:*	
B27, B58	24.0	28 Jan 86	0838	0838	B27: *	B58:13.30
		U				
* belo	w Do	Day 1	lensdirt	403		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H L	018	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-		Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	27 Jan 86	0838	0908	IV	nA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	27 Jule	0805	0805	#70	# 20	#OK.
	0.8	J. O					
	0.16		,				
	0.25	17	0853	0853	#68		
	0.33						
	0.42						
	0.50	h	0908	0908	# 78	# ND	# OC
	0.58						
	0.66						
	0.75	11	0923	0923	# 70		
	0.83						
	0.92						
	1.0	l)	0938	0938	#80	# NO	
	1.33	1,	0458	0958	# H6		
	1.66	1,	1018	1018	#82		
	2.0	1,	1038	1038	#76	# 20	
	2.5						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	W K K	D18	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	27 Gan 86	0838	098	IV	nA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	27 Jan 86	1138	1138	# 70		
	3.5	3					
	4.0	Į!	1238	1238	#76	# 10	
	5.0	li .					
	6.0	1.	1438	1438	#72	# 20	
	7.0	17					
	8.0	Ĭŧ					
	10.0	()					
	12.0	li .					
	24.0	28 Jan Ste	0838	0838	# 70	# わり	# ac
		Ü					
					//		
		3					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H U F M L	018	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	30 Jan 86	0900	0900	PO	nA
Sanina	o + nni	do = 7.63564	1/		

Springe + pipido = 7.63564 Syringe - pipido = 5.94243

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B28, B59	0	30 Jan 56	0843	1843	B28: *	B59: 13.45
B29, B60	0.25	٧,,	0915	0915	B29: *	B60: 13, 35
B30, B61	0.50	1/	0930	0930	взо:8.04	B61: 12.28
B31,B62	0.75	l/	0945	0945	B31:17.9	B62: 10.99
B32, B63	1.0	1,	1000	1000	вз2:/4.8	B63:10,77
B33, B64	1.33	1/	1020	1020	B33://.8	BE4: 10,48
B34, B65	1.66	li .	1040	1040	B34:11.9	B65: 10.65
B35, B66	2.0	17	1100	1100	B35:7,74	B66: 10.73
B36	2.5	<i>I</i> /.	1/30	1130	B36:6.0/	
B37, B67	3.0	11	1200	1200	B37:5.26	B67: 11.17
B38	3.5	ţ!	1230	1230	взв :4,28	
B39, B68	4.0	U.	1300	1300	B39:3,45	B68: 11.47
B40	5.0	//	1400	1400	B40:306	
B41,B69	6.0	1,	1500	1500	B41: 🗶	B69: 12 8
B42	7.0	1,	1600	1600	B42: *	
B43	8.0	Ž×.	1700	1900	B43:*	B69A:13.03
B44	10.0	t .	1900	1900	B44: *	13.22

* below assay sensitivity 406

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	WHU	018	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	30 Jan 86	0.900	0950	PO	NA

DOSAGE (total) ______/6 mg

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B45	12.0	30 Jan 86	2100	2100	B45: *	
B46, B70	24.0	3 Jau 86	0900	0900	B46:★	B70: 13,94
		V				
			-			
					•	
					·	
* bela	u assa	It som	o trivital			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	018	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	30 Jaule	0900	0900	PO	nA

DOSAGE (total) 16 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0			0900	#72	400	# OK
	0.25	·		0915	# 70		-
	0.50			0930	#12	# 10	# OK-
	0.75			0945	# 80		
	1.0			1000	#78	# 20	
	1.33			1020	# 80		
	1.66			1040	#80		
	2.00			1100	#76	# 20	
	2.5						
	3.0			1200	# 84		
	3.5						
	4.0			1300	# 80	# ND	
	5.0						
	6.0			1500	#80	# ND	
	7.0						
	8.0						
	10.0						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W M L	D18	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	30 Jun 86	0900	0900	PO	RA

DOSAGE (total) 16 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate .bpm	Hand Grip L/R	Coord Test
	12.0	30 Can 86	2100	2100	# 20		
	24.0	30 Can 86 31 Jan 86	0900	0900	6N#	4V5	N D#
		V					

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H LL	018	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VOLUES
Laboratory Johns Hipkins Haspidal

Screen Predrug

Study

TEST:NORMAL	21 Gan 86 ddmmmyy	ddmmmyy		289ans6 ddmininyy	31 Jan Ste ddminmyy	Da
NA:135-148 MEQ/L	141	141		141	141	
K:3.5-5.0 MEQ/L	4.2	4,1		4.3	4.3	
CL:96-109 MEQ/L	103	105		106	101	
CO2:24-30 MEQ/L	28	28		29	27	
SUN:12-25 MG/DL	8	13		//	12-	
CREAT:0.4-1.5 MG/DL	1.0	1,1		0.9	0.9	
GLU:70-115 MG/DL	130	76		82	76	
T.BILI:0.3-1.2MG/DL	0.7	0,7		0.6	0.6	
D. BILI:O. 1-O. 4MG/DL	0.1	0.1		0.0	0.0	
CA:9.0-11.0 MG/DL	8,8	9.2		9.0	9.7	
PD4:3.0-4.5 MG/DL	3.3	2.9		3.1	3,3	
URIC A:4.2-8.8MG/DL	4.6	4.8		4.3	5,5	
T. PROT:6.0-8.5G/DL	6.9	6.5			6.8	
ALB.:3.2-5.3 G/DL	4.3	4.1		4,1	4.6	
AST:0-35 IU/L	17.	12		12	22	
ALT:0-30 IU/L	10	19		8	14	1
ALK PHOS:0-95 IU/L	50	48		43	47	
CHOL:151-268 MG/DL	153	157		151	184	
LDH:0-200 IU/L	_	118	136	121		
CPK:0-160 U/L(male)		74	57	53	_	

page 13 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	d8	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Ahns. Hopkins Hopeful

Screen Predrug

Study

TEST	NORMAL	21 gan 86 ddmnmyy	36000 810	20 Jun 80	31 Okn 86 ddimmiyy	ddmmmyy	Date
НдЬ	13.9-16.3	15.0	14.7	14.6	15.1		
PCV	41.0-53.0	44.8	43.6	42,5	44,8		
Plt	150-350	161	216	257	337		
RBC	4.50-5.90	4.78	4.61	4.52	4.85		
MBC	4500-11000	6700	5300	6000	7300		e .
Bands	2-6%	16	5	6	3		
Polys	31-76%	58	49	50	47		
Eos	1-4%	3	4	35	3		
Bas		0	0	0	0		
Lymphs	24-44%	17	37	31	40		
Atyp Lym		0	/	U	/		
Monos	2-11%	6	9	10	6		
Other		0	0	0	0		
					,		
Retics	0.5-1.5%	0.3	ND	20	ND		
			•				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	W H L	018	PROTOCOL DAMD 17-85-C-5133-02

Laboratory China Hopkins University Clinical Pharmer cology

Screen Predrug

Study

TEST	NORMAL	32 Jan 80 ddmmmyy	28 Jan Sle ddmmmyy	31gan86 ddmmmyy	ddmmmyy	ddmmmyy	Date
Color/Ap		llear	yellow	Cloudy			
Sp. Gr.		1.022	1.020	1.019			
pН		7.0	6,5	6.0			
Protein		neg	- heg	NS G			
Ketones		neg	neg	neg			
Occ Bld		neg	neg	neg			
Bili.		nea	ntg	nsc			
RBC		NG O	Ð	D			
WBC		E	B	rare			
Casts		E	O	e			
Epi. Cel		t	R	R			
Crystals		Ð	Ŕ	R			
Bacteria		O	Ø	Ø			

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
22 Jan 86			
22 Jan 86 28 Jan 86	V		
31 Jan 86	/		

INVESTIGATOR'S NAME	PT INIT	IAL	PT #	COMPOUND	Pyridostigmine
Lietman	W K	<u> </u>	D18	PROTOCOL	DAMD 17-85-C-5133-02

ADVERSE EXPERIENCES

check if none occurred

-1	1	1
-1	1	/
	_\	
·ŧ		

ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	☐ Mod	PROB.	Treatment
			□ Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
			Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	MEC FML	019	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	17JM) 86	Screening laboratory
_	22 JAN 86	History, Physical Exam
0	D HATCE	Admission
2	28JAN8C	エソ
5	31JM86	ORM

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 15.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

m.D. 2V,Oct, & dd/mmm/yy

INVESTIGATOR'S NAME	PT	INI	TIAL	PT #	COMPOUND Pyridostigmine
Lietman	MF	E M	<u>C</u>	019	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation 22/401/86 dd min yy
		dd Minin yy

Examiner

BRENT G. PETTY, M.D.

Date of birth

10/APR/63

Age

2 yrs

Sex

01

Race

 \mathcal{B}

	No	Yes	Comments
Allergy	/		
Tobacco Use		V	1/2 ppd
Alcohol Use		/	6 pack/weck
Recreational Drug Use	/		<i>O</i>
Medications past 2 weeks	V		
Experimental Drug Exposure		V	Pharmakinetics 8/85
Blood or plasma donor	V		
Prior Surgery		/	Umbilical hermorrhaphy age 10
Eye, ear, nose, throat	V		
Endocrine(diabetes, thyroid)			
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal	V		
Genito-urinary		/	april 1985 GC & PCN
Musculoskeletal	V		
Neuropsychiatric	V		
Pesticide/herbicide use		/	Exterminators Springed 4/85
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	A E C F M L	019	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

PHYSICAL EXAMINATION Date 2 186 dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
36.6 c	_ <u>80</u> _/mir	16_/min	116_178_	178.0	-61-62

			GENERAL EXAMINATION:
(check)	Nor.	Abn.	Provide details of abnormalities
Head/Neck	~		
EENT	1		
Chest, lungs	1		
Heart	1		
Abdomen		·/	Deformed remoderces
Genitalia	no	0	
Rectal	n	W	
Extremities	1		
Skin		V	Patdy supersignoutation on truck , ned, and arms , poil tures successor. Sear (Foreaun from childhood burn
Neurologic	/		let handed, left goip > (R)

CHEST X-RAY
Date A Au / 86

NORMAL ABNORMAL	Describe	abnormalities:
,		

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	M E C F M L	019	PROTOCOL DAMD 17-85-C-5133-02

Pt's at 60.8 My MEDICATION RECORD
STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	28 JH-84	0917	0947	IV	NA

synge + pyride = 42.73

synige - gyride = 22.95 DOSAGE (total) 1.32 mg

PLASMA CONCENTRATIONS

NB: This page was created on 22 APR86. Times for BOZ-BOS								
	Schedule Time (hours)		Scheduled Clock Time (0-2400)	Actual	[PYRIDO]	RBC ACHe uM/m1/min		
B01, B47	0	28JAN86	0807	0807	B01:*	B47: //./8		
BOS	0.08	7 \	0922	0922	Bo5: 10'8			
B 03	0.16	٠. ١.,	0927	0927	BO3: /7./			
B04, B48	0.25	٠.	0932	0932	B04:/7.5	B48:9,08		
B05	0.33	и	0937	0937	B05: 8.6			
B06	0.42	٠,	0942	0942	BOE: 14.2			
B07, B49	0.50	-,	0947	0947	B07: 242	849: 7.21		
BOB	0.58	Q.	0952	0952	BOB:/7.6			
B09	0.66	Ĺ	0957	0957	B09:10,/			
B10, B50	0.75	C.	1002	1002	B10:8,42	B50: 7.95		
B11	0.83	٠,	1007	1007	B11:693			
B12	0.92	ζ,	1012	1012	B12:6.80			
B13, B51	1.0	٠,	1017	1017	B13:7/2	B51: 8.07		
B14, B52	1.33	٠,	1037	1037	B14:5.07	B52: 8,61		
B15, B53	1.66	\ \ \	1057	1057	B15:3,16	B53: 9.36		
B16, B54	2.0	٠,	1117	1117	B16: /.8/	B54: 9.10		
B17	2.5		1147	1147	B17: 6.59			
* Dula	ILLOUDDIE	Almaix	THETE!					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	MEC FML	019	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	28 Jan86	0917	0947	IV	MA

DOSAGE (total) 1.32 mg

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	[PYRIDO]	RBC ACHe uM/ml/min
B18, B55	3.0	28 Jan 86	1217	1217	B18:3,66	B55: _{10:38}
B19	3.5	١,	1247	1747	B19: 3.7/	
B20, B56	4.0	£ f	1317	1317	B20: 1,54	B56:10.58
B21	5.0	Ü	14 17	1417	B21: *	
B22, B57	6.0	10	15.17	1517	B55: *	B57: 10.83
B23	7.0	(i	1617	1617	B23: *	
B24	8.0	G	1717	1717	B24: *	
B25	10.0	¢ c	1917	1917	B25: *	
B26	12.0	(1	2117	2117	B26: NR	
B27, B58	24.0	29 Jan 56	0917	0910	B27: *	B58: 10,97
		J	,			
* beli	ow ax	Day se	nsitivity	V 418 N.R. =1	bt Run	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	A E C	DIG .	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	289ax86	0917	0947	IV	NI

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	28 Jansti	0915	0915	#60	# NO	# 016.
	0.8	J					
	0.16						
	0.25	D.	0932	0932	#62		
	0.33						
	0.42						
	0.50	ti -	0947	0947	#69	# 20	# OK
	0.58						
	0.66						
	0.75	11	1002	1002	#68		
	0.83						
	0.92						
	1.0	11	1017	1017	* 70	# ND	
	1.33	1/	1037	1637	# 70		
	1.66	11	1057	1057	# 68		
	2.0	1ì	1117	1117	#68	# NO	
	2.5						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	ME C	019	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	28 Can86	0917	0947	IV	NA

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	28 Gav. 86	1217	1217	#72		
	3.5	U					
	4.0	11	1317	1317	#68	# ND	
	5.0						
	6.0	11	1517	1517	#68	# ND	
	7.0						
	8.0						
	10.0						
	12.0						
	24.0	29 Janslo	0917	0910	#70	# ND	# OR
		J					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	MEC FML	019	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	31 Jan 86	0811	0811	PO	NA

Signings + pyrido = 7.63020 g DOSAGE (total) _ $\frac{1}{2}$ Signings - pyrido = 5.95638 g PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/MC	RBC ACHe uM/ml/min
B28, B59	0	31 Jan 86	0811	0808	B28: *	B59: 11.17
B29, B60	0.25	11	0826	0826	B29:4,16	B60: 10.06
B30, B61	0.50	1/	0841	0841	B30: 15A	B61: 8.96
B31, B62	0.75	11	0856	0856	B31:8,A	B62:8.22
B32, B63	1.0	I _l	0911	0911	B32:13.3	B63: 7,7/
B33, B64	1:33	l)	0931	0931	B33: //.0	B64: 7.78
B34, B65	1.66	- 11	0951	0951	B34:8.45	B65: 8.20
B35, B66	2.0	l)	1011	1011	B35:839	B66: 8.42
B36	2.5	11.	1041	1041	B36:6.22	
B37, B67	3.0	11 -	(1/1	1111	вз7:5.04	B67: 8,86
B38	3.5	lį	1141	1141	B38: 3,74	
B39, B68	4.0	ĺι	1211	1211	вз9:5,38	в68: 9,25
B40	5.0	Ĭţ	13/1	1311	B40: 3.99	
B41, B69	6.0	11	1411	1411	B41:7.02	B69: 9.80
B42	7.0	t_{i}	1511	1511	B42:259	
B43	8.0	Ėį	1611	1611	B43:1.78	B69A: 10.37
B44	10.0	11	1811	1811	B44: *	B69B: 10.30

* below assay sensitivity 421

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M E C	019	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	3i Jan 86	0811	0811	PO	nA

DOSAGE (total) 16 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/mir
B45	12.0	31 Jax 86	2011	3011	B45: *	
B46, B70	24.0	31 Jax 86	2011 0811	30 11 0811	B46: 💥	B70:/1.10
						
					•.	
	187					
	ur assa					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	ME C F M L	019	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	31 Jau 80	08/1	0811	PO	72 A

DOSAGE (total) 16 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	31 Junsto	08/1	0808	# 62	מע#	# 0 K.
	0.25	. (,	0826	0826	#60		_
	0.50	11	0841	0841	#58	# ND	# OK
	0.75	1,	0856	0856	#54		
	1.0	71	0911	0911	* 56	# NS	
	1.33	P.	0931	0931	#56		
	1.66	1:	0951	0951	# 56		
	2.00	U	1011	1011	#62	# ND	
	2.5						
	3.0	fi	////	////	#60		
	3.5						
	4.0	1/	12.11	1211	#70	# ND	
	5.0						
	6.0	11	1411	1411	*64	# NO	
	7.0						
	8.0						
	10.0						

N.B.: Bubget was something at 4.0 hours when

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M L C	019	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	31 Jan 80	0811	0811	PO	nA

DOSAGE (total)

16 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	3 Jan 84	2011	2011	# ND		
	24.0	3 Jan 84 13eb86	2011 0811	2011 0811	* 54	# 20	ac#
	,						
	*						
				-			

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	MEC	09	PROTOCOL <u>DAMD</u> 17-85-C-5133-02
		, 1	

CHEMISTRY VOLUES
Laboratory Chino Hopkins Laopital

	Screen	Predrug				Study
TEST:NORMAL	17 Mu86	27 Jun 80 ddmmmyy	26 Mr 80	29 Jan 810 dd iminingy	<u>01 76686</u> ddmmmyy	Date
NA:135-148 MEQ/L	145	140		138	140	
K:3.5-5.0 MEQ/L	4.1	4.5		4.2	3.9	
CL:96-109 MEQ/L	109	104		107	102	
CO2:24-30 MEQ/L	25	28		26	28	
SUN:12-25 MG/DL		16		14	17	
CREAT: 0. 4-1.5 MG/DL	1.0	1.1		1.1	1.1	
GLU:70-115 MG/DL	95	94	er e	75	ln 4	
T.BILI:0.3-1.2MG/DL	0.6	0.4		0.4	0,7	
D. BILI:0.1-0.4MG/DL	0.1	0.0		0.1	0.1	
CA:9.0-11.0 MG/DL	9.9	10,5		9.2	10.2	
PO4:3.0-4.5 MG/DL	4.0	4,1		3.1	4.6	
URIC A:4.2-8.8MG/DL	7.2	8.3		7.5	7.5	
T. PROT:6.0-8.5G/DL	7.2	7.3		6.3	10,9	
ALB.:3.2-5.3 G/DL	4.5	4.8		4.2	4,5	
AST:0-35 IU/L	19	17		4	13	
ALT:0-30 IU/L	(1	5		7	9	
ALK PHOS:0-95 IU/L	50	49		47		
CHOL:151-268 MG/DL	178	191		182	99X2	
LDH:0-200 IU/L		99		108	93	

CPK:0-160 U/L(male)

INVESTIGATOR'S NAME	PT INITIAL PT	
Lietman	ME CO	PROTOCOL DAMD 17-85-C-5133-02

Laboratory VALUES Hopkins Hospital

		Screen	Predrug	(5)01			Study
TEST	NORMAL	17 Jan 80 dd mmyy	27 Jan Slo ddmmmyy	29 941186 ddmmmyy	3 76686 ddmmmyy	ddmmmyy	Date
Ндь	13.9-16.3	16.0	16.9	15,2	16.3		
PCV	41.0-53.0	46.2	48.4	44.7	47.3		
Plt	150-350	216	211	191	197		
RBC	4.50-5.90	5.24	5.51	5.02	5.3		
WBC	4500-11000	5100	4910	6500	6200		
Bands	2-6%	1	4	5	3		
Polys	31-76%	40	48	57	54		
Eos	1-4%	3	2	C	2		
Bas		Û	/	/	0		
Lymphs	24-44%	44	34	30	31		
Atyp Lym		0	/	1	C		
Monos	2-11%	12	10	6	10		
Other		0	0	0	0		
Retics	0.5-1.5%	_			0.8		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M L C	019	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

Laboratory Clinical Pharmacology

		Screen Predrug					Stud
TEST	NORMAL	22 Jan 86	27 Jul 80 ddmmmyy	24 July ddidminyy	1 1eb 80 ddmmmyy	ddmmmyy	Dá
Color/Ap		yellow	Clear	clea R	_		
Sp. Gr.		1.021	1.017	1.017	1.030		
pН		6.5	5.0	5.0	5.0		
Protein		TRACE	IRACE.	N. G	neg		
Ketones		nich	neg	neg	neg		
Occ Bld		neg	neg	hey	neg		
Bili.		Neg	neg	nog	neg		
RBC		e	2	Ø	E		
WBC		e	Ø	Ø	0-1		
Casts		0	B	Ð	D		
Epi. Cel		8	Ø	Ø	2		
Crystals		Ø	Ø	8	Ø		
Bacteria		Ø	Ø	Ø	Ø		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
22 Jan 86	/		
39 Gay 86	V		
OI Feh 80			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M L 2	019	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

1	ı
	ı
	ı
	ŀ

ADVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	dd mmm yy	dd mmm yy	Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB.	Treatment
	,		Sev	DEF. NOT	Stop test drug
#	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	(0-2400)	(0-2400)	Mod	PROB.	 Treatment
			 Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FMR)53 C	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
	23JM 86	Screening laboratory
_	24 JAN 86	History, Physical Exam
0	28 WATERS	Admission
a	28JAN 86	IV
5	31JAN86	Po

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject #20.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

_M.D. <u>U/Ot/No</u>
re dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	J TR	20	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation 24/JAP/86	•
		dd mmm vv	

Examiner

,			
Bren	7 6.	Petty.	
	print	name 0	

Date of birth

17/JAN 86 dd mmm yy

....

Age

24 yrs

Sex

 $\overline{\omega}$

Race

B

	No	Yes	Comments
Allergy	~	3	
Tobacco Use		V	Igrd
Alcohol Use		/	6 pack + 2-3 nixed Links (weeken)
Recreational Drug Use		/	m-J 1985
Medications past 2 weeks			
Experimental Drug Exposure	/		
Blood or plasma donor		V	lost July 1985
Prior Surgery	/	=	
Eye, ear, nose, throat	~		
Endocrine(diabetes,thyroid)	~		
C-V (heart murmur, HBP)			
Pulmonary (cough, asthma)	V		
Hepatitis, gastro-intestinal	/		
Genito-urinary	0		
Musculoskeletal	V		
Neuropsychiatric			
Pesticide/herbicide use		/	Externinators in at. 1985
Other	-		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	J T R	020	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION

Date 24/ JAN 81 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height (cm	Wt. (kg)
37.60	_ & Omin	16/min	10	8164	18L.C	76.2

	GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities					
Head/Neck		/	slightly asymmetric through & A (D) lobe					
EENT		/	Slightly asymptric thyroid & A (D) lobe Nasal septem deviated to (D), text in Fair - poor ?					
Chest, lungs	V							
Heart	/							
Abdomen	1							
Genitalia			C'D					
Rectal			0'7					
Extremities	V	·						
Skin		~	small scar @ flook & st. Keloid formation					
Neurologic	V							

CHEST X-RAY
Date 24/ JAH 86

NORMAL	× ABNORMAL	Describe abnormalities:	
		- 	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	JIR	000	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study Day	Date ddmmmyy	Start drug admin- istration (0-2400)	End drug admin- istration (0-2400)	Route	Bottle I.D. #
2	28 JAN86	0923	0953	IV	NA

Syringe + gyride = 42.36g Syringe - pyride = 22.38g

DOSAGE (total) _

1,32 mg

PLASMA CONCENTRATIONS

Suby	subject's wt: 74.4 kg							
Sample No.		Date ddmmmyy	Scheduled	Actual Clock Time (0-2400)	DG/MC	RBC ACHe uM/m1/min		
B01, B47	0	28JAN86	NA 9	0800	B01:*	B47:12.53		
B02	0.08	1,	08.28	0878	B05:9.09			
B03	0.16	١.	0933	0933	B03:/2.6			
B04, B48	0.25	1,	0938	0938	B04:205	B48:10,17		
B05	0.33	••	0943	0943	B05: 199			
B06	0.42	11	0948	0948	BOE:26.8			
B07, B49	0.50	1.	0953	0953	B07:295	B49:8.53		
BOB	0.58	4	0958	0958	BOB:/8./			
B09	0.66	٠.	1003	1003	B09:16.8			
B10, B50	0.75	14	1098	1008	B10: 10.3	B50: 9,12		
B11	0.83		1013	1013	B11:6.98			
B12	0.92	1,	1018	1018	B12: 6.26			
B13, B51	1.0	4,	1023	1023	B13:5./8	B51:9,71		
B14, B52	1.33	1.	1043	1043	B14: 5,79	B52: 10,09		
B15, B53	1.66	11	1103	1103	B15:3.99	B53: 10,66		
B16, B54	2.0	14	1123	1123	B16:3,87			
B17	2.5	11	1153	1153	B17: 1.70			
the help	ir about	Demoitrie	432					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JIR	DO	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	28Jana	0923	0953	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	28 DANE	1223	1223	B18: *	B55: _{11.75}
B19	3.5	4,	1253	1253	B19:*	,
B20, B56	4.0	٠,	1323	1323	B50:*	B56:13.14
B21	5.0	٠,	1423	1423	B21: *	
B22, B57	6.0	٠,	1523	1523	B55:*	B57:12.25
B23	7.0	cı	1623	1623	B23:*	
B24	8.0	4.	1723	1723	B24:*	
B25	10.0	и	1923	1927	B25:*	
B26	12.0	٠.	2123	2123	B26:*	
B27, B58	24.0	29 JAN82	0923	0920	B27:*	B58: 12.55
			` // ` /			
* bell	rur assu	ay se	noitivity	433		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JIZ	020	PROTOCOL _DAMD_17085_C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	28 JAN&	0923	0953	IV	AN

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	28Jau86	0800	0800	# 64	#ND	# ok
	0.8	ν,	10828	8 SSC4			
	0.16	4,	0933	0933			
	0.25	9/	0938	0938	#56		
	0.33	lŋ .	0943	0943			
	0.42	ι_i	0948	0948			
	0.50	l _k	0953	0953	#60	# ND	#20
	0.58	i,	0958	0958			
	0.66	h	1003	1003			
	0.75	e i	1008	1008	#64		
	0.83	ι,	1013	1013			
	0.92	l,	1018	1018	-		
	1.0	٤ (1023	1023	#60	#ND	
	1.33	e _k	10 43	1043	# 60		
	1.66	۲,	1103	1103	#60		
	2.0	1-	1123	1123		4 20	
	2.5	4	1153	1153			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	J T R	020	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	28 JANSG	0923	0953	IV	28

DOSAGE (total) 1.32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	25 DANEL	1223	1223	# 72		
	3.5	٠ ٨	1253	1253			
	4.0	e,	1323	1323	# 68	# ~ 5	
	5.0	e _t	1423	1423			
	6.0	٠,	1523	1523	# 72	# ~5	
	7.0	4	1623	1623			
	8.0	4	1723	1723			
	10.0	ų	1923	1923			
	12.0	4.	2123	2123			
	24.0	29 JAN8	0923	0920	#80	# 20	#0K

INVESTIGATOR'S NAME	PT INITIAL PT	# COMPOUND Pyridostigmine
Lietman	JIR D	PROTOCOL <u>DAMD 17-85-C-5133-02</u>

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	31JAN82	0814	0814	PO	NA

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B28, B59	0	31 JAN86	0814	080	B28: *	B59: 12.66
B29, B60	0.25	٠,	0829	0829	B29:*	B60: 12.69
B30, B61	0.50	`.	0844	0844	B30:5,62	B61: 11.82
B31,B62	0.75	Ł i	0859	0859	B31:8,17	B62: 11.24
B32, B63	1.0	٠,	0914	0914	вз2: 6.30	B63: 10.81
B33, B64	1.33	41	0934	0934	взз:9.%	B64: 10.47
B34, B65	1.66	1	0954	0954	B34: 14.0	B65: 10.03
B35, B66	2.0	41	1014	1014	B35:13.6	B66: 9,48
B36	2.5	6	1044	1044	B36:10.3	
B37, B67	3.0	lı .	1114	1114	B37:/2./	B67: 8,94
B38	3.5	41	1144	1144	B38:12.1	
B39, B68	4.0	ι,	1214	1214	B39:8,14	B68: 9,73
B40	5.0	4,	1314	1314	B40:5,45	
B41, B69	6.0	ι,	1414	1414	B41:4.07	B69:10.69
B42	7.0	1,	1514	15/4	B42:5.90	
B43	8.0	4-	1614	1614	B43: 556	B69A://.73
B44	10.0	١,		1814	B44:4.58	11.49

* below assay sensitivity 436

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	77 Q F M L	D20	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	38UACIE	0814	0814	PO	ИА

DOSAGE (total) 16mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOD NG/ML	RBC ACHe uM/ml/mir
B45	12.0	31JAN86	2014 OSH LA	2014	B45: 1,58	
B46, B70	24.0	OIFeb86	0814	0814	B46: *	B70:12.51
					.	
	,					
	•					
			·			
A 1			1			
* Illou	vassay	Sensi	tuly			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	JIL	2 25	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	38 UAC 15	0814	0814	PO	NA

DOSAGE (total) 16mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	31JAU86	0814	0805	#62	#~D	# o L
	0.25	. 4	୍ଥ୍ୟ	0829	* 58		-
	0.50	£ ą	0844	0844	# 56	4 70	# ok
	0.75	s (0859	0859	#54		
	1.0	4.	0914	0914	#56	@ #	
	1.33	``	0934	0934	#5L		
	1.66	4,	0954	0954	# 60		
	2.00	4.8	1014	1014	#56	# 20)	
	2.5	lq .	1044	1044			
	3.0	44	1114	1114	#56		
	3.5	4.	1144	1144			
	4.0	b _g	1214	1214	#60	# NO	
	5.0	ı,	1314	1314			
	6.0	ε,	1414		# 68	* ND	
	7.0	\	1514	1514			
	8.0	λ.	1614	1614			
	10.0	*1	1814	1814			

INVESTIGATOR'S NAME	PT INITIAL PT	# COMPOUND Pyridostigmine
Lietman	JIL D	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	31JAN86	0814	0814	PO	NA

DOSAGE (total) 16mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	31JAL86	2014	2014	24 #		
	24.0	015086	0814	0814	#62	C1 #	ok#
	-						
	19	*					
		-					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	DIR. FM L	120	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hoghen Hospital

Screen Predrug

Study

TEST:NORMAL	24 JAN& ddmmmyy		28 JANS6 ddmmmyy		OIFeb86 ddmmmyy	Date
NA:135-148 MEQ/L	144	14/144A		140	146	
K:3.5-5.0 MEQ/L	4.3	4.1		4.3	4.0	
CL:96-109 MEQ/L	105	108	97	106	104	
CO2:24-30 MEQ/L	22	19		25	28	
SUN:12-25 MG/DL	21/5	14		18	14	
CREAT:0.4-1.5 MG/DL	1.3	1.2		1./	1. /	
GLU:70-115 MG/DL	76	81		79	65	
T. BILI:0.3-1.2MG/DL	0.4	0.5		0.6	0.6	
D. BILI:0.1-0.4MG/DL	0.0	0.6		0./	0.1	
CA:9.0-11.0 MG/DL	10.3	10.2		9.6	10.2	
PO4:3.0-4.5 MG/DL	5.0	4.3		4.2	4.9	
URIC A:4.2-8.8MG/DL	4.7	5.3	_	5, 3	5.3	
T. PROT:6.0-8.5G/DL	7.4	7.3		7.1	7.0	
ALB.:3.2-5.3 G/DL	5.0	5.0		4.7	4.7	
AST:0-35 IU/L	15	15		10	15	
ALT:0-30 IU/L	8	8		9	9	
ALK PHOS:0-95 IU/L	52	52		44	ND	
CHOL:151-268 MG/DL	217	230		229	218	
LDH:0-200 IU/L	ND	ND		128	102	
CPK:0-160 U/L(male)	ND	ND	85	76	80	

NB: No labo on 319AN86

INVESTIGATOR'S NAME	PT INITIAL		
Lietman	J T R F M L	20	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory July Hopkins Hageta

Screen Predrug

Study

TEST	NORMAL	24JAN& ddmmmyy	27JANSC ddmmmyy	292AVK ddmmmyy	01Feb86 ddmmmyy	ddmmmyy	Date
Hgb	13.9-16.3	14.7	14.7	14.2	14.2		
PCV	41.0-53.0	43.5	43.2	42.2	41.2		
Plt	150-350	264	252		258		
RBC	4.50-5.90	4.89	4.84	4.67	4.59		
WBC	4500-11000	5400	6100	4900	5400		
Bands	2-6%	5	5	4	7		
Polys	31-76%	40	247	54	46		
Eos	1-4%	3	1	4	3		
Bas		0		0	1		
Lymphs	24-44%	46	36	31	38		
Atyp Lym		1	α	0	0		
Monos	2-11%	5	8	7	9		
Other		0	0	0	0		
Retics	0.5-1.5%	KN	ND	ND	1.6		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	عد	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Clinical Pharmacolegy

Screen Predrug

Study

TEST	NORMAL	<u>N.D</u>	20) Jan 85 ddmmmyy			ddmmmyy	Date
Color/Ap			yellas	zelln	ND		
Sp. Gr.			1.019	1.024	1.023		
рH			6.0	6.5	60		
Protein			mel	nei	m est		
Ketones			ree.	0	8		
Occ Bld			ner	~ee /	ree		
Bili.			200	new	2		
RBC			0	0	9		
MBC			0	0	0-1		
Casts			0	0	0		
Epi. Cel			0	0	0		
Crystals			0	0	0		
Bacteria			0	0	9		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
24JAN86	/		
29JANGE	/		
2 Fel 86		V	developed Turave changes
			/

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	JIL	Do	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

1		-		1
ł		1		Į
Į	1	ĸ.		ı
١	1		\	ı
ŧ	-	_	_	ĸ,

	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#	¥	dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
	,	(0-2400)	(0-2400)	Mod	PROB.	Treatment
				Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB. POSS.	 Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M - P F M L	021	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	27JAN 86	Screening laboratory
_	BRUATEDE	History, Physical Exam
0	2 FEB 86	Admission
ع	3 FeB 84 6 FeB 84	P.O.
5	6 FEB 86	IV

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject 12
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Krun July M.D.
Investigator's signature

dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	M P	021	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

			30 JAN MEDICAL	HISTORY	0011
Date	of	evaluat		Examiner	SPELL
Date	of	birth	Hodd mmm yy 3, 14,66 dd mmm yy		Brent 6, Petty U.D.
Age			19 yrs		
Sex			$\overline{\mathcal{M}}$		
Race			B		

	No	Yes	Comments
Allergy	/		
Tobacco Use			1 ppd
Alcohol Use			
Recreational Drug Use			M-Journal, lest 14/31/75
Medications past 2 weeks	/		
Experimental Drug Exposure	/		
Blood or plasma donor			
Prior Surgery	/		
Eye, ear, nose, throat			
Endocrine(diabetes,thyroid)			
C-V (heart murmur, HBP)			
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal			
Genito-urinary			
Musculoskeletal			
Neuropsychiatric	/		
Pesticide/herbicide use	/		
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	M - P F M L	021	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION
Date dd mmm yy

			72 746		
Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
34.3	_80_/min	20/min	124,80	177.0	<u>_72.3</u>

-	GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities					
Head/Neck	/							
EENT			Office, White papelar levan (1) timeller from					
Chest, lungs								
Heart	1							
Abdomen								
Genitalia	N	P						
Rectal	N	0						
Extremities								
Skin		/	Membar bygggignented area- and arm framhet grease breme					
Neurologic			lefthanded					

CHEST X-RAY
Date 02/Feb/86

NORMAL	<	ABNORMAL	Describe	abnormalities:
	,		·	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	M - P F M L	021	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	06Feb86	0 818	0848	IV	NA

Syringer Byride = 41, 828 DOSAGE (total) 1.32 Mg Syringer - gride = 31.77 g DOSAGE (total)

PLASMA CONCENTRATIONS

0112	0 80 Br	bject's	me ight = 10	2.1 Kg		
Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B01,B47	0	06Feb86	N.A.	0814	B01: *	B47: 11.59
B02	0.08	٠,	0823	0823	B02:19,6	
воз	0.16	٠,	0828	0828	воз:16.3	
BO4, B48	0.25	ι,	0833	0833	B04:20,7	B4B: 9.63
B05	0.33	V,	0838	0838	B05:2/,7	
B06	0.42	l e	0843	0843	BOE: 19,0	
B07, B49	0.50	٠,	0848	0848	B07: 22,7	B49:8.21
вов	0.58	4.	085]	0853	BOB:/3,7	
B09	0.66	٠١	0858	0828	B09:10,5	
B10, B50	0.75	14	0903	0903	B10:7.54	B50: 8.99
B11	0.83	41	0908	0908	B11: 7.02	
B12	0.92	li .	0913	0913	B12:5.66	
B13, B51	1.0	Þη	0918	0918	B13:3,86	B51:9,70
B14, B52	1.33	t _e .	0938	0978	B14: 5,84	B52: 10.18
B15, B53	1.66	1.	0958	0958	B15: 2.41	B53: 10,40
B16, B54	2.0	t _e	1018	1018	B16: *	B54: 10, 84
B17	2.5	L	1048	1048	B17:*	

* below assay Sensitivity 447

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M-P FML	02 I	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	06 Feb86	0818	0 848	IV	an

DOSAGE (total) 1.32 mg

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	06Fab 86	1118	1118	B18: *	B55: 10.43
B19	3.5	11)148	1148	B19: *	
B20, B56	4.0	1.	1218	1218	B20: *	B56: 11:12
B21	5.0	h	1318	1318	B21:*	
B22, B57	6.0	Ł	1418	1418	B22:*	B57: 1,50
B23	7.0	l a	1518	1518	B23:*	
B24	8.0	()	1618	1618	B24: *	
B25	10.0	ŀη	1818	1818	B25:*	
B26	12.0	b _k	2018	2018	B26: *	
B27, B58	24.0	075686		0818	B27: *	B58: [1,55]
					ia.	
	E 0.0011					

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M - P	D 1	PROTOCOL _DAMD_17085-C-5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	06 Feb86	0818	0848	IV	NA

DOSAGE (total) 1.32 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	06Feb86	NA	0814	#68	# ND	# 0 人
	0.8	11	0823	0823			
	0.16	*4	0828	0828			
	0.25	*1	0833	0833	# ND		
	0.33	11	0838	0838			
	0.42	¢,	0843	0843			
	0.50	* (0848	0848	#68	# ND	# 02
	0.58	٠,	0853	085]			
	0.66	4	0828	0858			
	0.75	1,	0903	0903	# 6 U		
	0.83	11	0908	2908			
	0.92	h	0913	0913			
	1.0	/i	0918	0918	*64	# ND	
	1.33	n	0938	0978	# 68		
	1.66	n	0958	0958	# 64		
	2.0	45	1018	1018	# 60	# ND	
	2.5	1.	1048	1048			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	M-P FML	21	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	06 Feb86	0818	0848	IV	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	06 Feb 87	1118	1118	#64		
	3.5	. (1148	1148			
	4.0	• 1	1218	1218	# 60	# 20	
	5.0	•1	1318	1318			
	6.0	• (1418	1418	# 64	# ND	
	7.0	4.	1518	1518			
	8.0	1,	1618	1618			
	10.0	4	1818	1818			
	12.0	(2018	2018			
	24.0	on Febso	0818	0818	# 68	2 V #	# 32

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M - T F M L	121	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	03 Fab86	0856	0856	PO	29

Subje	itis wit:	73.1 k		CONCENTRA	*	
Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO3 NG/ML	RBC ACHe uM/m1/min
B28, B59	0	0372686	0810	080	B28: *	B59:11.56
B29, B60	0.25	٤.	0911	0911	B29:1,93	B60: 11.66
B30, B61	0.50	• 4	0926	0926	B30:5.02	B61: //, 00
B31, B62	0.75	ts	0941	0940	вз1:8,58	B62: 10.45
B32, B63	1.0	^	0956	0956	B32:11.5	B63: 9,69
B33, B64	1.33	٠.	1016	1016	B33:9,74	B64: 9.52
B34, B65	1.66	L,	1036	1076	B34:/0,/	B65: 9.24
835, B66	2.0	٠,	1056	1056	B35:10.9	B66: 9.02
B36	2.5	i,	1126	1126	B36:5,56	
B37, B67	3.0	٠, ٠	. 1156	115.6	B37: 7,35	B67: 9.57
B38	3.5	N.	1226	1226	B38:3,76	
B39, B68	4.0	١ر	1256	1256	B39:5,70	B68: 9 98
B40	5.0	• 、	1356	1356	B40:2.33	
B41,B69	6.0	• •	1456	1456	B41:3,90	B69: 10.83
B42	7.0	Α	1556	1556	B42:334	
B43	8.0	ι,	1656	1656	B43: 💥	B69A: ji.02
B44	10.0	4,	1856	1856	B44: *	11.01

* below assay sensitivity

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	M - C	Del	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	03 Feb 81-	0856	8556	PO	24

DOSAGE (total)

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	DE/ML	RBC ACHe uM/ml/mir
B45	12.0	03Fe686	2056	2056	B45:	
B46, B70	24.0		0856	0856	B46: ¥	B70: 11,58
					,	
					•.	
<u>-</u>						
* belay	wassay	1 Dens	tivita			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	F M L	021	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: DRAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	03 Feb86	0856	0856	PO	NA

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	03 7-686	0810	0810	# 74	94 W	# ok
	0.25	. • (0911	0911	# 60		_
	0.50	`.	0926	0926		# ND	# NO
	0.75	1	0941	0940	#72		
	1.0	41	0956	0956	#64	# ~ D	
	1.33	4 5	1016	1016	# 68		
	1.66	~ (1036	1036	#64		
	2.00	Lì	1056	1056	# 68	# 20	
	2.5	۴	1126	1126			
	3.0	4.4	1156	1156	# 68		
	3.5		1226	1226			
	4.0	£ (1256	1256	#72	۵۲ #	
	5.0		1356	1356			
	6.0	4.6	1456	1456	#72	# ND	
	7.0	4, "	1556	1556			
	8.0	- 4,	1656	1656			
	10.0	11	1856	1856			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>M</u> <u>-</u> <u>P</u>	021	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	03 Feb86	0856	0856	PO	24

DOSAGE (total) ______16 wy

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	03F-A56	2036	2026	# ND		
	24.0	04 Feb86	0856	0856		#ND	ok#
							-
	*.						
		ν.					

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL PT #	COMPOUND Pyridostigmine
Lietman	F M - P 2	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

9.9				045-686	06 Feb 526	Date
TEST: NORMAL	ddmmmyy	ddmmmyy	ddmmmyy	ddmmmyy	ddmmmyy	
NA:135-148 MEQ/L	141	141	Xen	139		
K:3.5-5.0 MEQ/L	4.2	4.3		4.6		
CL:96-109 MEQ/L	102	106		106		
CO2:24-30 MEQ/L	28	26		21		•
SUN:12-25 MG/DL	1)	9		14		
CREAT:0.4-1.5 MG/DL	1.0	1.2		1.)		
GLU:70-115 MG/DL	78	119		99		
T.BILI:0.3-1.2MG/DL	1.)	0.8		0.4		
D. BILI:0. 1-0. 4MG/DL	0.1	6.1		0.1		
CA:9.0-11.0 MG/DL	9.8	9.8		9.6		
PO4:3.0-4.5 MG/DL	4.6	3.4		3.8		
URIC A:4.2-8.8MG/DL	3.8	4.7		4.9		, .
T. PROT:6.0-8.5G/DL	7.1	64		7.1		
ALB.:3.2-5.3 G/DL	4.7	4.6		4.4		
AST:0-35 IU/L	16	19		26		
ALT:0-30 IU/L	10	12		14		
ALK PHOS:0-95 IU/L	44	43		46		
CHOL:151-268 MG/DL	122	120		183/		
LDH:0-200 IU/L	NP	ND	186	188	176	
CPK:0-160 U/L(male)	ND	ND	151	141	170	

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>M</u> - P	D2 1	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

07Feb86 NA NA Date TEST: NORMAL ddmmmyy ddmmmyy ddmmmyy ddmmmyy ddmmmyy NA:135-148 MEQ/L 141 4.5 K:3.5-5.0 MEQ/L CL:96-109 MEQ/L 106 CO2:24-30 MEQ/L SUN: 12-25 MG/DL 17 CREAT: 0. 4-1.5 MG/DL 1.0 GLU:70-115 MG/DL 85 T. BILI: 0.3-1.2MG/DL D. BILI: 0. 1-0. 4MG/DL QU 9.9 CA:9.0-11.0 MG/DL PD4:3.0-4.5 MG/DL 4.0 URIC A:4.2-8.8MG/DL 4.0 T. PROT:6.0-8.5G/DL ALB.:3.2-5.3 G/DL 41.7 AST:0-35 IU/L 44 ALT:0-30 IU/L 36 ALK PHOS:0-95 IU/L 45 CHOL: 151-268 MG/DL 148 LDH:0-200 IU/L 173 CPK:0-160 U/L(male)

INVESTIGATOR'S NAME	PT INITIAL PT #	COMPOUND Pyridostigmine
Lietman	m - P Q1	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

TEST	NORMAL	admumyy 30 Jun87	ddmmmyy	04Feb86	OJF-686 ddmmmyy	ddmmmyy	Date
НдЬ	13.9-16.3	15.5	15.410	140	15,2		
PCV	41.0-53.0	45.2	43736	ND	43.5		
Plt	150-350	241	262	ND	268		
RBC	4.50-5.90	5.21	5. 2	20	5.08		
MBC	4500-11000	10500	10700	20	9200		
Bands	2-6%	10	4	ND	7		
Polys	31-76%	57	68	70	48		
Eos	1-4%	3	0	ND	D		3
Bas		0	0	NO	0		
Lymphs	24-44%	25	20	ND	39		
Atyp Lym	*	2	1 .	ND	0		
Monos	2-11%	3	5	ND	6		
Other		0	D	ND	0		
Retics	0.5-1.5%	27	ND	ND	1.0		
	,						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	021	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Long Hosting University

Charical Pharmacology

		Screen	Predrug			Study
TEST	NORMAL	ddmmmyy	OZF686 ddmmmyy	04F288 ddmmmyy		Date
Color/Ap			clear	ND	yellow	
Sp. Gr.			1. 020		1.019	
рH			6.0		60	
Protein			mer		~~08	
Ketones			~01		~~ 6	
Occ Bld			200		29/	
Bili.			ner		~~/	
RBC			. 00		0	
WBC			0		ō	
Casts			0		0	
Epi. Cel		-	ی		٥	
Crystals			.0		Û	
Bacteria			0		0	

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
582FCO	1		
04Feb86	✓		
07Feb86	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	M - P F M L	021	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

ADVERSE EXPERIENCES

check if none occurred

Г	∇A	
ı	VI	
1	Λ I	
Ŀ	$_{I}$	

	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mww yy	dd faram yy	☐ Mild	DEF.	Norie
		(0-2400)	(0-2400)	Mod	PROB. POSS.	Treatment
				Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	 None
		(0-2400)	(0-2400)	Mod	PROB.	Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	972	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	17JAN &	Screening laboratory
	30JMV86	History, Physical Exam
0	2 FEB 86	Admission
2	03 Fers 86	P.O.
5	03 Fers 86 6 Fers 84	IV

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject # 0 2.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

_M. D.

Investigator's signature

INVESTIGATOR'S NAME	PT	INITIAL	PT	#	COMPOUND <u>Pyridostigmin</u> e
Lietman	F	-W M L	1 2	2	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

Date of evaluation of the second of the seco

	No	Yes	Comments
Allergy	/		
Tobacco Use		/	lord
Alcohol Use		1	2 parles a welled
Recreational Drug Use	/		
Medications past 2 weeks			
Experimental Drug Exposure		V	Ciftiraxone enteral, tobra
Blood or plasma donor			V
Prior Surgery	V		
Eye, ear, nose, throat			
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)			
Hepatitis, gastro-intestinal			
Genito-urinary			
Musculoskeletal	/		
Neuropsychiatric			
Pesticide/herbicide use		/	Vad modride + 3 woles ogo
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u>L</u> - (1) F M L	12 12	PROTOCOL DAMD 17-85-C-5133-02

PHYBICAL EXAMINATION
Date CH7507 & G

dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
360c	_ 8_Vmin	20/min	11.81.72_	177	70.7

GENERAL EXAMINATION:						
(check)	Nor.	Abn.	Provide details of abnormalities			
Head/Neck	/					
EENT		/	few lynghod declocenes peterio phangex			
Chest, lungs	/					
Heart						
Abdomen	V					
Genitalia	*	M				
Rectal	N	10				
Extremities			, buite lawatin,			
Skin		V	Scar(2) index frager vertical sear LUQ from bibs accident			
Neurologic			pur allocated (b) (b) (b) (b)			

CHEST X-RAY
Date 30/38N/86

NORMAL	X	ABNORMAL	Describe	abnormalities:
			····	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	2 - W F M L	022	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	06 Feb 86	0825	08.22	IV	NA

Syring - pyrido= 21.41 g DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Subjection wt. = 69.0						
	Schedule Time (hours)		Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B01, B47	0	06Fe686	0810	0810	B01: *	B47:15.48
Bos	0.08	٠,	0830	0830	BO2: 7,80	
B03	0.16	٠,	0835	0835	воз: 4.9	
BO4, B48	0.25	* (0840	0840	B04:22,4	B48:12.2
B05	0.33	٠,	0845	0845	B05:2//8	
B06	0.42	*1	0856	0850	BOE:24,8	
B07, B49	0.50	٧.	0855	0855	B07:20.6	B49: 10.91
BOB	0.58	٠,	0900	0900	BO8://.6	
B09	0.66	• •	0905	0905	B09: //.3	
B10, B50	0.75		2510	0910	B10:8,55	B50: 11.95
B11	0.83	- 1	0915	0915	B11:6.59	
B12	0.92		0920	0920	B12:480	
B13, B51	1.0	()	0925	0925	B13:6.36	B51:12.64
B14, B52	1.33	F i	0945	0945	B14:4.09	B52:13.29
B15, B53	1.66	(1)	1005	1005	B15:3.66	B53:1397
B16, B54	2.0	1.	1025	1025	B16:1,75	B54: 14.11
B17	2.5	١٠	1055	1055	B17:2.76	
+ Ibelow assay Sensitivity 463						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u> </u>	022	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
S	06 Feb 8 5	0825.	0855	IV	NA

DOSAGE (total) 1.32 mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ PNG/M	RBC ACHe uM/ml/mir
B18, B55	3.0	06 Feb 86	1125	1125	B18:1,73	B55:14,77
B19	3.5	٠,	1155	1155	B19: *	
B20, B56	4.0	٠,	1225	1225	B20:*	B56: 15.03
B21	5.0	11	1325	1325	B21: *	
B22, B57	6.0	١,	1425	1425	B22: *	B57: 15.31
B23	7.0	"(,	1525	1525	B23: *	
B24	8.0	٠,	1625	1625	B24: *	
B25	10.0	• \	1825	1825	B25: *	
B26	12.0	t,	2025	2025	B26: *	
B27, B58	24.0	07Feb86	0825	0825	B27: *	B58: 15, 44
* belo	rw awa	y seno	itivity 46	4		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	3 F M L	022	PROTOCOL <u>DAMD 17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	06 Feb 86	0825	0855	IV	NN

DOSAGE (total) 1.32 mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	06Feb 86	0800	0800	# 68	#20	#0/2
	0.8		0830	0835			
	0.16	*1	0835	0835			
	0.25	t (0.840	0840	# 60		
	0.33	• ,	0845	0845			
	0.42	`.	.0850	0856			
	0.50	٠,	0855	0855	#64	GN#	# ~ 0
	0.58	11	0900	0620			
	0.66	4.	0905	0905			
	0.75	١٤	0910	0910	# 64		
	0.83	4,	0915	0915			
	0.92	4	0920	0920			
	1.0	4	0925	0925	#60	# ND	
	1.33	1,	0945	0945	#68		
	1.66	٠,	1005	1005	#64		
	2.0	* *	1025	1025	#64	du #	
	2.5	1,	1055	1055			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmi</u> ne
Lietman	<u>β</u> <u>L</u> <u>ω</u>	022	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	06Fe686	0825	0855	IV	NA

DOSAGE (total) 1,32 mg.

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	06Feb86	1125	1125	# 60		
	3.5	((1155	1155			
	4.0	٠,	1225	1225	# 20	# 10	
	5.0	٠,	1325	1325			
	6.0	11	1425	1425	# 60	# 20	
	7.0	fà	1525	1525			
	8.0	c j	1625	1625			
	10.0	(1	1825	1825			
	12.0	r	2025	2025			
	24.0	07 Feb86	0825	0825	#64	# ND	# 0/2
							-

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	<u> </u>	Ø2	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	03 Fa 686	0859	0859	PO	NA

Syring + 8 yride: 7.65638

Syring - pyride: 5.97471 DOSAGE (total) 16 mg

PLASMA CONCENTRATIONS

Subject's weight = 70.7 kg									
Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/m1/min			
B28, B59	0	03Fe686	0820	೧೯ ೩೦	B28: *	B59: 15.32			
B29, B60	0.25	• (0914	0916	B29:2.8/	B60: 14.75			
B30, B61	0.50	~	0929	0929	B30: 6.13	B61: 13.56			
B31, B62	0.75	ι,	0944	0944	B31:/4/	B62:12.53			
832, B63	1.0	٠,	0959	0959	B32: 11.4	B63: 11.76			
B33, B64	1.33	ı,	1019	1019	взз: 9,23	B64: /1,27			
B34, B65	1.66	• •	1039	1039	B34: /0./	B65: 11.76			
B35, B66	2.0	e (1059	1059	B35: 757	B66: 11.47			
B36	2.5	·	1129	1129	B36:8.17				
B37, B67	3.0	٠,	1159	1159	B37:8,/7	B67: 11.19			
B38	3.5	v ₄	1229	1229	B38: 7.67				
B39, B68	4.0	h	1259	1259	B39:59/	B68: 11.94			
B40	5.0	c (1359	1359	B40: 4.77				
B41, B69	6.0	e Į	1459	1459	B41:356	B69: 13.83			
B42	7.0	٤.	1559	1559	B42: X				
B43	8.0	ιį	1659	1659	B43: 1.93	B69A:14,33			
B44	10.0	h	1859	1859	B44: *	14.71			
at lul	MIT AN	0011	1. 1. 1						

* below assay sensitivity 467

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	<u> </u>	022	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	03 Feb 86	0859	0859	PO	NA

DOSAGE (total) 16mg.

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/MC	RBC ACHe uM/ml/mir
B45	12.0	03 Feb86	2059	2059	B45: ¥	
B46, B70	24.0	04 Feb86	0859	0859	B46: *	B70: 15.53
				4		
						(a)
				-		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	<u> </u>	022	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	03 Feb &	0859	2859	PO	20

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	03 Feb 86	0820	0825	#70	# ~5	#ok
	0.25	. 4	0914	09/6	# 76		-
	0.50	t,	0929	0929	#60	1 ND	#NO
	0.75	4	0944	0944	# 56		
	1.0	· c	0959	0959	* 5 0	@y #	
	1.33	ιį	1019	1019	#64		
	1.66	*,	1039	1039	#60		
	2.00	(t	1059	1059	# 52	CN #	
	2.5	6.	1129	1129			
	3.0	41	1159	1159	# 60		
	3.5	c,	1229	1229			
	4.0	٤,	1259	1259	#64	Q C #	
	5.0	4	1359	1359			
	6.0	1.	1359	1459	* 60	# NO	
	7.0	4,	1559	1559			
	8.0	14	1659	1659			
	10.0		1859	1659 1859 18684			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	- <u> 2</u> F M L	022	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	03 Feb86	0859	0859	PO	NA

DOSAGE (total) 16mg.

PHYBIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	03 Feb86	2059	2059	# 20		
	24.0	04Febsa	i	0859	#N0	*NO	~D#
					1.		

page 12 of 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	FML	022	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopkins Hospital

Screen Predrug

Study

TEST:NORMAL	17 JAN &		03 Feb86 ddmmmyy	04 Febsa ddmmmyy	Objeb86 ddmmmyy	D
NA:135-148 MEQ/L	145	143		139		
K:3.5-5.0 MEQ/L	4.6	4.1		4.2		
CL:96-109 MEQ/L	110	109	+1	105		
CD2:24-30 MEQ/L	24	28		22		
SUN:12-25 MG/DL	11	16		12		
CREAT:0.4-1.5 MG/DL	1./	1.1		1.0		
GLU:70-115 MG/DL	40	117		156		
T.BILI:0.3-1.2MG/DL	0.6	0.9		1.0		
D. BILI:0.1-0.4MG/DL	ND	0.1		0.1		
CA:9.0-11.0 MG/DL	9.6	9.5		9.2		
PO4:3.0-4.5 MG/DL	3.6	ND		3.6		
URIC A:4.2-8.8MG/DL	5.3	5.8		5.0		
T. PROT:6.0-8.5G/DL	7.0	6.8		6.4		
ALB.:3.2-5.3 G/DL	5.1	4.7		4.4		
AST:0-35 IU/L	11	24		ಎಂ		
ALT:0-30 IU/L	3	3		11		
ALK PHOS:0-95 IU/L	40	49		41		
CHOL:151-268 MG/DL	207	180		184		
LDH:0-500 IN/F	20	ND	128	104	119	
CPK:0-160 U/L(male)	ND	ND	3713	2 58	207	

page 12Aof 15

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	22	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

Date

	NA		07Feb86			Da
TEST:NORMAL	ddmmmyy	ddmmmyy	ddmmmyy	ddmmmyy	ddmmmyy	
NA:135-148 MEQ/L			140			
K:3.5-5.0 MEQ/L			4.1			
CL:96-109 MEQ/L			104			
CD2:24-30 MEQ/L			25			
SUN:12-25 MG/DL			13			
CREAT:0.4-1.5 MG/DL			1.0			
GLU:70-115 MG/DL			83			
T.BILI:0.3-1.2MG/DL			1. 1			
D. BILI:0. 1-0. 4MG/DL			RS			
CA:9.0-11.0 MG/DL			9.4			
PO4:3.0-4.5 MG/DL			4.2			
URIC A:4.2-8.8MG/DL			5,2			
T. PROT:6.0-8.5G/DL			6.8			
ALB.:3.2-5.3 G/DL			4.8			
AST:0-35 IU/L			33			
ALT:0-30 IU/L			18			
ALK PHOS:0-95 IU/L			41			
CHOL:151-268 MG/DL			187			
LDH:0-200 IU/L			104			
CPK:0-160 U/L(male)			210			

INVESTIGATOR'S NAME		
Lietman	Z- W 22	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hopkins Hospital

Screen Predrug

Study

TEST	NORMAL	MUAC CI	01 Feb 86 ddmmmyy	04Fd86 ddmmmyy	07F185 ddmmmyy	ddmmmyy	Date
Hgb	13.9-16.3	14.9	13.8	13.9	13.7		
PCV	41.0-53.0	43.5	40.0	40.8	39.5		
Plt	150-350	220	225	205	218		
RBC	4.50-5.90	4,98		4.69	4.48		
MBC	4500-11000	4800	4800	4100	5900		
Bands	2-6%	5	4	1	12		
Polys	31-76%	53	48	41	59		
Eos	1-4%	j	3	2	0		
Bas		0	D	0	0		
Lymphs	24-44%	35	33	46	25		
Atyp Lym		0	7	1	0		
Monos	2-11%	6	9	9	4		
Other	-	O	Ö	0	0		
Retics	0.5-1.5%	119	ND	0,4	0.7		
	,						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	2 - W	022	PROTOCOL DAMD 17-85-C-5133-02

Laboratory Chincol Pharmocole Sty

Screen Predrug

Study

TEST	NORMAL	44	N 0	ddminmyy	04F588 ddmmmyy	ddmmmyy	ddmmmyy	Date
Color/Ap				clear	N.O.	grilly		
Sp. G r.				1.019		1.020		
pН				5.5		5.0		
Protein				mer		(ance)		
Ketones				رمكا		0	-	
Occ Bld				200		~ex		
Bili.				Leix		nay		
RBC				0		0		
MBC				0		0		
Casts				0		6		
Epi. Cel				0		0	·	
Crystals			Ü	0		0		
Bacteria		1	/	D		O		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
30JAN8	V		
30JAN86 04F486 07F488	1		
075486	/		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	F M L	022	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred



ADVERSE	DNSET	CESSATION			ACTION
EVENT	(Date/Time)	Date/Time	(Check)	TEST DRUG	(Check)
#	dd mmm yy	dd mm yy	☐ Mild	DEF.	None
				PROB.	
	(0-2400)	(0-2400)	Mod	POSS.	Treatment
				DEF. NOT	
			Sev	UNKNOWN	Stop test drug
# .					
	dd mmm yy	dd mmm yy	Mild	DEF.	None
-				PROB.	
	(0-2400)	(0-2400)	Mod	POSS.	Treatment
				DEF. NOT	
				DEF. NOT	
			Sev	UNKNOWN	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D M L	027	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	30JAN 86	Screening laboratory
_	06 FEB 86	History, Physical Exam
0	09 FEB 860	Admission
2	10 FEB 84	I.V.
5	13 Fes 84	P.O.

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject #123.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M F M L	D23	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

Date of evaluation

Object

October MEDICAL HISTORY

Date of evaluation

October Medical History

Add mmm yy

Date of birth

Date of birth

October Medical History

Age

Date of evaluation

October Medical History

Brent 6. Petty

print name

October Medical History

Age

October Medical History

Brent 6. Petty

print name

October Medical History

Age

October Medical History

Brent 6. Petty

print name

October Medical History

Age

October Medical History

Brent 6. Petty

print name

October Medical History

Age

October Medical History

Brent 6. Petty

print name

October Medical History

Brent 6. Petty

Date of birth

Brent 6. Petty

Date of birth

October Medical History

Brent 6. Petty

Date of birth

Brent 6. Petty

Date of birth

October Medical History

Brent 6. Petty

Date of birth

Date of birth

October Medical History

Brent 6. Petty

Date of birth

Brent 6. Petty

Date of birth

Date of birth

October Medical History

Brent 6. Petty

Date of birth

Date of birth

October Medical History

Brent 6. Petty

Date of birth

Date of birth

October Medical History

Brent 6. Petty

Date of birth

	No	Yes	Comments
Allergy		1	Tomator - lives; Por- + king
Tobacco Use		/	Tomatoe - lives; Por - kine
Alcohol Use	V		
Recreational Drug Use	/		
Medications past 2 weeks			
Experimental Drug Exposure		V	Pharmalineties, last 1983
Blood or plasma donor		/	ist 2 martie ago
Prior Surgery	/		
Eye, ear, nose, throat			
Endocrine(diabetes,thyroid)	/		
C-V (heart murmur, HBP)	/		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal			
Genito-urinary		1	GC ~10 plans ago, Rid & njection
Musculoskeletal	/		
Neuropsychiatric	/		
Pesticide/herbicide use		/	last exterimeters orient - Typ ago
Other			/~/

	1			COMPOUND <u>Pyridostigmin</u> e
Lietman) F	<u>CM</u>	023	PROTOCOL DAMD 17-85-C-5133-02

PHYSICAL EXAMINATION Date 06/Feb/F0 dd mmm yy

Temperature	Pulse	Respir	Blood	Pressure	Height	(cm)	Wt. (kg)
37.5c	_ 72/min	21/min		0,70	L !	<u>3.0</u>	83.0

GENERAL EXAMINATION:							
(check)	Nor.	Abn.	Provide details of abnormalities				
Head/Neck	/						
EENT	V		Few lymphoid excresones posterior plangue				
Chest, lungs	W		·				
Heart	/		II systolic flow (w), physiologic 5,				
Abdomen							
Genitalia	N						
Rectal	1	0	,				
Extremities	V		Mil seaving anteental forose from Nort/ Levers direttino				
Skin		/	Time resocolor (Dann, low back, aheka, lowchest.				
Neurologic	1		Scarly fream, Ownot				

CHEST X-RAY
Date O6/F36/86

NORMAL &	ABNORMAL	Describe abnormalities:	
		-	

Examiner

Brent F. Pott

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M F M L	623	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	10Feb86	0842	0912	IV	NA

Syringe + pyrido = 43.15 & DOSAGE (total) 1.32 MY

pt. wt = 83.3 kg } recribe to this plasma CONCENTRATIONS

pt. wt = 84.5 kg } det in separate

places in Source document

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO3 NG/ML	RBC ACHe uM/ml/min
B01,B47	0	10 Feb 86	N.A.	0807	B01:*	B47:13.34
B02	0.08		0847	0847	BO2:12.3	
B03	0.16	lį	0823	0852	воз: 8.39	
B04, B48	0.25	.*.	0857	0857	B04:16.7	B48: //. 38
B05	0.33	i ş	0902	0902	B05: 226	
B06	0.42	• .	0907	0907	BOE: 14.9	
B07, B49	0.50	• ,	0912	0912	B07:205	B49:9,45
BOB	0.58		0917	0917	BOB: 12,7	
B09	0.66	/	0922	0922	BO9: //.Ó	
B10, B50	0.75	٨	0927	0927	B10:/48	B50: 9.7/
B11	0.83	4,	0932	0932	B11:9.26	
B12	0.92	1,	0937	2937	B12:7.74	
B13, B51	1.0	-	0942	0942	B13:3.86	B51: 10.82
B14, B52	1.33	h	1002	1002	B14:4.24	B52://.08
B15, B53	1.66	Fi	1022	1022	B15:3,86	B53: //-32
B16, B54	2.0	· ų	1042	1042	B16: ¥	B54: 11.96
B17	2.5	l.	1112	1112	B17: 2.92	

* below assay sensitivity 479

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	D C M F M L	023	PROTOCOL DAMD 17-85-C-5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	10Feb86	0842	0912	IV	NA

DOSAGE (total) 1.32 mg/

PLASMA CONCENTRATIONS

page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	10Feb 86	1142	1142	B18: *	B55:12.47
B19	3.5	٠,	1212	1212	B19: *	
B20, B56	4.0	1,	1242	1242	B50: *	B56: 12.94
B21 .	5.0	6 ₈₀	1342	1342	B21: NR	
B22, B57	6.0	* (1442	1442	BSS: NE	B57: 12.72
B23	7.0	٠,	1542	1542	B23: NR	
B24	8.0	-1	1642	1642	B24: NR	
B25	10.0	44	1842	1842	B25: *	
B26	12.0	4	2042	2042	B26: *	
B27, B58	24.0	11 Feloss	0842	0842	B27: *	B58:/3,50
					_	

* below assay sensitivity 480 NR :- not hun

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C A	023	PROTOCOL <u>DAMD 17085-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	16Feb86	0842	0912	IV	NA

DOSAGE (total) 1.32 MY
PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	1070686	NA	0837	# 54	# ND	#0)_
	0.8		0847	0847			
	0.16	٩	0852	5280			
	0.25	١,	0857	0857	# 60		
	0.33	4	0902	0902			
	0.42	r (0907	0907			
	0.50	•4	0912	0912	# 64	#ND	#ok
	0.58	ъ	0917	0917			
	0.66	٤	0922	0922			
	0.75	ęj.	0927	0927	#60		
	0.83	* 1	0932	0932			
	0.92	t _i	0937	0937			
	1.0	14	0942	0942	# 60	1 nD	
	1.33	*1	1002	1002	#64		
	1.66	6),	1022	1022	#60		
	2.0	à,	1042	1242		# ND	
	2.5	k	11)2	1112			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M	D3	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	10Feb86	0842	0912	IV	NN

DOSAGE (total) 1.32mg

PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	10 Feb 86	1142	1142	#64		
	3.5	٠ ١,	1212	1212			
	4.0	١,	12.42	1242	#64	# 20	
	5.0	V ₄	1342	1342			
	6.0	4	1442	1442	#60	# ND	
	7.0	٩	1542	1542			
	8.0	h	1642	1642			
	10.0	.,	1842	1842			
	12.0	ŋ	2042	2042			
	24.0	11 Feb88	0842	0842	# 60	eu #	# ND
					,		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M F M L	023	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13 Feb 86	0828	0858	PO	NA

ots weight = 82.0 Kg [PYRIDO] Sample Schedule Actual RBC Date Scheduled No. Time ddmmmyy Clock Time Clock Time ACHe NG/ML (0-2400)(0-2400)uM/ml/min (hours) B28: * B28, B59 B59:/3.34 0846 13 Feb86 NA B29:480 | B60: 12.87 B29, B60 0.25 1 0913 0913 B30, B61 0.50 B30: 7.77 B61: 11.83 h 8690 0928 B31, B62 B31:10.9 B62:11.36 0.75 0943 0943 B32: 14.0 B63: 10.60 B32, B63 1.0 0958 0958 B33: 16, / B64: 10.14 B33, B64 1.33 1018 1018 4 B34, B65 1.66 B34:15,3 B65: 9,88 1038 1 1038 B35:/3,9 | B66: 9,94 B35, B66 2.0 4, 1058 1058 2.5 B36: /3,4 **B36** 1128 1128 B37:12,3 B67:9,77 B37, B67 3.0 14 1158 1158 1, B38:/3/ 3.5 **B38** 1228 1228 B39, B68 4.0 B39: 857 B68: 10, 23 1258 1328 B40: 7,46 **B40** 5.0 1358 1358 B41:4,89 B69: 11.32 B41, B69 6.0 1458 1458 ١, B42:4.60 B42 7.0 1558 1 1558 1 B43:5,29 B69A: 11.61 **B43** 8.0 1658 1658 B44: 169 **B44** 1858 10.0 1858 12.29

* below assay sensitivity 483

INVESTIGATOR'S NAM	E PT INITIAL	PT # COMPOUND _Pyridostigmine
Lietman	D C M	PROTOCOL DAMD 17-85-C-5133-

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13 Feb 86	0858	0828	PO	NA

PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDO) NG/ML	RBC ACHe uM/ml/mir
B45	12.0	13Feb86	2058	2058	B45:*	
B46, B70	24.0	14 Feb 86	0828	8280	B46: 2,20	B70:13.21
					•	
						-
* bell	w associ	y ser	Victority	484		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	D C IN F M L	023	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13Feb86	0828	0858	PO	24

DOSAGE (total) 16 MY
PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	13JAN86	27	0849	#60	# 20	# ok
	0.25	ė (0913	0913	# 60		1
	0.50	n	0928	0928	#64	# 20	# ok
	0.75	i,	0943	0943	#68		
	1.0	ካ '	0958	0958	#64	0 N #	
	1.33	6%	1018	1018	# 60		
	1.66	4,	1038	1038	#64		
	2.00	11	1058	1058	#60	4 ND	
	2.5	i i	1128	1128			
	3.0))	1158	1158	#60		
	3.5	4	1228	1228			
	4.0	Ŋ	1258	1258	# ND	# ~)	
	5.0	٨	1358	13 58			
	6.0	''	1458	1458	# ND	# 20	
	7.0	4,	1558	1558			
	8.0	١,٠	1658	1658			
	10.0	1,	1858	1858			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M F M L	023	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

		Start drug admin- istration (0-2400)			Bottle I.D. #
5	13 Feb 86	0828	0858	PO	an

DOSAGE (total) _____ /6 MV
PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	13 Feb 82	2028	2058	#25		
	24.0	14F-686		0858	#68	# ND	ok#
	1						
			19				
					·		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C M	23	PROTOCOL <u>DAMD</u> 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hopeful

Screen Predrug

Study

TEST:NORMAL	30 DANS			11Feb86 ddmmmyy		ddmmmyy	Date
NA:135-148 MEQ/L	146			141	139		
K:3.5-5.0 MEQ/L	4.3			4.3	4.2		
CL:96-109 MEQ/L	107			106	103		
CO2:24-30 MEQ/L	23			24	25		
SUN:12-25 MG/DL	15			16	12		
CREAT: 0.4-1.5 MG/DL	1.1			1.0	1.1		
GLU:70-115 MG/DL	61			89	109		
T.BILI:0.3-1.2MG/DL	8.0			1.3	0.6		
D. BILI:0. 1-0. 4MG/DL	0.0			0.1	2		
CA:9.0-11.0 MG/DL	10.1			9.7	10.0		
PO4:3.0-4.5 MG/DL	4.5			3.7	2.9		
URIC A:4.2-8.8MG/DL	4287.4			6.8	7.0		
T. PROT:6.0-8.5G/DL	7.6			7.3	7.5		
ALB.:3.2-5.3 G/DL	4.7			4.7	4-8		
AST:0-35 IU/L	16			8	16		
ALT:0-30 IU/L	22			9	11		
ALK PHOS:0-95 IU/L	59			60	51		
CHOL:151-268 MG/DL	20			C4	156		
LDH:0-200 IU/L	184			121	84		
CPK:0-160 U/L(male)	ND	(/	76	82		

NB. CPK + LDH not done on days of doning

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	DCM FML	⊃ 3	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hopkurs Hospital

Screen Predrug

Study

								_
TEST	NORMAL	30 JAN8L ddmmmyy			LL Feb &L	14F-186 ddmmmyy	ddmmmyy	Date
Hgb	13.9-16.3	15.2			15.1	15.1		
PCV	41.0-53.0	45.7			45.4	44.8		
Plt	150-350	162			193	176		
RBC	4.50-5.90	4.95			4.93	4.87		
WBC	4500-11000	8200			7600	7300		
Bands	2-6%	7			6	8		
Polys	31-76%	49			50	59		
Eos	1-4%	0			3	0		
Bas		0			0	0		
Lymphs	24-44%	33			31	24		
Atyp Lym		2			Q	0		
Monos	2-11%	9			8	9		
Other		D			0	0		
Retics	0.5-1.5%	ND	J	/	1.5	1.4		
			galate					

INVESTIGATOR'S NAME	PT	INI	TIAL	PT	#	COMPOUND	Pyridostigmine
Lietman	D F	C_ M	713	0	}	PROTOCOL	DAMD 17-85-C-5133-02

URINALYBIS VALUES
Laboratory Johns Hopkins Harmacology
Clinical Charmacology

Scre	eri	Pre	drug

Study

TEST	NORMAL	06Feb86 ddmmmyy	qqw _77	D,	11Fe	<u>ь 86</u> mmyy	14F486 ddmmmyy	ddmmmyy	Date
Color/Ap		wellow			2	. P.	willow		
Sp. Gr.		1.020					1,029		
pН		6.0					6.0		
Protein		nex	•				6.0 (trace) rey		
Ketones		ner					ner		
Occ Bld		ner					rex		
Bili.		rev					nex		
RBC		0					0		
WBC		0					0		
Casts		0					0		
Epi. Cel		0					0		190
Crystals		0					0		
Bacteria		0	١		(/	0		

ELECTROCARDIOGRAM

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
06F.686	~		
11 Feb 86	v	,	
14 Feb 86	1		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	D C K	023	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

ADVERSE EXPERIENCES

check if none occurred

	OVERSE EVENT	ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)
#		dd mmm yy	dd mmm yy	Mild	DEF.	None
		(0-2400)	(0-2400)	Wod	PROB. POSS.	Treatment
			,	Sev	DEF. NOT	Stop test drug
#		dd mmm yy	dd mmm yy	Mild	DEF.	None
		(0-2400)	(0-2400)	Mod	PROB.	 Treatment
				Sev	DEF. NOT	Stop test drug

COMMENTS (Indicate # and event)

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R M B	024	PROTOCOL DAMD 17-85-C-5133-02

PROTOCOL

Study Day	Date ddmmmyy	Procedures
_	03FEB 86	Screening laboratory
-	06 FEB 86	History, Physical Exam
0	09 FeB 86	Admission
2	10 FEB 84	I.V.
5	13FEB86	I.V. P.O.
		·

I certify that:

- 1) I have carefully examined all entries on the data collection forms, consisting of 15 pages, for subject # 24.
- 2) All information entered onto the data collection forms for this subject, to the best of my knowledge, is correct.

Investigator's signature

24,0ct,80 dd/mmm/yy

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	RMB	024	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

MEDICAL HISTORY

Date	of	evaluation	<u>Feb</u>	

Examiner

Brent & Petty
print name

Date of birth

24/Feb/53 dd mmm yy

Age

_32_yrs

Sex

Race

B

	No	Yes	Comments
Allergy			
Tobacco Use	/		
Alcohol Use		/	Rauly
Recreational Drug Use		/	last tried craine, 1/2 yes ago
Medications past 2 weeks	1		
Experimental Drug Exposure		V	Phamalintie at ye ago
Blood or plasma donor		/	last 1978
Prior Surgery			
Eye, ear, nose, throat	/		
Endocrine(diabetes, thyroid)	/		
C-V (heart murmur, HBP)	V		
Pulmonary (cough, asthma)	/		
Hepatitis, gastro-intestinal	/		
Genito-urinary			GC 1977, AH RN
Musculoskeletal	V		
Neuropsychiatric			
Pesticide/herbicide use	/		V. pets, us insertrides
Other			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB	024	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

PHYSICAL EXAMINATION Date 06/Feb/86 dd mmm yy

Temperature	Pulse	Respir	Blood Pressure	Height (cm)	Wt. (kg)
~~~ c	_ND_/min	ND/min	-ND	ND.	ND.

	GENERAL EXAMINATION:									
(check)	Nor.	Abn.	Provide details of abnormalities							
Head/Neck										
EENT			Slightly red Treather pellaco, few lapsquard accumines							
Chest, lungs										
Heart										
Abdomen	/									
Genitalia	N	D								
Rectal	Ą									
Extremities	/									
Skin		/	Modest acue in ban glot face buthing (P) our-							
Neurologic	V		left handed (Bushed things - buttons							

CHEST X-RAY
Date 06/FEB/86

NORMAL	×	ABNORMAL	Describe	abnormalities:	

Examiner

print name

INVESTIGATOR'S NAME	PT INITIAL		COMPOUND Pyridostigmine
Lietman	RMB FML	024	PROTOCOL DAMD 17-85-C-5133-02

#### STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
Q	10 Feb86	0846	0916	IV	NA

Syringe + gyids = # 43.16 g DOSAGE (total) 1.32 mg.

# PLASMA CONCENTRATIONS

pt's w	pt's weight = 68.0 kg								
Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOS NG/ML	RBC ACHe uM/ml/min			
B01, B47	0	10Feb88	NA	0810	B01:*	B47:10.51			
BOS	0.08	٠,	0851	0855	B02:9,91				
B03	0.16	٠.	0856	0856	B03:11.2				
B04, B48	0.25	• •	0901	0901	B04:/3.0	B4B: 9,02			
B05	0.33	•,	0906	0906	B05: 19.2	,			
B06	0.42	••	0911	0911	BOE:22,3				
B07, B49	0.50	٤,	0916	0916	B07:20\$	B49: 7.29			
BOB	0.58	8	0921	0921	BOB:/7,0				
B09	0.66	t,	0926	0926	во9:12.4				
B10, B50	0.75	C ₄	0931	0931	B10:9.04	B50: 8.07			
B11	0.83	l a	0936	0936	B11: 6.98				
B12	0.92	1,	0941	0941	B12: 6.77				
B13, B51	1.0	4	0946	0946	B13:5.37	B51: 8,51			
B14, B52	1.33	1,	1006	1006	B14:3.19	B52: 8,87			
B15, B53	1.66	1.	1026	1026	B15:3,77	B53: 9,30			
B16, B54	2.0	(i	1046	1046	B16: 4,40	B54: 9.22			
B17	2.5	6,	1/16	1116	B17:4,64				
* helai	v assau	s sensi	twity 494						

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB FML	24	PROTOCOL DAMD 17-85-C-5133-02

# STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	10 Fe 686	0846	0916	IV	NA

DOSAGE (total) 1,32 mg.

#### PLASMA CONCENTRATIONS

#### page 2

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	CPYRIDOJ NG/ML	RBC ACHe uM/ml/min
B18, B55	3.0	10Feb86	1146	1146	B18:1.57	B55: 9.81
B19	3.5	٠,	1216	1218 20	B19:/.93	
B20, B56	4.0	·	1246	1246	B20: *	B56: 10,22
B21	5. 0	٠,	1346	1346	B21: 🗶	
B22, B57	6.0	4,	1446	1446	B55:*	B57:10,25
B23	7.0	l. į	1546	1546	B23:*	
B24	8.0	• (	1646	1646	B24: *	
B25	10.0	٠,	1846	1846	B25: *	
B26	12.0	4	2046	2046	B26: *	
B27, B58	24.0	11 Feb 86	0846	0846	B27: *	B58: 10,88
			1			

* below assay sensitivity 495

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB	024	PROTOCOL _DAMD_17085_C-5133-02

# STUDY: IV PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
2	10 Feb 86	0846	0916	IV	NA

DOSAGE (total) 1.32mg.

# PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	O	10 Feb 86	NA	0835	# 72	# 20	# o/c
	0.8		0851	0851			
	0.16	٠,	0856	0856			
	0.25	" (	0901	0908	# 78		
	0.33	**	0906	0906	,		
	0.42	ε,	0911	0911			
	0.50	N	0916	0.916	#ŋ2	# NO	#ot
	0.58	٠,	0921	0921			
	0.66	٠,	0926	0926			
	0.75	٠,	0931	0931	#86		
	0.83	(,	0936	0936			
	0.92	٠,	0941	094,			
	1.0	V _X	0946	0946	*80	# 20	
	1.33	٠,	. 1006	1006	#80		
	1.66	٤,	1026	1026	# 80		
	2.0	• `	1046	1046	#80	# 20	
	2.5	ft	11/6	1116			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmine</u>
Lietman	R M R	0a4	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

# STUDY: IV PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
2	105.686	0846	0916	IV	NA

DOSAGE (total) 1.32 mg.

# PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	3.0	10Feb 86	1146	1146	#72		
	3.5		1216	1216			
	4.0	e _t	1246	1246	# NO	# 20	
	5.0	١,	1346	1346			
	6.0		1446	1446	@U#	# N D	
	7.0	4.(	1546	1546			
	8.0	l _k	1646	1646			
	10.0	4,	1846	1846			
	12.0	~,	2046	2046			
	24.0	11 Feb 86		0842	# 80	2N#	#0 K
							*
<u> </u>							
			4				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	R M B	024	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

#### STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13 Fe 682	0903	0903	PO	

# PLASMA CONCENTRATIONS

14 Feb 86	tw 2 tg	= 68.6	)			
Sample No.	Schedule Time (hours)	<b>Date</b> ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/min
B28, B59	0	13 Feb86	NA	0849	B28: *	B59:10.60
B29, B60	0.25	٠,	0918	0918	B29:2.00	B60: 10.67
B30, B61	0.50	• •	0933	0933	B30:10.6	B61: 9,07
B31,B62	0.75	• •	0948	0948	B31: 10.9	B62: 8,75
B32, B63	1.0	٠,	1003	1003	B32: 11.6	B63:7,94
B33, B64	1.33	4,	1023	1023	0,0\:EE8	B64: 2.95
B34, B65	1.66	-1	1043	1043	B34:/3,0	B65:7,46
B35, B66	2.0	-1	1103	1103	B35:/0,6	B66:7.28
B36	2.5	ci,	1133	1133	B36: 9.38	
B37, B67	3.0	مر	1203	1203	B37:9.98	B67: 7.81
B38	3.5	e,	. 1233	1233	B38:9.04	
B39, B68	4.0	4,	1303	1303	вз9:9,45	B68: 8,33
B40	5.0	* (	1403	1403	B40:4.62	
B41,B69	6.0	" (	1503	1503	B41: 3,41	B69: 9.31
B42	7.0	۲.	1603	1603	B42:2,63	
B43	8.0	٠,	1703	1703	B43: 2,/4	B69A: 9,73
B44	10.0	΄(	1903	1903	B44: 1,84	9.43
* held	v assay	y sens	itivity 498			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND _Pyridostigmine
Lietman	R M B	24	PROTOCOL <u>DAMD 17-85-</u> C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13 Feb 86	0903	0903	PO	NA

DOSAGE (total) 16mg.

# PLASMA CONCENTRATIONS

Sample No.	Schedule Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	NG/ML	RBC ACHe uM/ml/mir
B45	12.0	13Feb86	2103	2103	B45: *	
B46, B70	24.0	14 Fe686	0903	0903	B46:*	B70: 1021
				-		
					•	
* held	w MM	DUL DE	nsitivity			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND <u>Pyridostigmin</u> e
Lietman	R M B F M L	024	PROTOCOL <u>DAMD 17-85-C-</u> 5133-02

STUDY: ORAL PYRIDOSTIGMINE

Study	Date	Start drug admin-	End drug admin-	Route	Bottle
Day	ddmmmyy	istration (0-2400)	istration (0-2400)		I.D. #
5	13 Felso	0903	0903	PO	NA

DOSAGE (total) /6mg.

# PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	0	13 Feb86	0847	0847	<b>*</b> 80	#ND	# ok
	0.25	. 4	0918	0918	# 80		-
	0.50	**	0933	0933	*84	# ~0	# ok
	0.75	.,	0948	0948	<b>*</b> 80		
	1.0	C _C	1003	1003	#76	# ND	
	1.33	r _t	1023	1023	#76		
	1.66	£4	1043	1043	# 80		
	2.00	4 4	1103	1103	#80	24 NO	
	2.5	£ ę	1133	1133			
	3.0	14	1203	1203	08#		
	3.5	e,	1233	1233			, .
	4.0	• 4	1303	1303	# 20	#ND	
	5.0	٠,	1403	1403			
	6.0	ار	1503	1503	en#	2 U W	
	7.0	در	1603	1603			
	8.0	tç	1703	1703			
	10.0	4	1903	1903			

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB FML	024	PROTOCOL DAMD 17-85-C-5133-02

STUDY: ORAL PYRIDOSTIGMINE

			End drug admin- istration (0-2400)		Bottle I.D. #
5	13 Feb 84	0903	0903	PO	29

# PHYSIOLOGIC VARIABLES

Sample No.	Scheduled Time (hours)	Date ddmmmyy	Scheduled Clock Time (0-2400)	Actual Clock Time (0-2400)	Pulse Rate bpm	Hand Grip L/R	Coord Test
	12.0	13Feb86	2103	2103	# ND		
	24.0	145-656	1	0903	#88	# ND	a)<#
	-	-					
	14						
			,				

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB	024	PROTOCOL DAMD 17-85-C-5133-02

CHEMISTRY VALUES
Laboratory Johns Hospital

Predrug Screen Study 11 Feb 86 14 Feb86 Date 03 Feb 86 ddmmmyy ddmmmyy TEST: NORMAL ddmmmyy ddmmmyy ddmmmyy NA:135-148 MEQ/L 142 144 141 K:3.5-5.0 MEQ/L 4.3 4.7 4.1 CL:96-109 MEQ/L 107 104 105 CO2:24-30 MEQ/L 24 24 29 SUN: 12-25 MG/DL 11 16 10 CREAT: 0.4-1.5 MG/DL 0.8 1.0 1.1 GLU:70-115 MG/DL 61 77 96 T. BILI: 0. 3-1. 2MG/DL 0.4 N.D. 0.5 D. BILI: 0. 1-0. 4MG/DL ND 0.0 0.0 CA:9.0-11.0 MG/DL 9.6 N.D. 9.9 PO4:3.0-4.5 MG/DL 3.6 N.D. 4.2 URIC A:4.2-8.8MG/DL N.D. 5.6 5.5 T. PROT:6.0-8.5G/DL N.D 7.2 7.7 ALB.:3.2-5.3 G/DL N.P. 4.7 4.9 AST:0-35 IU/L 15 15 12 ALT:0-30 IU/L 12 11 10 ALK PHOS:0-95 IU/L 35 25-37 CHOL: 151-268 MG/DL 141 150 ND LDH:0-200 IU/L 103 ND 120

N.B.: CPK+LDH not done on days of drug daing

ND

CPK:0-160 U/L(male)

77

75

INVESTIGATOR'S NAME		
Lietman	IL M B D4	PROTOCOL DAMD 17-85-C-5133-02

HEMATOLOGY VALUES
Laboratory Johns Hospital

Screen Predrug

Study

TEST	NORMAL	03 Feb 81 ddmmmyy		.D.		14 Feb 86 ddmmmyy	ddmmmyy	Date
НдЬ	13.9-16.3	13.7			14-1	14.5		
PCV	41.0-53.0	41.5			42.2	43.3		
Plt	150-350	250			201	193		
RBC	4.50-5.90	5.03			5.18	5.28		
MBC	4500-11000	6100			6100	9100		
Bands	2-6%	9			2	6		
Polys	31-76%	64			48	66		
Eos	1-4%	0			4	3		
Bas		O			1	1		
Lymphs	24-44%	27			36	14		
Atyp Lym		0			1	2		
Monos	2-11%	0			8	8		
Other	-	0			0	0		
Retics	0.5-1.5%	ND	$\rightarrow$		0.8	0.8		

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RMB	294	PROTOCOL DAMD 17-85-C-5133-02

URINALYSIS VALUES
Laboratory Johns Hospital
Clinical Pharmaculogy

		Screen	Predi	^ug				Study
TEST	NORMAL	06 F-186 ddmmmyy						 Date
Color/Ap		zellow			N,	<i>D</i> .	yellas	
Sp. Gr.		1.018					1.008	
рН		6.0					6.0	
Protein		(tree)	•				ne.	
Ketones		new					rey	 ,
Occ Bld		need					res	
Bili.		nev					~av	
RBC		0					0	
WBC		B					0	
Casts		0					0	
Epi. Cel		0					0	
Crystals		0		*			0	
Bacteria		O	]	/			0	

# **ELECTROCARDIOGRAM**

Date ddmmmyy	NORMAL check	ABNORMAL check	Describe abnormalities
06 Fels6	✓		
11 Feb86			
14 Feb 86	Sys.		est atrial abnormality, questionable lead
	,		placement

INVESTIGATOR'S NAME	PT INITIAL	PT #	COMPOUND Pyridostigmine
Lietman	RM B FM L	024	PROTOCOL <u>DAMD 17-85-C</u> -5133-02

# ADVERSE EXPERIENCES

check if none occurred

_	1	1
T	$\mathcal{I}$	7
ı	12	K I
1	/	
1	1	

ADVERSE EVENT		ONSET (Date/Time)	CESSATION Date/Time		REL'N TO TEST DRUG	ACTION (Check)	
#		dd mmm yy	dd mmm yy	Mild	DEF.	None	
		(0-2400)	(0-2400)	Mod	PROB.	 Treatment	
		e e e e e e e e e e e e e e e e e e e		Sev	DEF. NOT	Stop test drug	
#		dd mmm yy	dd mmm yy	☐ Mild	DEF.	None	
		(0-2400)	(0-2400)	☐ Mod	PROB.  POSS.	 Treatment	
				Sev	DEF. NOT	Stop test drug	

COMMENTS (Indicate # and event)